

# UNIVERSIDAD DE LA FRONTERA

Facultad de Ingeniería, Ciencias y Administración  
Programa de Doctorado en Ciencias de Recursos Naturales



## **MANGANESE (Mn) EXCESS IN BLUEBERRY CULTIVARS (*Vaccinium corymbosum* L.): POSSIBLE MECHANISMS OF TOXICITY AND RESISTANCE**

---

TESIS PARA OPTAR AL GRADO ACADÉMICO  
DE DOCTOR EN CIENCIAS  
DE RECURSOS NATURALES

---

**RAYEN XIMENA MILLALEO MILLALEO**

**TEMUCO – CHILE  
2014**

# UNIVERSIDAD DE LA FRONTERA

Facultad de Ingeniería, Ciencias y Administración  
Programa de Doctorado en Ciencias de Recursos Naturales



## **MANGANESE (Mn) EXCESS IN BLUEBERRY CULTIVARS (*Vaccinium corymbosum* L.): POSSIBLE MECHANISMS OF TOXICITY AND RESISTANCE**

**Doctoral Thesis  
In Fulfillment of the  
Requirements for the Degree  
Doctor of Sciences in Natural Resources  
By**

**RAYEN XIMENA MILLALEO MILLALEO**

**TEMUCO – CHILE  
2014**

**MANGANESE (Mn) EXCESS IN BLUEBERRY CULTIVARS (*Vaccinium corymbosum* L.): POSSIBLE MECHANISMS OF TOXICITY AND RESISTANCE**

Esta tesis fue realizada bajo la supervisión del director de Tesis Dra. MIREN RITA ALBERDI LAG, perteneciente al Departamento de Ciencias Química y Recursos Naturales de la Universidad de La Frontera y es presentada para su revisión por los miembros de la comisión examinadora.

**RAYEN XIMENA MILLALEO MILLALEO**

.....  
DIRECTOR PROGRAMA  
DOCTORADO EN CIENCIAS DE  
RECURSOS NATURALES

.....  
Dra. MIREN ALBERDI

.....  
Dr. LUIS CORCUERA

.....  
DIRECCIÓN DE POSTGRADO  
UNIVERSIDAD DE LA FRONTERA

.....  
Dra. LILIANA CARDEMIL

.....  
Dra. MARIA DE LA LUZ MORA

.....  
Dr. FERNANDO BORIE

.....  
Dra MARJORIE REYES

A Dios, mi familia, mis amigos.

## *Agradecimientos/Acknowledgements*

Muchos agradecimientos debo realizar. En primer lugar, agradezco profundamente a Dios por darme una familia, ánimo, salud y personas queridas que siempre están alrededor. Gracias por ayudarme en este desafío.

Mis agradecimientos van también a mi querida familia, que siempre me apoyaron en esta etapa. A mis padres, Ruth y Leo, mis pilares y en quienes siempre recibo su apoyo, consejos, amor y comprensión y veo su esfuerzo y amor hacia la familia. Son mi apoyo emocional y espiritual. A mis queridos hermanos Luis y Mauricio, que con su alegría, consejos y amistad me han ayudado siempre. A mis tios, Elena, Claudio y Lautaro, donde también recibo su apoyo y cariño.

Agradezco profundamente a mis tutores de tesis, Dra. Miren Alberdi y Dra Marjorie Reyes, quienes me brindaron su enorme apoyo, paciencia, confianza, sus consejos invaluable y en momentos complicados, me ayudaron y orientaron para desarrollar esta tesis doctoral. Aprecio mucho sus valiosos y sabios aportes y criticas constructivas hacia el trabajo desarrollado. A Marjorie, no sólo agradezco su apoyo académico, si no también su cariño y aprecio que es recíproco a ella y su linda familia.

Mis agradecimientos van también a la Dra. María de la Luz Mora por sus valiosos consejos, su experiencia y su constante apoyo en todos los momentos, que ha contribuido a una buena orientación para mejorar esta tesis. También agradezco al Director del Programa en Ciencias de Recursos Naturales de la Universidad de La Frontera, Dr. Francisco Matus porque también me ayudado y apoyado en esta última etapa de la tesis.

Agradezco al grupo de Fisiología Vegetal de la Universidad de La Frontera, a Ximena Garrido, Jessica Yañez, Mariela Mora y Karin Berndt que junto con su ayuda en el laboratorio, colocando en marcha diversas metodologías y experimentos en ardua labor, fueron importantes para realizar la tesis, sin las cuales se hubiera hecho mucho más difícil. También me brindaron su amistad, apoyo y alegría en momentos adversos. Muchas gracias por todos esos momentos de alegría, conversaciones y humor!!!

Mis agradecimientos van también al querido grupo técnico del Laboratorio de Suelo de la Universidad de La Frontera, pues muchas veces su ayuda fue imprescindible. Además, su calidez humana y alegría me animaban para continuar trabajando en los experimentos. Gracias Juanita, Olguita, Evelyn, Ceci, Jaque, Lore, Fanny, Ale, Dani, Marce... me ayudaron y su alegría fue un motor para seguir adelante. Doy gracias también a la Sra Brigida Sobarzo que me ayudó a analizar mis muestras en el Laboratorio de Suelo.

Quisiera agradecer en especial a mis estimadas amigas: “las Ediths”: Edith Alarcón, Edith Cantero, a Elizabeth Ulloa, Ligia Pinilla, gracias por su inmenso apoyo, por sus palabras de

ánimo constantemente, y por esas largas conversaciones que sirvieron para trabajar con alegría y entusiasmo.

Quisiera agradecer también a la Dra. Alejandra Ribera, que siempre con su disposición, no sólo me ayudó con los escritos y críticas muy buenas para mejorar mi tesis, sino me brindó su amistad y alegría. Muchac gracias Ale!.

También, quisiera agradecer a la Universidad Western Ontario de Canadá, Departamento de Biología y en especial especial al grupo de trabajo del Dr. Norman Hüner. Profundamente agradezco al Dr. Hüner por darme la oportunidad de estar en su excelente grupo de trabajo y sus excelentes colaboradores: el Dr. Alex Ivanov y Dra. Marianna Krol, que siempre me apoyaron con sus consejos, su confianza y su calidez humana. Doy infinitamente las gracias también a la Dra. Alejandra Zuñiga, a quien conocí allá y me brindó su apoyo, consejos y generosa amistad. Allá conocí también a mi “abuela argentina” Norma Nuñez, que me brindó su amor y atención. Su cariño y atención emocional y espiritual fue valiosísimo para mí durante mi estadía en Canadá.

Una profunda gratitud doy también a la Dra Marysol Alvear y mi compañera y amiga Paula Aguilera, porque ambas me ayudaron en una parte esencial de mi tesis, al enseñarme técnicas relacionadas a la extracción de acidos organicos en mis plantas. Y no sólo eso, me brindaron consejos útiles importantes y sobre todo su cariño y amistad.

Agradezco a mis amigos y compañeros del doctorado: Loreto Manosalva, Andrea Silva, Elizabeth Garrido, Cecilia Paredes, Paula Aguilera, Karen Manquian, Sebastián Meier, Marcelo Panichini, Raúl Orrego, Emilio Jorquera, Claudio Inostroza, Edgardo Cruces y Oscar Martínez. En forma muy, muy, muy especial quiero agradecer a mi gran amiga Pilar Ulloa por su incondicional amistad, ánimo, valentía y apoyo en todos los momentos. Sus experiencias y consejos siempre son útiles en todo los aspectos. Gracias Pili!!!

Por último, agradezco a la Beca de Apoyo de Tesis Conicyt, Becas de Pasantía en el extranjero Becas-Chile y Beca de Asistencia a Reuniones Científicas Conicyt. Agradezco también el apoyo económico del Programa de Postgrado y del Doctorado en Ciencias de Recursos Naturales de la Universidad de La Frontera por financiar muchas veces trabajos científicos a diferentes congresos nacionales e internacionales.

## Thesis outline

Acid soils, such as Ultisol and Andisol of southern Chile, have volcanic ash origin. The main characteristics of these soils are the high organic matter (OM) content, low availability of essential elements for plants such as nitrogen (N), phosphorus (P), calcium (Ca), magnesium (Mg) and micronutrients (e.g. molybdenum, Mo), and pH values lower than 5.5. This last characteristic is associated with high available contents of aluminum (Al) and manganese (Mn), which can be toxic for crops reducing their productivity. Mn is influenced by soil redox conditions, where the reduced form of this metal,  $Mn^{2+}$  is the available form for plants.

Manganese is an essential micronutrient for plants. In plant cells, it participates in various metabolic cell processes, such as photosynthetic pathways, enzyme activation, and diverse metabolites syntheses. Nevertheless, Mn excess in acid soils causes various alterations at physiological and biochemical level in plants according to its toxicity. Thus, these toxic effects in plants alter the photosynthetic capacity, stimulating the reactive oxygen species (ROS). ROS are highly harmful to biological membranes, which can cause alterations including to the cell death. To counteract this damage, plant cells have developed several mechanisms that remove ROS.

In Chile, highbush blueberry (*Vaccinium corymbosum* L.) is grown in the southern regions where is well adapted to acid soils (pH ranges from 4.0 to 5.5), with high levels of organic matter (OM), even in flooded conditions. Fruits are a good source of natural antioxidants, considered beneficial for human health.

The general aim of this thesis is to evaluate the Mn toxicity effect on physiological and biochemical properties of photosynthetic apparatus, as well as to analyze its resistance mechanisms to Mn excess in highbush blueberry cultivars. Additionally, the effect of Mn excess on photosynthetic performance in *Arabidopsis thaliana* as a model plant under Mn excess was studied.

In the first Chapter a General Introduction is presented, where an overview about the Mn excess in plants is showed. In Chapter 2, an overview on Mn transport, accumulation and resistance mechanism are discussed.

In Chapter 3 biochemical and physiological differential responses of three cultivars of blueberry (Legacy, Brigitta and Bluegold) subjected to increasing Mn concentrations in nutrient solutions were analyzed. The results showed no evidence of visual symptoms, but significant differences in biomass and photosynthetic parameters were found. Significant reductions in aerial biomass (33%) in Bluegold with the highest Mn treatment (1000  $\mu\text{M}$ ) compared with the control were observed. Brigitta and Legacy showed the best growth of the aerial biomass remaining stable under Mn treatments. In contrast, root growth was not significantly different in all cultivars subjected to Mn excess. Moreover, some mechanisms of tolerance as the antioxidant system and the exudation of organic compounds indicated less damage in Legacy, where radical scavenging activity (RSA) values had no significant changes in leaves. However, Bluegold exhibited the higher RSA values in leaves with the highest Mn treatment. With respect to organic acid anions root exudates, oxalate, and citrate presented the greater values in Legacy and Brigitta.

In Chapter 4, we reported that an excess of Mn in plants has a negative effect on the photosynthetic performance, especially in the first light reactions involving the photosystems I and II (PSI and PSII). The aim of this research was evaluate the role of specific Mn-induced changes in the structure and function of PSII and PSI, to help to understand the mechanisms by which Mn excess may cause a decrease in  $\text{CO}_2$  assimilation in *Arabidopsis thaliana*. We analyzed PSII and PSI operation, together with the protein composition of the photosynthetic apparatus against increasing levels of Mn. In agreement with earlier studies Mn excess caused minimal changes in the maximal photochemical efficiency of PSII measured as  $F_v/F_m$ , although the characteristic peak temperature of the  $S_{2/3}Q_B$ - charge recombinations was shifted to lower temperatures at the highest Mn concentration. SDS-PAGE and immunoblot analyses also did not exhibit any significant change in the relative abundance of PSII-associated polypeptides: PSII reaction center protein D1, Lhcb1 (major light harvesting protein of LHCII complex) and PsbO (OEC33 - 33kDa protein of the oxygen-evolving complex). In addition, the abundance of Rubisco did

not change with Mn treatments. Nonetheless, plants grown under excess Mn exhibited increased susceptibility to PSII photoinhibition. In contrast, *in vivo* measurements of the redox transients of PSI reaction centre (P700) showed a considerable gradual decrease in the extent of P700 photooxidation (P700<sup>+</sup>) under increased Mn concentrations compared to control. This was accompanied by a slower rate of P700<sup>+</sup> re-reduction indicating a down-regulation of the PSI-dependent cyclic electron flow. The abundance of PSI reaction center polypeptides (PsaA and PsaB) in plants under the highest Mn concentration was also significantly lower compared to the control. The results demonstrate for the first time that PSI is the major target of Mn toxicity within the photosynthetic apparatus of *Arabidopsis* plants. The possible involvement mechanisms of Mn toxicity targeting specifically PSI are discussed.

Finally, in Chapter 5 and 6 we present the general discussion and conclusions of the thesis, concluding that a Mn excess (1000 µM Mn) has frequently negative effects on physiological and biochemical parameters of the essayed blueberry cultivars. The most noticeable negative changes were exhibited by Bluegold, where the photosynthetic performance and aerial biomass were more reduced than in Legacy and Brigitta.

## **Table of contents**

---

<b>Thesis outline</b>	<b>i</b>
<b>Table of contents</b>	<b>iv</b>
<b>Table index</b>	<b>viii</b>
<b>Figure index</b>	<b>x</b>

### **Chapter 1. General Introduction**

1.1 Introduction	2
1.1.1 Acid soils	2
1.1.2 Mn in soils	2
1.1.3 Mn as micronutrient in plants	3
1.1.4 Physiological and biochemical effects of Mn excess in plants	3
1.1.5 Mechanism against Mn excess	4
1.1.6 Blueberry crop in Chile	6
1.2 Hypotheses of this thesis	7
1.3 General aim	7
1.4 Specific aims	7

**Chapter 2. Theoretical background: Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms.**

Abstract	9
2.1 Introduction	10
2.2 Manganese forms and dynamics in soils	10
2.2.1 Manganese dynamics in the rizosphere	12
2.3 Manganese transport and accumulation in plants	13
2.4 Manganese as an essential element in plant metabolism	16
2.5 Manganese phytotoxicity and injury symptoms	19
2.5.1 Manganese toxicity and oxidative stress	23
2.6 Tolerance mechanisms to manganese toxicity	24
2.7 Conclusions and perspectives	27
2.8 References	29

**Chapter 3. Effect of Mn excess on biochemical and physiological properties of Highbush blueberry (*Vaccinium corymbosum* L.) cultivars growing in nutrient solution.**

Abstract	41
3.1 Introduction	42
3.2 Materials and Methods	43
3.3 Results	46
3.4 Discussion	48
3.5 References	53

## **Chapter 4. Excess manganese (Mn) differentially inhibits Photosystem I (PSI)**

### **versus Photosystem II (PSII) in *Arabidopsis thaliana*.**

Abstract	69
4.1 Introduction	70
4.2 Materials and methods	72
4.2.1 Plant material and growth conditions	72
4.2.2 Mn treatments	72
4.2.3 Plant growth measurements	72
4.2.4 Manganese concentration	73
...4.2.5 Thylakoid preparation, SDS-PAGE and immunoblotting	73
...4.2.6 Measurement of the redox state of P <sub>700</sub> .	74
...4.2.7 Modulated chlorophyll fluorescence measurements	74
...4.2.8 Thermoluminescence measurements	75
...4.2.9 Experimental design and statistical analysis	76
4.3 Results	76
4.4 Discussion	79
4.5 References	84

## **Chapter 5. General discussion**

5.1 General discussion	107
------------------------	-----

## **Chapter 6. General conclusions**

6.1 General conclusions	115
6.2 Futures perspectives	115
6.3 General References	116
<b>Appendix</b>	126

## Table Index

---

<b>Chapter 2.</b>	<b>Theoretical background: Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms.</b>	
2.1	Some transporter proteins implicated in Mn <sup>2+</sup> and other cations transport and their cellular localization	18
2.2	Symptoms of Mn toxicity and Mn concentrations in organs of some plant species subjected to toxic Mn concentrations.	21
<b>Chapter 3</b>	<b>Effect of Mn excess on biochemical and physiological properties of Highbush blueberry (<i>Vaccinium corymbosum</i> L.) cultivars growing in nutrient solution.</b>	
3.1	Maximal photochemical efficiency of PSII (Fv/Fm), effective photochemical efficiency of PSII (ΦPSII) and non-photochemical quenching (NPQ) of three blueberry cultivars subjected to increasing Mn levels (25 days).	61
<b>Chapter 4</b>	<b>Excess manganese (Mn) differentially inhibits Photosystem I (PSI) versus Photosystem II (PSII) in <i>Arabidopsis thaliana</i>.</b>	
4.1	Relative growth rates (RGR) of <i>Arabidopsis thaliana</i> (leaves and roots) growing at increased Mn treatments under acidic conditions (pH 5.3) for 21 d in comparison with the control.	95
4.2	Maximal photochemical efficiency of PSII (Fv/Fm) in leaves of <i>Arabidopsis thaliana</i> measured at 0, 5, 10, 15 and 21 days subjected to increasing Mn treatments (μM).	96
4.3	Characteristic thermoluminescence (TL) peak emission temperatures (T <sub>M</sub> ) and the overall TL emission area (A) of S <sub>2</sub> S <sub>3</sub> Q <sub>B</sub> - glow peaks of control and Mn-treated (21 days)	97

- Arabidopsis* plants.
- 4.4** Effect of high light treatments ( $1000 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ , PFD, for 1, 2 and 4 hours) on the maximal photochemical efficiency of PSII measured as Fv/Fm in control *Arabidopsis thaliana* leaves and plants exposed for 21 days to different Mn doses. 98
- 4.5** Effects of Mn treatments on far-red light induced steady state oxidation of P700 ( $\Delta A_{820-860}$ , P700<sup>+</sup>), the relative intersystem electron donor pool size to PSI ( $e^- / \text{P700}$ ) and half times for P700<sup>+</sup> reduction ( $t_{1/2}$ ) of *Arabidopsis* leaves at 21 days of treatment. 99

## Figure Index

---

<b>Chapter 2.</b>	<b>Theoretical background: Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms.</b>	
2.1	The role of soil rhizosphere in the mobilization and immobilization of Mn in soils.	14
<b>Chapter 3</b>	<b>Effect of Mn excess on biochemical and physiological properties of Highbush blueberry (<i>Vaccinium corymbosum</i> L.) cultivars growing in nutrient solution.</b>	
3.1	Effect of Mn treatments on dry weight of shoot (leaves + stems) ( <b>A</b> ) and roots dry weight ( <b>B</b> ) in three blueberry cultivars grown in nutrient solution for 25 days.	59
3.2	Mn contents in leaves and roots of blueberry cultivars with Mn excess-doses at the end of experiment (25 days).	60
3.3	Net photosynthesis ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) and stomatal conductance ( $\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$ ) of three highbush blueberry cultivars subjected to increase Mn treatments at the end of experiment (25 days).	62
3.4	Lipid peroxidation of leaves and roots of three highbush blueberry cultivars subjected to different Mn treatments for 25 days.	63
3.5	Radical-scavenging capacity (RSA) of three highbush blueberry cultivars subjected to different Mn treatments at leaves and roots at 25 days, measured as Trolox equivalents (TE).	64

3.6	Effect of Mn-excess on the activity of SOD enzyme in leaves and roots of three blueberry cultivars with excess Mn treatments at 25 days of treatment.	65
3.7	Organic acid anions exuded by roots of blueberry cultivars grown with increasing Mn levels for 25 d.	66
<b>Chapter 4</b>	<b>Excess manganese (Mn) differentially inhibits Photosystem I (PSI) versus Photosystem II (PSII) in <i>Arabidopsis thaliana</i>.</b>	
4.1	Total biomass (leaves and roots, g DW) of <i>Arabidopsis thaliana</i> plants subjected to Mn treatments ( $\mu\text{M}$ ) at the end of experiment (21 days).	101
4.2	Mn concentration in leaves and roots of <i>Arabidopsis thaliana</i> plants subjected to Mn treatments ( $\mu\text{M}$ ) at the end of experiment (21 days).	102
4.3	Thermoluminescence (TL) glow curves of $\text{S}_2/\text{S}_3\text{Q}_\text{B}^-$ charge recombinations in control (solid lines) and Mn-treated ( $500 \mu\text{M Mn}^{2+}$ - dashed lines, $1500 \mu\text{M Mn}^{2+}$ - dotted lines) <i>Arabidopsis</i> leaves after illumination with two single turnover flashes.	103
4.4	Typical traces of <i>in vivo</i> measurements of P700 oxidation by far-red light (FR) in control (A, $18 \mu\text{M Mn}$ ) and Mn-treated (B, $1500 \mu\text{M Mn}$ ) <i>Arabidopsis</i> plants.	104
4.5	A - Representative immunoblots of SDS-PAGE separated proteins from thylakoid membranes probed with antibodies raised against PSI (PsaA and PsaB), PSII (D1, PsbO, Lhcb1)-associated polypeptides and Rubisco (PbcL) in control (C, $18 \mu\text{M Mn}$ ) and <i>Arabidopsis</i> plants treated with $500$ and $1500 \mu\text{M Mn}$ . B - The relative polypeptide abundance was quantified by densitometric analysis (area x	105

intensity bands) of the immunoblots and the presented data were normalized to the relative abundance of PsaA, PsaB, D1, PsbO, Lhcb1 and PbcL in control *Arabidopsis* plants.

## **Chapter 5**

### **General discussion**

- 5.1 Theoretical model showing effect of Mn excess on physiological and biochemical features in leaves and roots of Mn- tolerant and Mn-sensitive blueberry cultivars. 112

## Chapter 1

### General Introduction

## **1.1 Introduction**

### **1.1.1 Acid soils**

Acid soils represents up to 50% worldwide arable soils. Moreover, tropical and subtropical zones have a special significance because around 60% of their soils are acids, limiting the crops growth in several developing countries (Kochian et al., 2004). Acidification can be caused by high pluviometry in winter, which leads to a loss of interchangeable bases (Mora, 1993; Mora et al., 2006), high use of fertilizers acid reaction (ammonium type), with consequent decrease of calcium (Ca), magnesium (Mg), phosphorus (P) and/or molybdenum (Mo) availability (Foy et al., 1978; Mora et al., 1999).

Acid soils are found in southern Chile, with Andisol and Ultisol types mainly, from volcanic origin. These soils are developed under a cold and wet climate, with high organic matter content, and low availability of P and nitrogen (N) (Aguilera et al., 1997, 2002; Peñaloza et al., 2004). The main constraint reported in acid soils by pH among 4 to 5.5, is an increase of metals cation solubility, as aluminum (Al) and manganese (Mn) (Jarvis, 1987; Watmough et al., 2007), being Mn the second phytotoxic element after Al which limits growth of plants (Foy, 1984; Kochian et al. 2004).

### **1.1.2 Mn in soils.**

Apart of low pH, redox changes conditions of soils determine the Mn availability, solubility (Marschner, 1995; Pittman, 2005; Sposito, 1989; Gherardi and Rengel, 2004) and Mn content (Farasova and Beinrohr, 1998). According, Boojar and Goodarzi (2008), low pH enhance the Mn solubility to accelerate its release from soil solution. Under reduction conditions, soils have major amount of free electrons, which may be increased by water excess, poor drainage, waterlogged places or heavy organic matter applications, among others, even in soil with high pH (Sparrow and Uren 1987; Hue, 1988). In addition, to remove electron acceptor agents (as oxygen) from surrounding soil, there are an increase in the Mn availability (Porter et al., 2004). Therefore, Mn phytotoxicity is favored in acid and/or waterlogged soils, where both pH (<5.5) and redox conditions increase the concentration of the soluble Mn ( $Mn^{2+}$ ), which is the predominant Mn form in the soil acid solution (Marschner 1995; Porter et al. 2004).

### **1.1.3 Mn as micronutrient in plants.**

Mn is considered an essential micronutrient to plant growth and development. It can be absorbed mainly as  $Mn^{2+}$  ions (Mukhopadhyay y Sharma, 1991). Specifically, it is involved in water splitting in the photosystem II (PSII) of the photosynthetic process (Goussias et al., 2002). Also, Mn is a constitutive element associated with the oxygen evolving complex (OEC) of photosystem II (PSII), an important multi-protein pigment complex embedded in the thylakoid membranes (Hankamer et al., 1997; Enami et al., 2008). Therefore, the Mn cluster, together with other ions and extrinsic proteins that constitute OEC are required to oxidize water and reduce P680, the reaction center of PSII (Kern and Renger, 2007; Ferreira et al., 2004; Rutherford and Boussac, 2004). In conjunction with photosystem I (PSI) and linear electron transport, these reducing equivalents (electrons) are used primarily in the conversion of  $CO_2$  into carbohydrate (Ferreira et al., 2004). Mn also participates as metallic cofactor in superoxide dismutase enzyme (MnSOD), located in mitochondria of eukaryotic organisms and into SOD iso-enzymes group such as Fe (FeSOD), copper and zinc enzymes (Cu/ZnSOD) (Blokhina et al., 2003).

It has been estimated that Mn normal values of various agricultural crops range from 30 to 500  $mg\ kg^{-1}$  dry weight (DW) (Humpries et al., 2007). Pasture species had values fluctuating between 5 to 1200  $mg\ kg^{-1}$  in different zones (Whitehead, 2000). In southern Chile, species as white clover (*Trifolium repens* L.) and perennial ryegrass (*Lolium perenne* L.) have values of Mn concentrations between 50 to 1000  $mg\ kg^{-1}$ . Into bush species as blueberries (*Vaccinium corymbosum* L.), values up to 1600  $mg\ kg^{-1}$  in autumn and values between 280 to 560  $mg\ kg^{-1}$  of Mn are reported (Universidad de La Frontera, Soil and Plant Analysis Laboratory Services).

### **1.1.4 Physiological and biochemical effects of Mn excess in plants.**

As mentioned, trace element Mn can be a toxic element in plants when it is in excess (Marschner, 1995). Mn toxicity can show visual symptoms as chlorosis, brown speckles in the leaves (Wissemeier and Horst, 1992; González et al., 1998). Besides, Mn excess can produce several detrimental effects such as growth reduction (Mora et al., 2009), modifying

various processes such as activity of enzymes, absorption, translocation and utilization of other mineral elements (Ca, Mg, Fe and P) (Ducic and Polle, 2005; Lei et al 2007).

The Mn excess seems also to be particularly damaging to physiological processes as well as the photosynthetic apparatus (Mukhopadhyay and Sharma, 1991). In this context, it has been demonstrated that high Mn accumulation in thylakoids it is associated with inhibition of the net photosynthesis and carboxylation efficiency, suggesting that this metal may interfere with the energy fluxes by thylakoid stacking (Lidon and Teixeira, 2000; Lidon et al., 2004). The decline of photosynthesis is one of the mechanisms constituting the toxic effects of Mn excess, being an early indicator for Mn toxicity in tobacco (Nable et al., 1988). Furthermore, an inhibition of photosynthesis and photochemical parameters of Cucumber leaves was observed by increasing Mn concentrations, producing stomatal closure with a lower accumulation of biomass (Feng et al., 2009).

#### **1.1.5 Mechanism against Mn excess.**

To counteract and ameliorate the detrimental effects of Mn excess, plant tissues have Mn resistance mechanisms that can include avoidance or exclusion (Delhaize et al., 2007a) and tolerance or inclusion. The first one involves a protective role that prevents that the metal ions entering the cytoplasm of plant cells (Blamey et al 1986; Marschner 1991) and the second one (tolerance or inclusion) implies a detoxification of metal ions after they have crossed the plasma cell membrane or internal biomembranes of organelles (Taylor, 1991). Thus, organic acids exudations can be considered as resistance mechanisms against metal toxicity in several plant species (Delhaize et al., 2007a; Mora et al., 2009; Ryan et al., 2009). Besides, in case of Mn deficiency, some organic acids act as metal chelators in the rhizosphere, increasing the availability of metallic soil micronutrients (Dakora and Phillips, 2002).

Otherwise, one research area that have relevance to the field of Mn tolerance mechanisms in plant cell, involves the identification of several and numerous plant genes that encode transporters with the ability to internal transport, sequestrations, distribution and

homeostasis of  $Mn^{2+}$  (Pittman, 2005; Kochian et al 2004; Delhaize et al., 2007b; Sasaki et al., 2011; Hirschi et al., 2000). In the case of Mn, researches about genes expression suggested that genes associated to internal Mn tolerance are more abundant than those involved with Mn exclusion (Horst, 1988; Delhaize et al., 2003).

Other mechanism that can withstand an abiotic stress (as metal toxicity) is the activation of antioxidant systems. Reactive oxygen species (ROS) are intermediates products of molecular oxygen reduction to water molecules which cause damage at cell level (Schützendübel and Polle, 2002). Under normal conditions, plant cell generate ROS, which may increase when plant organelles are under both biotic and abiotic environmental stresses (Fecht-Christoffers et al., 2003). Effects of ROS involve oxidative damages on cell biomolecules such as lipids, proteins and nucleic acids, leading to a biomembrane peroxidation, loss fluidity, protein hydrolysis, inactivation of enzyme and even DNA chain breakdown (Demirevska-Kepova et al., 2004; Guo et al., 2007; Hegedüs et al., 2001; Feng et al., 2009; Srivastava and Dubey, 2011).

Antioxidant defense mechanisms can prevent and diminish the detrimental effects of ROS. These systems involved enzymatic mechanisms, as superoxide dismutase (SOD), peroxidase (APX), catalase (CAT), glutation peroxidase (GPX) and non-enzimatic mechanisms as ascorbate,  $\alpha$ -tocopherol, alkaloids, carotenoids, flavonoids and glutation (Foyer and Noctor, 2003; Apel and Hirt, 2004). Studies on Mn excess showed significant changes in antioxidant systems of some plants. Specifically, SOD activity is reported as a Mn-tolerance mechanism in rye grass cultivars subjected to Mn excess (Mora et al. 2009). Rosas et al., (2007) reported high activity of APX values in Mn toxic treatments (355  $\mu$ M Mn) in prairie species, indicating an oxidative stress. In the same way, several of these enzymes are considered scavengers of ROS produced by Mn toxicity in rice and cucumber plants (Srivastava and Dubey, 2011; Feng et al., 2009). In addition, Mn is required as a cofactor for Mn-SOD enzyme, essential to the protection against oxidative stress in plants (Burnell, 1988; Bowler et al., 1994).

### **1.1.6 Blueberry crop in Chile.**

In Chile, *Vaccinium corymbosum* L. (highbush blueberry) is an important crop, with 12,000 ha, concentrated in the South Central area. Recently, in the region where our study was carried out, there are approximately 1,500 ha of blueberry crop (ODEPA and CIREN, 2012). Likewise, its production reached 100,000 ton approximately in year 2011, being Chilean blueberry industry a global leadership and the largest exporter of this species (ODEPA, 2012).

Blueberry fruits are considered the richest sources of antioxidant compounds of different species of fresh fruits and vegetables which contribute to a high antioxidant capacity (Prior et al., 1998; Howard et al., 2003). Phenolic acids, anthocyanins and flavonoids are the phytochemicals that mostly influence the antioxidant capacity of plants (Prior et al., 1998).

Although, blueberry crop is well adapted to soils under acidic conditions (Ireland and Wilk, 2006), the effect of Mn excess on this crop is unknown. Therefore, the main focus of this investigation intends to clarify the physiological and biochemical response of *V. corymbosum* under excess of Mn, in order to elucidate the tolerance mechanisms involved in the Mn excess. In addition, a complementary study was conducted to obtain information concerning to the toxic effect(s) of excess of Mn on the polypeptides of both PSII and especially PSI using the model plant *Arabidopsis thaliana*.

## 1.2 Hypotheses of this thesis

High concentrations of available Mn ( $Mn^{2+}$ ) are associated with soil acidity. These ions can be accumulated in plant cell tissues producing toxicity, which alters physiological and biochemical features. This can stimulate defense plant mechanisms to exclude or tolerate this toxicity.

We hypothesized:

- i)* Blueberry cultivars will have differential capacities to accumulate Mn and therefore different capacity to tolerate oxidative stress.
- ii)* The Mn-resistant cultivar will have a higher photosynthetic capacity and better Mn-tolerance mechanisms. Also, in a model plant, Mn excess decreases the activity and/or performance of the photosystems (PSI and PSII) involved in the photosynthetic apparatus.

## 1.3 General aim

To evaluate Mn toxicity effects on physiological and biochemical properties of photosynthetic apparatus in highbush blueberry cultivars, as well as to analyze their resistance mechanisms by Mn excess. Additionally, to evaluate the susceptibility of the proteins of PSI and PSII to Mn excess in a model plant (*Arabidopsis thaliana*).

## 1.4 Specific aims

1. To evaluate specific mechanisms of Mn resistance (inclusion and/or exclusion) and antioxidant mechanisms in a Mn tolerant and a Mn sensitive cultivar of blueberry in the presence of Mn excess.
2. To compare physiological changes of photosynthetic apparatus in a Mn tolerant and a Mn sensitive blueberry cultivar under Mn excess.
3. To evaluate changes in proteins related to photosystems (PSI and PSII) in a plant model exposed to an excess of Mn.

## Chapter 2

Theoretical background.

Manganese as essential and toxic element for plants:  
transport, accumulation and resistance mechanisms.

Paper published in Journal of Soil Science and Plant Nutrition (2010) 10:470-481

DOI: 10.4067/S0718-95162010000200008

# **Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms**

R. Millaleo<sup>a</sup>, M. Reyes-Díaz<sup>b</sup>, A.G. Ivanov<sup>c</sup>, M. L. Mora<sup>d</sup>, M. Alberdi<sup>d\*</sup>

<sup>a</sup>Programa de Doctorado en Ciencias de Recursos Naturales, Universidad de La Frontera, Casilla 54-D, Temuco, Chile. <sup>b</sup>Center of Plant-Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Temuco, Chile. <sup>c</sup>Department of Biology and the Biotron, University of Western Ontario, London, Ontario, N6A 5B7 Canada. <sup>d</sup>Departamento de Ciencias Químicas, Center of Plant-Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Temuco, Chile.

## **Abstract**

Manganese is an essential element for plants, intervening in several metabolic processes, mainly in photosynthesis and as an enzyme antioxidant-cofactor. Nevertheless, an excess of this micronutrient is toxic for plants. Mn phytotoxicity is manifested in a reduction of biomass and photosynthesis, and biochemical disorders such as oxidative stress. Some studies on Mn toxicity and Mn translocation from soil to plant cells in Mn<sup>2+</sup> form have demonstrated their importance under low pH and redox potential conditions in the soil. When Mn is inside the cells, mechanisms that can tolerate this toxicity are also observed, being important the compartmentalization of this metal in different organelles of shoot and leaf plant cells. A key role of antioxidative systems in plants in relation to high Mn amounts has also been reported as a defense mechanism. The purpose of this review is to show the role of Mn as an essential micronutrient and as a toxic element to higher plants as well as to their transport and tolerance mechanisms. The forms and dynamics of this element in soils and the importance of the acidity for this dynamic and availability for plants are also given.

*Keywords:* Manganese, Mn toxicity, resistance mechanisms.

## 2.1 Introduction

Manganese (Mn) is an essential micronutrient in most organisms. In plants, it participates in the structure of photosynthetic proteins and enzymes. Its deficit is dangerous for chloroplasts because it affects the water-splitting system of photosystem II (PSII), which provides the necessary electrons for photosynthesis (Buchanan, 2000). However, its excess seems also to be particularly damaging to the photosynthetic apparatus (Mukhopadhyay and Sharma, 1991). Thus, Mn has two roles in the plant metabolic processes: as an essential micronutrient and as a toxic element when it is in excess (Kochian et al., 2004; Ducic and Polle, 2005). Mn toxicity is favored in acid soils (Pendias and Pendias, 1992). With decreasing pH, the amount of exchangeable manganese – mainly  $Mn^{2+}$  form – increases in the soil solution. This Mn form is available for plants and can be readily transported into the root cells and translocated to the shoots, where it is finally accumulated (Marschner, 1995). In contrast, other forms of Mn predominate at higher pH values, such as Mn (III) and Mn (IV), which are not available and cannot be accumulated in plants (Rengel, 2000).

Excessive Mn concentrations in plant tissues can alter various processes, such as enzyme activity, absorption, translocation and utilization of other mineral elements (Ca, Mg, Fe and P), causing oxidative stress (Ducic and Polle, 2005; Lei *et al.*, 2007). The threshold of Mn injury as well as the tolerance to an excess of this metal is highly dependent on the plant species and cultivars or genotypes within a species (Foy *et al.*, 1988, Horst, 1988).

The purpose of this review is to illustrate the most current understanding about Mn role as an essential micronutrient and as a toxic element to higher plants, the long distance and cellular transport in plants as well as the mechanisms or strategies involved for to resist an overload of this metal. The forms and dynamics of this element in soils and the importance of the acidity for this dynamic and availability to plants are also given.

## 2.2 Manganese forms and dynamics in soils.

Manganese biogeochemistry in soils is complex, because it is present in several oxidation states (0, II, III, IV, VI and VII), while in biological systems it occurs preferably as II, III and IV. Divalent manganese (Mn II) is the most soluble species of Mn in soil, whereas the solubility of Mn III and Mn IV are very low (Guest *et al.*, 2002). Mn oxides can form co-

precipitates with iron (Fe) oxides, exhibiting amphoteric behavior. In addition, Mn interacts both with cations and anions in oxidation-reduction reactions involving Mn. These reactions are influenced by a variety of physical, chemical and microbiological processes (Bradl, 2004).

Both pH and redox conditions influence Mn bioavailability in soils (Marschner, 1995; Porter *et al.*, 2004). In most acid soils at low pH (<5.5) and an increased redox potential of Mn, oxides can be easily reduced in the soil exchange sites (Kogelmann and Sharpe, 2006), increasing the concentration of soluble  $Mn^{2+}$  (Watmough *et al.*, 2007), which is the predominant Mn form in the soil solution (Adriano, 2001) and the most available Mn form for plants (Marschner, 1995). At higher soil pH (up to pH 8), chemical  $Mn^{2+}$  auto-oxidation is favored over  $MnO_2$ ,  $Mn_2O_3$ ,  $Mn_3O_4$  and even  $Mn_2O_7$ , which are not normally available to plants (Ducic and Polle, 2005; Humpries *et al.*, 2007; Gherardi and Rengel, 2004). Furthermore, high pH allows Mn adsorption into soil particles, decreasing their availability (Fageria *et al.*, 2002). Nevertheless, some reports have suggested that an excess of available Mn is produced under reduced soil conditions, even at high soil pH values (Hue, 1988). A reducing environment can be produced when there is an excess of water, poor drainage or applications of organic material (Hue, 1988; El-Jaoual and Cox, 1998). Different organic molecules can dissolve solid Mn oxides through transfer of electrons, transforming them into an available Mn form for plants (Laha and Luthy, 1990). Soil acidification is also accentuated by abundant pluviometry during winter, causing the main cations to leak from the soil (Mora *et al.*, 2006). On the other hand, lime application is a key factor in decreasing soluble Mn in acid soils with a high Mn content, given that it can increase soil pH (Hue and Mai, 2002). The total Mn content in soils is variable. Sparks (1995) reported small amounts of Mn in soils, fluctuating from 20 to 10,000  $mg\ kg^{-1}$  soil, whereas other authors have registered total Mn contents between 450 and  $\sim 4,000\ mg\ Mn\ kg^{-1}$  soil (Adriano, 2001). In addition, the total Mn soil content was from 15 to 17  $mg\ kg^{-1}$  in acid soils without liming (pH about 4.4) (Hue and Mai, 2002). In liming acid soils, the interchangeable Mn concentration varied from 14 to 96  $mg\ kg^{-1}$  soil in one year, with higher concentrations under high moisture and temperature conditions (Conyers *et al.*, 1997). In Chilean volcanic soils, so-called Andisols, Mn concentrations fluctuate between 4.5 and 80  $mg\ kg^{-1}$  depending on the agronomic management. Moreover, the Mn amount is


higher in pasture soils (up to 400 mg kg<sup>-1</sup>) mainly in winter (Data from Laboratorio de Análisis de Suelo y Planta, Universidad de La Frontera, Temuco, Chile). Environmental conditions also affect Mn soil contents. The highest concentrations of soluble and exchangeable Mn are found after hot, dry summers and under warm waterlogged conditions in acid soils. This is probably due to the inhibition of Mn-oxidizing organisms, thereby allowing the chemical reduction of Mn oxides in these soils (Sparrow and Uren, 1987; Conyers *et al.*, 1997).

### **2.2.1 Manganese dynamics in the rizosphere**

Rhizosphere, which is the narrow zone of soil immediately surrounding the root system, is of great importance for mineral plant nutrition. In this zone, both the mobilization and immobilization of nutrients occur (Marschner, 1995). As shown in Figure 2.1, a mobilization of Mn<sup>2+</sup> is produced by the rhizosphere acidification due to the release of H<sup>+</sup> or low molecular weight organic acids (LMWOA) from plants (Rengel and Marschner, 2005). Organic acids released in anion forms from roots can chelate Mn<sup>2+</sup> released from the MnO<sub>x</sub> (Mn oxides) (Ryan *et al.*, 2001). Neumann and Römheld (2001) reported that mobilization of micronutrients (including Mn) into the rhizosphere is due mainly by its acidification and complexation with the organic acids (citrate) in various plant species. It has been reported that organic amendments (chip compost and pine bark) applied to melon plants released organic compounds such as arabinose and malic acid that can dissolve MnO<sub>x</sub> (Tsuji *et al.*, 2006). Soil microorganisms can also help Mn mobilization and immobilization, depending on soil conditions (Marschner, 1995). In aerated soils, microorganisms may mobilize Mn through MnO<sub>x</sub> reduction favored by H<sup>+</sup> root excretion. In contrast, Mn-oxidizing bacteria can decrease Mn availability in aerated and calcareous soils or in poorly aerated and/or submerged soils. Another key factor in the Mn dynamics in soil is organic matter (OM). Given that OM is negatively charged, it has a great Mn adsorption capacity, forming Mn complexes which decrease the amount of exchangeable Mn. However, the Mn adsorbed by OM can be exchanged by the H<sup>+</sup> released from the roots (Bradl, 2004).

### 2.3 Manganese transport and accumulation in plants

As mentioned above, reduced Mn ( $Mn^{2+}$ ) form is the only available metal form for plants. It can be taken up via an active transport system in epidermal root cells and transported as divalent cation  $Mn^{2+}$  into the plants (Marschner, 1995; Gherardi and Rengel, 2003; Pittman, 2005). Manganese uptake by roots is characterized as a biphasic process. The initial and rapid uptake phase is reversible and non-metabolic, with  $Mn^{2+}$  and  $Ca^{2+}$  or other cations being freely exchanged in the rhizosphere. In this phase,  $Mn^{2+}$  appears to be adsorbed by the negatively charged cell wall constituents of the root-cell apoplastic spaces (Humphries *et al.*, 2007; Clarkson, 1988). The second phase is slow, with  $Mn^{2+}$  being less readily exchanged. Its uptake into the symplast is dependent on plant metabolism (Maas and Moore, 1968), although the exact mechanisms are not clear (Humphries *et al.*, 2007). It has been shown that in transgenic tobacco transformed with a tomato root protein with a metal binding site at its N-terminus (LeGlp1), Mn binds to this protein. This strongly suggests the involvement of LeGlp1 in Mn uptake from the soil (Takahashi and Sugiura, 2001). Kinetic measurements have demonstrated 100 to 1,000 times higher rates of Mn transport than the estimated plant requirement for this element (Clarkson, 1988). These transport rates are explained by the high capacity of ion carriers and channels in the Mn ion transportation through the plasma membrane at a speed of several hundred to several million ions per second per protein molecule (Humphries *et al.*, 2007). According to these authors, Mn distribution from root cells within the whole plant involves primary transport in the xylem, transference from the xylem to the phloem and re-translocation into the phloem. Xylem transport from roots to the above-ground parts of plants is performed by the transpiration stream, whereas phloem transport is more selective, taking place from sources to sinks (Marschner, 1995).



**Figure 2.1.** The role of soil rhizosphere in the mobilization and immobilization of Mn in soils.

Nonetheless, a low mobility in phloem has been reported for Mn, and its redistribution may depend on the plant species and stages of development (Herren and Feller, 1994). In fact, it has been reported that Mn transport from roots to grains is frequently insufficient at a mature stage of wheat. The relatively poor Mn mobility in the phloem emphasizes the importance of xylem in the transport of this element, even in wheat grain discharge (Rengel, 2001). Manganese generally tends to accumulate predominantly in the plant shoots than in the roots, as demonstrated in Mn labeling experiments with  $^{54}\text{Mn}$  at an early stage of wheat (*Triticum aestivum* cv. Arina) development, where a fast Mn transport from roots to shoots was visualized in the xylem and was essentially immobile in the phloem (Page and Feller, 2005). Similar effects on Mn translocation have been shown by the same technique in young (28 days) white lupine plants (*Lupinus albus*) (Page *et al.*, 2006). Nevertheless, Mn was present in a large amount in the root system, hypocotyls and stem in older lupine plants, immediately after the labeling phase (day 0). Seven days later (day 7) almost all  $^{54}\text{Mn}$  had moved to the youngest fully expanded leaves and only a small fraction to the other leaves. Mn accumulation was observed in the periphery of the oldest leaves. These authors reported that Mn was rapidly released from the roots into the xylem,

reaching photosynthetically active leaves via the transpiration stream. Furthermore, the low mobility of this element via phloem in the shoot may be due to a restricted loading of soluble Mn into the phloem or by insolubilization in the leaves, although the issue remains to be clarified (Page *et al.*, 2006). Page and Feller (2005) emphasized that little is known about the mechanisms involved in the loading of Mn into the phloem and the chemical transport forms.

In addition to the long distance transport of Mn, short distance transport mechanisms are important for the translocation of this metal into the cell and cell organelles. These mechanisms involve Mn translocation throughout the plasma membrane and the biomembranes of organelles (Ducic and Polle, 2005; Pittman, 2005). Possible mechanisms of homeostasis and Mn transport, based on studies performed in yeast (*Saccharomyces cerevisiae*) cells and in *Arabidopsis thaliana* plants, have been discussed (Delhaize *et al.*, 2007; Reddi *et al.*, 2009). They pointed out that transport proteins play an important role for the maintenance of adequate Mn concentrations in the cytoplasm. Moreover, a variety of metal transporter family proteins with a broad-specificity such as Fe<sup>2+</sup> and Ca<sup>2+</sup> transporters have also the ability to transport Mn into the plant cells (Pittman, 2005). Migocka and Klobus (2007) demonstrated the activation of an antiport system in *Cucumis sativus* with different affinities for Pb, Mn, Ni and Cd in the root plasma membrane. This antiport system participates as part of the general defense mechanism activated under heavy metal stress. Using tonoplast-enriched vesicles, Shigaki *et al.*, (2003) suggested that a Cd/H antiporter might also be involved in Mn accumulation in vacuoles.

In Table 1 we have summarized some of the metal transporter proteins, their families, cellular localization and the transported metals. Transporter proteins are described according to their localization. In the plasma membrane, four Mn<sup>2+</sup> uptake transporters are identified: AtIRT1, Nramp, AtYSL (Ducic and Polle, 2005; Pittman, 2005) and PHO84 (Ducic and Polle, 2005). IRT1 can transport Mn when expressed in yeast (Korshunova *et al.*, 1999). Nramp is also considered a metal transporter protein that can transport Mn away from other cations. This protein may be localized in the tonoplast rather than in the plasma membranes (Thomine *et al.*, 2003). AtYSL belongs to the yellow stripe-like (YSL) proteins, also involved in metal-complex transport in the plasma membrane (Roberts *et al.*,

2004). This complex can be formed by nicotianamine (NA), which is a strong chelator of metals including  $Mn^{2+}$  (Pittman, 2005). PHO84 is a transporter protein identified in *S. cerevisiae* and it has a high-affinity phosphate uptake (Mitsukawa *et al.*, 1997). Luk *et al.* (2003) reported a new form in Mn transport as  $MnHPO_4$ . However, there is no evidence for  $Mn^{2+}$  or  $MnHPO_4$  accumulation by a plant phosphate transporter (Pittman, 2005).

Some transport proteins have been related to  $Mn^{2+}$  transport and accumulation into the intracellular compartments, such as the vacuole. It has been suggested that a metal transporter (specifically antiporter CAX2, calcium exchanger 2) originally identified as a  $Ca^{2+}$  transporter (which can also transport  $Cd^{2+}$ ) located in the cytosol. It has also the ability to transport Mn to the vacuole in tobacco plants (*Nicotiana tabacum*) and yeast (Hirschi *et al.*, 2000; Pittman, 2005) (Table 1). ATP-binding cassette (ABC) protein transporters are considered to be involved in detoxification processes (Martinoia *et al.*, 2002). Studies on cyanobacteria also suggested the putative role of these proteins in  $Mn^{2+}$  transport (Bartsevich and Pakrasi, 1996).

Another protein, considered indirectly as a metal transport protein is the ShMTP1, which is able to sequester metal ions within cells or efflux them out of the cells (Delhaize *et al.*, 2003). Therefore, these authors considered it a metal-tolerant protein. Ducic and Polle, (2005) and Pittman (2005) highlighted that, despite the available information about Mn transport across membranes in plant cells, the Mn transport and efflux strategies into the mitochondria, chloroplasts and Golgi are not completely understood. Nonetheless, Mills *et al.* (2008) have recently identified a Ca-ATPase that also transports  $Mn^{2+}$  into Golgi apparatus (Table 1).

#### **2.4 Manganese as an essential element in plant metabolism**

The main Mn role in photosynthesis is its involvement in the water-splitting system of photosystem II (PSII), which provides electrons necessary for photosynthetic electron transport. In water photolysis, a group of four Mn atoms (Mn cluster) is associated with the oxygen evolving complex (OEC) bound to the reaction center protein (D1) of PSII (Goussias *et al.*, 2002). The Mn cluster in PSII accumulates four positive charges, which

oxidize two water molecules, releasing one O<sub>2</sub> molecule and four protons. Therefore, this metal cluster is considered a catalyst compound of water oxidation (Zouni *et al.*, 2001), where Mn ions are close to a redox-active tyrosine residue (Z and D) (Goussias *et al.*, 2002).

Manganese also plays a role in ATP synthesis (Pfeffer *et al.*, 1986), in RuBP carboxylase reactions (Houtz *et al.*, 1988) and the biosynthesis of fatty acids, acyl lipids and proteins (Ness and Woolhouse, 1980). In addition, Mn plays a primary role in the activation and as cofactor of various enzymes in plants (~35) (Burnell, 1988), such as: Mn-superoxide dismutase, Mn-catalase, pyruvate carboxylase and phospho-enolpyruvate carboxykinase (Ducic and Polle, 2005). Manganese is also essential for the biosynthesis of chlorophyll (through the activation of specific enzymes), aromatic amino acids (tyrosine), secondary products, like lignin and flavonoids (Lidon *et al.*, 2004). It also participates in the biosynthetic pathway of isoprenoids (Lidon *et al.*, 2004) and assimilation of nitrate (Ducic and Polle, 2005). Hence, Mn is involved in metabolic processes such as respiration, photosynthesis, synthesis of aminoacids and hormone activation (indol acetic acid, IAA) throughout the IAA-oxidases (Burnell, 1988).

**Table 1.** Some transporter proteins implicated in Mn<sup>2+</sup> and other cations transport and their cellular localization. (Summarized from Ducic and Polle, 2005; Pittman, 2005 and Mills *et al.*, 2008). Abbreviations as follows: At= *Arabidopsis thaliana*; Sh= *Stylosanthes hamata*).

Transporter proteins	Protein cellular localization and protein family transporters	Transported ions
AtIRT1	Plasma membrane protein (ZIP, zinc-regulated transporter/iron regulated transporter (ZRT/IRT1) related protein) family transporter	Mn <sup>2+</sup> and Fe <sup>2+</sup> , Zn <sup>2+</sup> and Cd <sup>2+</sup> under Fe-deficiency conditions.
AtECA1	Endoplasmic reticulum (ER) Ca <sup>2+</sup> - and Mn <sup>2+</sup> -transporting P-type ATPase	Ca <sup>2+</sup> and Mn <sup>2+</sup>
AtCAX2	Vacuolar cation/H <sup>+</sup> antiporter CAX (the cation exchanger)	Mn <sup>2+</sup> , Ca <sup>2+</sup> and Cd <sup>2+</sup>
AtNramp3	Vacuolar Nramp transporter. Also, possible plasma membrane localization (Nramp?)	Mn <sup>2+</sup> , Fe <sup>2+</sup> and Cd <sup>2+</sup> in Fe-deficiency conditions
ShMTP1	Vacuolar-localized cation diffusion facilitator (CDF) family transporter	Related to the <i>Stylosanthes hamata</i> Mn <sup>2+</sup> transporter
ABC	Vacuolar- localized ATP binding cassette transporter families	Related to the cyanobacterium <i>Synechocystis</i> Mn <sup>2+</sup> transporter
AtOPT3	Probably located in the plasma membrane (AtOPT3?). It is an oligopeptide transporter-like protein (OPT).	Possible transport of Cu <sup>2+</sup> and Fe <sup>2+</sup> and Mn <sup>2+</sup>
AtYSL	Probably located in the plasma membrane. Yellow stripe-like transporter with equivalent function to rice OsYSL2, a Mn <sup>2+</sup> -nicotianamine (NA) and Fe <sup>2+</sup> -NA transporter	Mn <sup>2+</sup> and Fe <sup>2+</sup>
AtECA3	Ca <sup>2+</sup> transporters (Ca-ATPases) in Golgi.	Mn <sup>2+</sup> and Ca <sup>2+</sup>
PHO84	Probably located in the plasma membrane. It is MnHPO <sub>4</sub> transporter.	Mn binding to phosphate.

As a cofactor of superoxide dismutase (SOD), manganese participates in the plant's defense against oxidative stress, produced by elevated levels of activated forms of oxygen and free radicals (reactive oxygen species, ROS), which are harmful to plants. It has been proposed that Mn can act as a scavenger of superoxide ( $O_2^-$ ) and hydrogen peroxide ( $H_2O_2$ ). However, this mechanism is still unclear (Ducic and Polle, 2005).

Manganese SOD (MnSOD) belongs to the group of metal-containing SOD enzymes which are classified according to their metal cofactor: iron SOD (Fe-SOD), localized in the chloroplast; copper-zinc SOD (Cu/Zn-SOD), located in the chloroplast, cytosol, and possibly in the extracellular space. Manganese SOD is found mainly in mitochondria (Clemens *et al.*, 2002) and peroxisomes (Alscher *et al.*, 2002). MnSOD may play an important role in the adaptive responses of plant cells under environmental stresses such as salt stress, enhancing their tolerance. This tolerance has been demonstrated in transgenic plants of *Arabidopsis*, where Mn-SOD was overexpressed (Wang *et al.*, 2004). Similarly, in tomato transgenic plants which overexpressed MnSOD, an improvement in tolerance to NaCl stress in seedlings was concomitant with an improvement in seed germination and root development (Wang *et al.*, 2007).

## **2.5 Manganese phytotoxicity and injury symptoms**

As an essential micronutrient, low Mn levels are absolutely necessary for normal nutrition and development of plants. Normal Mn contents of leaves differ greatly between species (30-500 mg kg<sup>-1</sup> Mn dry mass, Clarkson, 1988). Nonetheless, when it is present in excessive amounts, it is extremely toxic to plant cells (Migocka and Klobus, 2007). The injury extent of Mn toxicity is approximately proportionate to the concentration of accumulated Mn excess. However, there is considerable inter- and intra- specific variation among Mn levels that induce toxicity as well as the symptoms of this toxicity in plant species (Foy *et al.*, 1988).

In addition to a decrease in growth rate, symptoms of Mn toxicity such as chlorosis in leaves (interveinal and marginal) and necrotic leaf spots are very common and have been reported in the whole plants of canola (Moroni *et al.*, 2003), clover (Rosas *et al.*, 2007),

ryegrass (Mora *et al.*, 2009) as well as in leaves of barley and cowpea (Demirevska-Kepova *et al.*, 2004; Fühns *et al.*, 2008) (Table 2). Necrotic brown spots and chlorotic leaves are frequently reliable indicators of the severity of Mn toxicity in plants (Wissemeier and Horst, 1991). The interveinal chlorosis due to Mn toxicity can have an appearance similar to that observed under Fe deficiency (Sarkar *et al.*, 2004). Moreover, the Mn toxicity is intensified when other available elements such as Ca, Mg, K, Fe and Si are in a low quantity (Abou *et al.*, 2002). However, a decrease in productivity by Mn toxicity without the appearance of leaf visual symptoms is sometimes observed (Miner and Sims, 1983). It is important to know that all these symptoms induced by Mn toxicity are preceded by an alteration of the photosynthetic apparatus and the photosynthetic performance of plants.

Other studies have shown that in rice (*Oryza sativa* cv. Safari) exposed to Mn excess in a nutrient solution, Mn was predominantly accumulated in leaves compared with roots (Lidon, 2001), whereas in *Sinapis alba* Mn mostly accumulated in the shoots (Farasova and Beinrohr, 1998).

**Table 2.** Symptoms of Mn toxicity and Mn concentrations in organs of some plant species subjected to toxic Mn concentrations according references from the last decade (earlier references in the text). Mn treatments were performed in nutrient solutions with MnCl<sub>2</sub> or MnSO<sub>4</sub>.

Species	Mn treatment	Mn concentrations in different plant organs	Symptoms of Mn toxicity	References
Rice ( <i>Oryza sativa</i> L.)	583 µM	Shoots: 2020 µg g <sup>-1</sup> dw	Decrease of shoot growth rate.	Lidon and Texeira (2000a)
Barley ( <i>Hordeum vulgare</i> L.)	1830 µM	Leaves: 656 mg g <sup>-1</sup> dw	Dark-brown necrotic spots, individually or in groups.	Demirevska-Kepova <i>et al.</i> (2004)
	18300 µM	Leaves: 1615 mg g <sup>-1</sup> dw		
Ryegrass ( <i>Lolium perenne</i> L.)	355 µM	Shoot: 2357 mg kg <sup>-1</sup>	Chlorosis and necrotic leaves.	Rosas <i>et al.</i> (2007)
		Roots: 2408 mg kg <sup>-1</sup>	Decrease of dry weight in roots.	
Ryegrass ( <i>L. perenne</i> L.)	150 µM	Shoot: 902 mg kg <sup>-1</sup>	Dry weight reduction.	Mora <i>et al.</i> (2009)
		Roots: 1342 mg kg <sup>-1</sup>	Dry weight reduction.	
Clover ( <i>Trifolium repens</i> L.)	355 µM	Shoot: 2050 mg kg <sup>-1</sup>	Reddish borders on leaves.	Rosas <i>et al.</i> (2007)
		Root: 7481 mg kg <sup>-1</sup>	Decrease of dry weight.	
Soybean ( <i>Glycine max</i> L.)	200 µM	Leaves: 806 mg kg <sup>-1</sup>	Chlorotic leaves.	Lavres Jr <i>et al.</i> (2009)
		Roots: 502 mg kg <sup>-1</sup>	No visual symptoms, increase in root diameter.	
Cowpea ( <i>Vigna unguiculata</i> L.)	50 µM	Leaves: ~ 25 µmol g <sup>-1</sup> dw	Brown spots on leaves.	Fühns <i>et al.</i> (2008, 2009)
Canola ( <i>Brassica napus</i> L.)	200 µM	Shoot: ~ 3500 µg g <sup>-1</sup> dw	Necrotic leaf spots, chlorosis in leaf margin.	Moroni <i>et al.</i> (2003)
<i>Juncus effuses</i> L. (wetland plant)	500 µM	176 mg kg <sup>-1</sup>	Reduction in plant dry biomass and height, no phytotoxic visual symptoms.	Najeeb <i>et al.</i> (2009)
<i>Populus cathayana</i>	1000 µM	Leaves: 713 mg kg <sup>-1</sup> dw	Decrease in shoot height, total biomass, and total leaf area.	Lei <i>et al.</i> (2007)

---

Despite the importance of Mn excess in the photosynthetic performance of plants, only a few studies about this issue are available. A reduction in photosynthesis, in chlorophyll *a* and *b* contents and their biosynthesis, as well as a reduction in carotenoids is frequently found in plants and also in algae under Mn excess (Macfie and Taylor, 1992; Hauck *et al.*, 2003). In rice cultivated at different Mn concentrations (from 2.3 to 583  $\mu\text{M}$ ), a significant decrease in chlorophyll *a* content has been reported at the highest Mn concentration (Lidon and Teixeira, 2000a). Nable *et al.* (1988) reported an early inhibition of photosynthesis concomitant with a high Mn accumulation in the leaves in *Nicotiana tabacum* cultivated in nutrient solutions with a Mn excess (1,000  $\mu\text{M}$ ). The authors concluded that the inhibition of photosynthesis is an early indicator for Mn toxicity in tobacco leaves. Lidon *et al.*, (2004) also observed a decline in net photosynthesis ( $\mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$ ) and photosynthetic capacity ( $\mu\text{mol O}_2 \text{ m}^{-2}\text{s}^{-1}$ ) in rice plants subjected to 9.1 and 36.4  $\mu\text{M}$  Mn treatment, respectively, with no changes in the levels of the ratio between variable (Fv) and maximum (Fm) chlorophyll fluorescence (Fv/Fm). However, an increase in the photochemical quenching and the quantum yield of non-cyclic electron transport was found up to 36.4  $\mu\text{M}$  Mn. Since Mn is accumulated in thylakoids, this element may interfere with thylakoid stacking, decreasing net photosynthesis (Lidon and Teixeira, 2000a).

Similarly to Mn toxicity, Mn deficiency also depressed leaf photosynthetic capacity in plants of *Carya illinoensis*, primarily by reducing the number of PSII units per unit leaf area, but these PSII units maintained abilities similar to those of the control plants (Henriques, 2003). Increased Mn concentrations (0 – 1,000 and 10,000  $\mu\text{M}$  Mn) in wheat plants inhibited the biosynthesis of chlorophyll and carotenoids, inducing a decrease in photosynthetic electron transport rates and therefore a decrease in the rate of photosynthesis (Macfie and Taylor, 1992). Higher Mn concentrations are also involved in a shortening of root and shoot and in a decreased chlorophyll concentration in the Mn accumulator plant, *Alyssum murale* (Abou *et al.*, 2002). Recently, Amao and Ohashi (2008) suggested that high Mn amounts in spinach leaves inhibited the activity of oxygen evolved complex of PSII.

---

### 2.5.1 Manganese toxicity and oxidative stress

Mn toxicity can also trigger oxidative stress in plant cells (Demirevska-Kepova *et al.*, 2004). As a toxic metal, Mn can cause metabolic alterations and macromolecular damage that disrupt the cell homeostasis (Hegedüs *et al.*, 2001; Polle, 2001). According to Lynch and St. Clair (2004), Mn toxicity in plants generates reactive oxygen species (ROS), mainly OH<sup>·</sup>, the most reactive oxidant and harmful species in cells (Lidon and Henriques, 1993). With respect to the oxidative stress responses to Mn excess, Lidon and Teixeira (2000b) reported that at the first growth stages of rice, two kinetic phases can be distinguished: in the first one, there is an increase in Mn accumulation in the thylakoid lamellae (McCain and Markley, 1989), which inhibits electron leakage from the Hill and Mehler reactions, limiting ROS formation. In the second kinetic phase, higher Mn amounts inhibit the non-cyclic photophosphorylation process, promoting an increase in ROS production that parallels an injury increase in the thylakoid peroxidase system (Lidon and Teixeira, 2000b). These authors concluded that Mn excess increases the disorganization in chloroplast lamellae, but elevated activity of superoxide dismutase (SOD) still limits cell damage.

It has also been reported that in *Cucumis sativus* plants both Mn excess and optimum light intensity determine an enhancement in oxidative stress by increased Mn content in the tissues concomitant with an inhibition of plant growth (Shi *et al.*, 2006). Investigations performed by González *et al.*, (1998) showed that lipid peroxidation was not induced by Mn-toxicity stress in the mature leaves of *Phaseolus vulgaris*, although other studies have shown that lipid peroxidation occurred in isolated chloroplast of wheat (Panda *et al.*, 1986). However, González *et al.* (1998) mentioned that this damage process could be related to the development stage of leaves, with the damage being more intense in immature than in mature leaves.

The geographical origin of the species and climatic conditions also affect the degree of Mn toxicity in plant species or populations as shown in two populations of *Populus cathayana*, coming from a wet and dry climate cultivated in an acid solution with increasing Mn concentrations (Lei *et al.*, 2007). The results showed that the wet climate population accumulated more Mn in plant tissues especially in leaves, decreasing their growth, chlorophyll contents, and activities of antioxidant enzymes than the dry climate population.

---

## 2.6 Tolerance mechanisms to manganese toxicity

The ability of plant for to grow and survive in a metal-contaminated environment, commonly called resistance, can be achieved through different mechanisms: avoidance and/or tolerance. The former involves a protective role that prevents the metal ions from entering the cytoplasm of plant cells (Blamey *et al.*, 1986; Marschner, 1991). The latter strategy (tolerance) implies a detoxification of metal ions after they have crossed the plasma cell membrane or internal organelle biomembranes (Macfie *et al.*, 1994). However, as shown in the following paragraphs, the differentiation of these mechanisms in the pertinent literature is very confusing.

The sequestering of Mn in the apoplast is considered an avoidance mechanism. However, some researchers have included this feature as a tolerance mechanism. For example, Horst *et al.*, (1999) suggested that tolerance to Mn excess in *Vigna unguiculata* is performed by the reduction of Mn<sup>2+</sup> activity in the apoplast throughout complexation by organic acids. In this species, symptoms such as brown leaf cell spots are also identified as oxidized Mn, and phenolic compounds present in the cell walls are considered a Mn tolerance mechanism (Wissemeier and Horst, 1992). On the other hand, the Mn<sup>2+</sup> oxidation by peroxidases in the cell walls of roots is considered by Marschner (1991) as an avoidance mechanism, although the existence of such a mechanism was not considered in the study of Horiguchi (1987). In this study, it is suggested that oxidized Mn deposition in plant tissues corresponds to a tolerance mechanism to Mn toxicity, with *Cucumis sativus* being more tolerant to high Mn deposition in tissues than melon (*Cucumis melo*). Blamey *et al.*, (1986) reported the accumulation and secretion of Mn<sup>2+</sup> in and around the trichomes of sunflower plants (*Helianthus annuus*) as a Mn tolerance mechanism. Another strategy that plants use to prevent the toxic effects of heavy metals as well as of Mn can be the efflux from the cell. In this process, the Mn cell is delivered into the Golgi apparatus and finally exported from the cell via secretory pathway vesicles that carry the metal to the cell surface (Ducic and Polle, 2005).

Summarizing the available literature regarding tolerance mechanisms to Mn, it appears that the main Mn tolerance mechanism is the sequestration by organic compounds in metabolically less active cells or organelles. The vacuole is considered the biggest and most important compartment, because it can store many toxic compounds (Pittman, 2005). Hence, an increase in phenolic compounds was found in the hydrophyte (*Trapa natans*) leaves exposed to high Mn

---

levels (130  $\mu\text{M}$ ) (Baldisserotto *et al.*, 2004). These compounds chelate Mn inside the vacuole, segregating the metal ion in the protoplasm and thus reducing the damage (Davis *et al.*, 2001). A similar key role has recently been assigned to oxalic acid in Mn internal sequestration by chelating specifically the Mn excess in vacuoles of Mn hyperaccumulator plants (*Phytolacca americana*) (Dou *et al.*, 2008). Furthermore, it has been observed that a Mn excess can accumulate dark material in the vacuoles, probably for deposition of Mn oxides or an increase in polyphenol oxidase activity in *Citrus volkameriana* plants (Papadakis *et al.*, 2007). Similarly, studies about the effect of Mn excess in varieties of the conifer *Pseudotsuga menziesii* showed dark deposits of Mn-complexes in plant root vacuoles, which were associated with phosphate, establishing “free”  $\text{Mn}^{2+}$  to form insoluble complexes, giving a greater tolerance (Ducic and Polle, 2007). Additionally, these studies showed that both, root elongation and biomass production were inhibited by Mn treatment above 2,500  $\mu\text{M}$ , mainly in *P. menziesii* var. *glauca*, confirming that it is a species with lower tolerance than *P. menziesii* var. *viridis*.

Another strategy to confer Mn tolerance inside plant cells is associated with several metal transporter proteins identified in the Mn transport mechanisms described in section 4. Metal transporter proteins located in the tonoplast (CAX2 in tobacco plants, ShMTP1 in *Arabidopsis* plants) conferred greater tolerance on elevated  $\text{Mn}^{2+}$  levels due to the internal sequestering of this element (Hirschi *et al.*, 2000; Delhaize *et al.*, 2003). Other transporter proteins (ECA1) can maintain low cytosolic Mn, since it moves into the endoplasmic reticulum (Wu *et al.*, 2002). In this research, ECA1 was able to reduce cytosolic  $\text{Mn}^{2+}$ , preventing an interference with the internal distribution of other ions ( $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$  or  $\text{Ca}^{2+}$ ).

Unlike other metal stresses, the accumulation of Mn excess does not have a single cell target. Depending on the plant species, different organelles can serve as stores for this accumulation (Lidon *et al.*, 2004). Under high Mn levels apart from vacuoles, chloroplasts are important sinks of this metal in *Citrus volkameriana*. This feature, together with the larger size of this organelle, is considered to be an adaptive response of this plant to Mn excess (Papadakis *et al.*, 2007). In general, cultivated plants like rice are considered tolerant to Mn toxicity (Lidon, 2001) because their leaf tissues can accumulate from 5 to 10 times more Mn than other grasses (Foy *et al.*, 1978). The Mn tolerance mechanism in this species included the inhibition of apoplastic influx

---

from the cortex toward the stele and symplastic Mn assimilation in the shoot protoplast, where the chloroplast is the main target (Lidon, 2001).

The distribution of Mn excess in both roots and shoots is dependent on plant species and genotype. Early research associated Mn tolerance in some plants with a greater retention of Mn excess in the roots, as mentioned by Andrew and Hegarty (1969) in regard to tropical and temperate legume species. The root retention of heavy metals has been attributed to the formation of metal complexes in roots (Foy *et al.*, 1978). Metals with high electro-negativity accumulate in roots in larger amounts than metals with low electro-negativity. In the latter instance, Mn and Zn metals can easily be translocated to the tops (Chino, 1981).

In proteomics studies, a comparison of Mn-sensitive and Mn-tolerant cultivars of cowpea (*Vigna unguiculata*.) has shown relevant features of leaf apoplast in the expression of Mn toxicity: formation of brown spots, induction of callose formation and an enhanced release of phenols and peroxidases into the apoplast (Fecht-Christoffers *et al.*, 2003; Fecht-Christoffers *et al.*, 2006). Specific proteins involved in the regulation processes, such as CO<sub>2</sub> fixation, stabilization of the Mn cluster of the photosystem II, pathogenesis-response reactions and protein degradation, were affected at low or high Mn levels, mainly in the Mn-sensitive cowpea cultivar (Führs *et al.*, 2008). Chloroplastic proteins, which are important for CO<sub>2</sub> fixation and photosynthesis, were of lower abundance upon Mn-induced stress, suggesting the scavenging of metabolic energy for a specific stress response. Führs *et al.*, (2008) concluded that a coordinated interplay of apoplastic and symplastic reactions seems to be important during the Mn-stress response in plants.

To alleviate metal toxicity in plants, the antioxidant systems are also considered an important tolerance mechanism. The antioxidant systems include antioxidant enzyme “scavengers” such as superoxide dismutase (SOD), catalase (CAT), peroxidases (phenol peroxidase, POX, ascorbate peroxidase, APX, guaiacol peroxidase, GPX) and the non-enzymatic antioxidant molecules: ascorbate,  $\alpha$ -tocopherol, carotenoids, flavonoids and glutathione (Foyer and Noctor, 2003; Apel and Hirt, 2004). These constitute other mechanisms against ROS, produced by Mn toxicity.

Higher activities of antioxidant enzymes are found in response to a Mn excess in woody plants (Lei *et al.*, 2007), in herbs such as white clover (*Trifolium repens*.) and in ryegrass (*Lolium perenne*), which suggest a lower oxidative stress (Rosas *et al.*, 2007; Mora *et al.*, 2009). Another

---

plant tolerance mechanism to metal toxicity is associated with lower metal uptake and translocation to other organs (Hall, 2002). The exudation of organic acid anions (carboxylates) to the rhizosphere may minimize the absorption by roots of such metals as aluminium and nickel (Ma *et al.*, 2001; Yang *et al.*, 1997). There are few reports regarding the release of organic acid anions by roots in the event of Mn toxicity (González and Lynch, 1999). Recently, Mora *et al.* (2009) reported that the root exudates of oxalate and citrate may decrease Mn availability in the rhizosphere, enhancing their Mn tolerance in ryegrass subjected to Mn toxicity.

Other nutrient applications can help minimize the effects of Mn toxicity. Thus, it has been reported that silica (Si) addition significantly decreases lipid peroxidation caused by an Mn excess decreasing the symptoms of Mn phytotoxicity and improving plant growth in some plants (Iwasaki *et al.*, 2002; Shi *et al.*, 2005). There is also an association between Mn toxicity and the decrease in Ca concentration in barley plants (*Hordeum vulgare*), indicating a competition and a specific interaction during the absorption and/or translocation of these elements (Alam *et al.*, 2001). Another study, in which Ca is applied to reduce the Mn toxicity, showed that Ca additions inhibited Mn translocation from roots to shoots in barley plants, but did not affect the Mn absorption in roots (Alam *et al.*, 2006). This suggested that Ca could avoid Mn accumulation in shoots, protecting the photosynthetic apparatus from the dangerous effect of an excess of Mn. In barley, Mn toxicity could also be repressed by high K contents, which inhibit both the absorption and the translocation of Mn (Alam *et al.*, 2005).

## **2.7 Conclusions and perspectives**

Manganese is considered an essential micronutrient for the metabolic process in plants. Nevertheless, both deficiency and excess alter these processes. Acid soils make excessive Mn amounts toxic for the plants. Mn toxicity is a world-wide problem in areas with acid soils. This toxicity alters physiological, biochemical and molecular processes at the cell level. It is crucial to know the limitations of these soils for the purpose of soil-plant interaction management, especially in relation to the presence and Mn excess. Thus, the knowledge of Mn uptake, translocation, accumulation and resistance mechanisms in crop plants under Mn excess and toxicity is of great importance to crop improvement.

---

Most molecular and physiological approaches to Mn transport inside plant cells have recently been analyzed, as these are useful tools for understanding resistance mechanisms. However, the question as to which is the best candidate gene related to Mn toxicity in acid soil continues to elude plant scientists due to Mn toxicity in plants being a complex trait and involving multiple physiological and biochemical mechanisms and a wide array of genes. The understanding of these mechanisms will contribute to improving the yield and quality of cultivated plants in acid soils. Future efforts for developing Mn-tolerant plants should take all these aspects into account.

#### Acknowledgments

Financial support for this review was provided by Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT N° 1080372) and Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) Doctoral Fellowship of the Government of Chile.

---

## 2.8 References

- Abou, M., Symeonidis, L., Hatzistavrou, E., Yupsanis, T. 2002. Nucleolytic activities and appearance of a new DNase in relation to nickel and manganese accumulation in *Alyssum murale*. J. Plant Physiol. 159, 1087-1095.
- Adriano, D.C. 2001. Trace elements in terrestrial environments. Biogeochemistry, bioavailability and risks of metals. Springer-Verlag, New York, 867 p.
- Alam, S., Kamei, S., Kawai, S. 2001. Amelioration of manganese toxicity in barley with iron. J. Plant Nutr. 28, 1421-1433.
- Alam, S., Akiha, F., Kamei, S., Huq, S., Kawai, S. 2005. Mechanism of potassium alleviation of manganese phytotoxicity in barley. J. Plant Nutr. 28, 889-901.
- Alam, S., Kodama, R., Akiha, F., Kamei, S., Kawai, S. 2006. Alleviation of manganese phytotoxicity in barley with calcium. J. Plant Nutr. 29, 59-74.
- Alscher, R., Erturk, N., Heath, L. 2002. Role of superoxide dismutase (SODs) in controlling oxidative stress in plants. J. Exp. Bot. 53, 1331-1341.
- Andrew, C.S., Hegarty, M.P. 1969. Comparative responses to manganese excess of eight tropical and four temperate pasture legume species. Aust. J. Agric. Res. 20, 687-696.
- Amao, Y., Ohashi, A. 2008. Effect of Mn ion on the visible light induced water oxidation activity of photosynthetic organ grana from spinach. Catal. Commun. 10, 217-220.
- Apel, K., Hirt, H. 2004. Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. Annu. Rev Plant Biol. 55, 373-379.
- Baldisserotto, C., Ferroni, L., Medici, V., Pagnoni, A., Pellizzari, M., Fasulo, M.P., Fagioli, F., Bonora, A., Pancaldi, S. 2004. Specific intra-tissue responses to manganese in the floating lamina of *Trapa natans* L. Plant Biol. 6, 578-589.
- Bartsevich, V., Pakrasi, H.B. 1996. Manganese transport in the cyanobacterium *Synechocystis* sp PCC 6803. J. Biol. Chem. 271, 26057-26061.
- Blamey, F., Joyce, D., Edwards, D., Asher, C. 1986. Role of trichomes in sunflower tolerance to manganese toxicity. Plant Soil. 91, 171-180.
- Bradl, H. 2004. Adsorption of heavy metal ions on soils and soils constituents. J. Colloid. Interf. Sci. 277, 1-18.
- Buchanan, B., Grusen, W., Jones, R. 2000. Biochemistry and molecular biology of plants. American Society. of Plant Physiologists. Maryland. 1367 p.

- 
- Burnell, J. 1988. The biochemistry of manganese in plants. In: R.D. Graham, R.J. Hannam, N.J. Uren. (eds). *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 125–137.
- Chino, M. 1981. Metal stress in rice plants. In: K. Kitagisi, I. Yamane (eds). *Heavy metal pollution in soils of Japan*. Japan Scientific Societies Press, Tokyo, pp. 65-80.
- Clarkson, D.T. 1988. The uptake and translocation of manganese by plant roots. In: R.D. Graham, R.J. Hannam, N.J. Uren. (eds). *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 101-111.
- Clemens, K.L., Force, D.A., Britt, R.D. 2002. Acetate binding at the photosystem II oxygen evolving complex: an S<sub>2</sub>-State multiline signal ESEEM study. *J. Am. Chem. Soc.* 124, 10921-10933.
- Conyers, M., Uren, N., Helyar, K., Poile, G., Cullis, B. 1997. Temporal variation in soil acidity. *Aust. J. Soil Res.* 35, 1115-1129.
- Davis, M.A., Pritchard, S.G., Boyd, R.S., Prior, S.A. 2001. Developmental and induced responses of nickel-based and organic defenses of the nickel-hyperaccumulating shrub, *Psychotria douarrei*. *New. Phytol.* 150, 49-58.
- Delhaize, E., Kataoka, T., Hebb, D.M., White, R.G., Ryan, P.R. 2003. Genes encoding proteins of the cation diffusion facilitator family that confer manganese tolerance. *Plant Cell.* 15, 1131- 1142.
- Delhaize, E., Gruber, B.D., Pittman, J.K., White, R.G., Leung, H., Miao, Y., Jiang, L., Ryan, P.R., Richardson, A.E. 2007. A role for the AtMTP11 gene of *Arabidopsis* in manganese transport and tolerance. *Plant J.* 51, 198–210.
- Demirevska-Kepova, K., Simova-Stoilova, L., Stoyanova, Z., Holzer, R., Feller, U. 2004. Biochemical changes in barley plants after excessive supply of copper and manganese. *Environ. Exp. Bot.* 52, 253-266.
- Dou, C., Fu, X, Chen, X, Shi, J., Chen, Y. 2008. Accumulation and detoxification of manganese in hyperaccumulator *Phytolacca americana*. *Plant Biol.* 11, 664-670.
- Ducic, T., Polle, A. 2005. Transport and detoxification of manganese and copper in plants. *Braz. J. Plant Physiol.* 17, 103-112.
- Ducic, T., Polle, A. 2007. Manganese toxicity in two varieties of Douglas fir (*Pseudotsuga menziesii* var. *viridis* and *glauca*) seedlings as affected by phosphorus supply. *Funct. Plant Biol.* 34, 31-40.
- El-Jaoual, T., Cox, D. 1998. Manganese toxicity in plants. *J. Plant Nutr.* 21, 353-386.

- 
- Fageria, N., Baligar, V., Clark, R. 2002. Micronutrients in crop production. *Adv. Agron.* 77, 185-268.
- Farasova, A., Beinrohr, E. 1998. Metal-metal interaction in accumulation of V, Ni, Mo, Mn, and Cu in under and above ground parts of *Sinapis alba*. *Chemosphere* 36, 1305-1317.
- Fecht-Christoffers, M., Braun, H., Lemaitre-Guillier, C., VanDorsselaer, A., Horst, W. 2003. Effect of manganese toxicity on the proteome of the leaf apoplast in cowpea. *Plant Physiol.* 133, 1935-1946.
- Fecht-Christoffers, M., Führs, H., Braun, H., Horst, W. 2006. The role of hydrogen peroxide-producing and hydrogen peroxide-consuming peroxidases in the leaf apoplast of cowpea in manganese tolerance. *Plant Physiol.* 140, 1451-1463.
- Foy, C., Chaney, R., White, M. 1978. The physiology of metal toxicity in plants. *Ann. Rev. Plant Physiol.* 29, 511-566.
- Foy, C., Scott, B., Fisher, J. 1988. Genetic differences in plant tolerance to manganese toxicity. In: R.D. Graham, R.J. Hannam, N.J. Uren. (eds). *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 293–307.
- Foyer, C., Noctor, G. 2003. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiol. Plant.* 119, 355-364.
- Führs, H., Hartwig, M., Buitrago, L., Heintz, D., Van Dorsselaer, A., Braun, H., Horst, W. 2008. Early manganese-toxicity response in *Vigna unguiculata* L.– a proteomic and transcriptomic study. *Proteomics* 8, 149-159.
- Führs, H., Götze, S., Specht, A., Erban, A., Gallien, S., Heintz, D., Van Dorsselaer, A., Kopka, J., Braun, H.P., Horst, W.J. 2009. Characterization of leaf apoplastic peroxidases and metabolites in *Vigna unguiculata* in response to toxic manganese supply and silicon. *J. Exp. Bot.* 60, 1663–1678.
- Gherardi M., Rengel, Z. 2003. Genotypes of lucerne (*Medicago sativa* L.) show differential tolerance to manganese deficiency and toxicity when grown in bauxite residue sand. *Plant Soil* 249, 287-296.
- Gherardi M., Rengel, Z. 2004. The effect of manganese supply on exudation of carboxylates by roots of lucerne (*Medicago sativa*). *Plant Soil* 260, 271-282.
- González, A., Steffen, K., Lynch, J. 1998. Light and excess manganese. *Plant Physiol.* 118, 493-504.
- González, A., Lynch, J. 1999. Subcellular and tissue Mn compartmentation in bean leaf under Mn toxicity stress. *Aust. J. Plant Physiol.* 26, 811-822.

- 
- Goussias, C., Boussac, A., Rutherford, W. 2002. Photosystem II and photosynthetic oxidation of water: an overview. *Phil. Trans. R. Soc. Lond. B* 357, 1369-1381.
- Guest, C., Schulze, D., Thompson, I., Huber, D. 2002. Correlating manganese X-ray absorption near-edge structure spectra with extractable soil manganese. *Soil Sci. Soc. Am. J.* 66, 1172-1181.
- Hall, J.L. 2002. Cellular mechanisms for heavy metal detoxification and tolerance. *J. Exp. Bot.* 53, 1-11.
- Hauck, M., Paul, A., Gross, S., Raubuch, M. 2003. Manganese toxicity in epiphytic lichens: chlorophyll degradation and interaction with iron and phosphorus. *Environ. Exp. Bot.* 49, 181-191.
- Hegedüs, A., Erdei, S., Horváth, G. 2001. Comparative studies of H<sub>2</sub>O<sub>2</sub> detoxifying enzymes in green and greening barley seedlings under cadmium stress. *Plant Sci.* 160, 1085-1093.
- Henriques, F. 2003. Gas exchange, chlorophyll a fluorescence kinetics and lipid peroxidation of pecan leaves with varying manganese concentrations. *Plant Sci.* 165, 239-244.
- Herren, T., Feller U. 1994. Transfer of zinc from xylem to phloem in the peduncle of wheat, *J. Plant Nutr.* 17, 1587-1598.
- Hirschi, K., Korenkov, V., Wilganowski, N., Wagner, G.J. 2000. Expression of *Arabidopsis* CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. *Plant Physiol.* 124, 125-133.
- Horiguchi, T. 1987. Mechanism of manganese toxicity and tolerance of plants. II. Deposition of oxidized manganese in plant tissues. *J. Soil Sci. Plant Nutr.* 33, 595-606.
- Horst, W.J. 1988. The physiology of manganese toxicity. In: R.D. Graham, R.J. Hannam, N.J. Uren (eds). *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 175-188.
- Horst, W., Fecht, M., Naumann, A., Wissemeier, A., Maier, P. 1999. Physiology of manganese toxicity and tolerance in *Vigna unguiculata* (L.) Walp. *J. Plant Nutr. Soil Sci.* 162, 263-274.
- Houtz, R.L., Nable, R.O., Cheniae, G.M. 1988. Evidence for effects on the in vivo activity of ribulose-biphosphate carboxylase/oxygenase during development of Mn toxicity in tobacco. *Plant Physiol.* 86, 1143-1149.
- Hue, N. 1988. A possible mechanism for manganese toxicity in Hawaii soils amended with a low-Mn sewage sludge. *J. Environ. Qual.* 17, 473-479.

- 
- Hue, N., Mai, Y. 2002. Manganese toxicity in watermelon as affected by lime and compost amended to a Hawaiian acid Oxisol. *Hortscience* 37, 656-661.
- Humphries, J., Stangoulis, J., Graham, R. 2007. Manganese. In: A. Barker, D. Pilbeam (eds). *Handbook of Plant Nutrition*, Taylor and Francis, USA, pp. 351-366.
- Iwasaki, K., Maier, P., Fecht, M., Horst, W. 2002. Effects of silicon supply on apoplastic manganese concentrations in leaves and their relation to manganese tolerance in cowpea (*Vigna unguiculata* (L.)Walp). *Plant. Soil* 238, 281-288.
- Kochian, L., Hoekenga, O., Piñeros, M. 2004. How do crops plants tolerate acid soils? Mechanisms of aluminum tolerance and phosphorus efficiency. *Annu. Rev. Plant Biol.* 55, 459-93.
- Kogelmann, W., Sharpe, W. 2006. Soil acidity and manganese in declining and nondeclining sugar maple stands in Pennsylvania. *J. Environ. Qual.* 35, 433-441.
- Korshunova, Y., Eide, D., Clark, W., Guerinot, M., Pakrasi, H. 1999. The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range. *Plant Mol. Biol.* 40, 37-44.
- Laha, S., Luthy, R. 1990. Oxidation of aniline and other primary aromatic amines by manganese dioxide. *Environ. Sci. Technol.* 24, 363-373.
- Lavres J., Malavolta, E., Nogueira, N.L., Moraes, M.F., Rodrigues, A., Lanzoni, M., Pereira, C. 2009. Changes in anatomy and root cell ultrastructure of Soybean genotypes under manganese stress. *R. Bras. Ci. Solo* 33,395-403.
- Lei, Y., Korpelainen, H., Li, C. 2007. Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. *Chemosphere* 68, 686-694.
- Lidon, F. 2001. Tolerance of rice to excess manganese in the early stages of vegetative growth. Characterization of manganese accumulation. *J. Plant Physiol.* 158, 1341-1348.
- Lidon, F.C., Barreiro, M., Ramalho, J. 2004. Manganese accumulation in rice: implications for photosynthetic functioning. *J. Plant Physiol.* 161, 1235-1244.
- Lidon, F.C, Teixeira, M.G. 2000a. Rice tolerance to excess Mn: Implications in the chloroplast lamellae and synthesis of a novel Mn protein. *Plant Physiol. Biochem.* 38, 969-978.
- Lidon, F.C., Teixeira, M.G. 2000b. Oxy radicals production and control in the chloroplast of Mn-treated rice. *Plant Sci.*152, 7-15.

- 
- Lidon, F.C., Henriques, F. 1993. Oxygen metabolism in higher plant chloroplasts. *Photosynthetica* 29, 249-279.
- Luk, E., Jensen, L.T., Culotta, V.C. 2003. The many highways for intracellular trafficking of metals. *JBIC* 8, 803-809.
- Lynch, J.P., St.Clair, S.B. 2004. Mineral stress: the missing link in understanding how global climate change will affect plants in real world soils. *Field Crop Res.* 90, 101–115.
- Ma, J.F. Ryan, P. R., Delhaize, E. 2001. Aluminium tolerance in plants and the complexing role of organic acids. *Trends Plant Sci.* 6, 273-278.
- Maas E., Moore D. 1968. Manganese absorption by excised barley roots. *Plant Physiol.* 43, 527-530.
- McCain, D.C., Markley, J:L. 1989. More manganese accumulates in maple sun leaves than in shade leaves. *Plant Physiol.* 90: 1417–1421.
- Macfie, S., Taylor, G. 1992. The effects of excess manganese on photosynthetic rate and concentration of chlorophyll in *Triticum aestivum* grown in solution culture. *Physiol. Plant.* 85, 467-475.
- Macfie, S., Cossins, E., Taylor, G. 1994. Effects of excess manganese on production of organic acids in Mn-tolerant and in Mn-sensitive cultivars of *Triticum aestivum* L. (wheat). *J. Plant Physiol.* 143, 135-144.
- Marschner, H. 1991. Mechanisms of adaptation of plants to acid soils. In: R. Wright, V. Baligar, R. Murrmann (Eds). *Development in Plant and Soil Sciences. Plant-soil interactions at low pH.* Kluwer Academic Publisher, West Virginia, pp. 1-20.
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants.* Academic Press, San Diego, 889 p.
- Martinoia, E., Klein, M., Geisler, M., Bovet, L., Forestier, C., Kolukisaoglu, H., Mueller-Roeber, B., Schulz, B. 2002. Multifunctionality of plant ABC transporters: more than just detoxifiers. *Planta* 214, 345-355.
- Migocka M., Klobus G. 2007. The properties of the Mn, Ni and Pb transport operating at plasma membranes of cucumber roots. *Physiol. Plant.* 129, 578-587.
- Mills, R., Doherty, M., López-Marqués, R., Weimar, T., Dupree, P., Palmgren, M., Pittman, J., Williams, L. 2008. ECA3, a Golgi-localized P2A-type ATPase, plays a crucial role in manganese nutrition in *Arabidopsis*. *Plant Physiol.* 146, 116-128.
- Miner, G., Sims, J. 1983. Changing fertilization practices and utilization of added plant nutrient for efficient production of tobacco. *Recent Adv. Tobacco Sci.* 9, 4-76.

- 
- Mitsukawa, N., Okumura, S., Shirano, Y., Sato, S., Kato, T., Harashima, S., Shibata, D. 1997. Overexpression of an *Arabidopsis thaliana* high-affinity phosphate transporter gene in tobacco cultured cells enhances cell growth under phosphate-limited conditions. *Proc. Natl. Acad. Sci.* 94, 7098-7102.
- Mora, M., Alfaro, M., Jarvis, S., Demanet, R., Cartes, P. 2006. Soil aluminium availability in Andisols of southern Chile and its effect on forage production and animal metabolism. *Soil Use Manage.* 22, 95-101.
- Mora, M., Rosas, A., Ribera, A., Rengel, R. 2009. Differential tolerance to Mn toxicity in perennial ryegrass genotypes: involvement of antioxidative enzymes and root exudation of carboxylates. *Plant Soil* 320, 79–89.
- Moroni, J., Scott, B., Wratten, N. 2003. Differential tolerance of high manganese among rapeseed genotypes. *Plant Soil* 253, 507-519.
- Mukhopadhyay M., Sharma, A. 1991. Manganese in cell metabolism of higher plants. *Bot. Rev.* 57, 117-149.
- Nable, R., Houtz, R., Cheniae, G. 1988. Early inhibition of photosynthesis during development of Mn toxicity in tobacco. *Plant Physiol.* 86, 1136-1142.
- Najeeb, U., Xu, L., Shafaqat, A., Jilani, G., Gong, H.J., Shen, W.Q., Zhou, W.J. 2009. Citric acid enhances the phytoextraction of manganese and plant growth by alleviating the ultrastructural damages in *Juncus effusus* L. *J. Hazard. Mater.* 170, 1156-1163.
- Ness, P.J., Woolhouse, H.W. 1980. RNA synthesis in *Phaseolus* chloroplasts. 1. Ribonucleic acid synthesis and senescing leaves. *J. Exp. Bot.* 31, 223-233.
- Neumann, G., V., Romheld. 2001. The release of root exudates as affected by the plants physiological status. In: R. Pinto, Z. Varanini, P. Nannipieri (eds). *The rhizosphere: Biochemistry and organic substances at the soil-plant interface*, Marcel Dekker. New York, pp. 41–93.
- Page, V., Feller, U. 2005. Selective transport of zinc, manganese, nickel, cobalt and cadmium in the root system and transfer to the leaves in young wheat plants. *Ann. Bot.* 96, 425-434.
- Page, V., Weisskopf, L., Feller, U. 2006. Heavy metals in white lupin: uptake, root-to-shoot transfer and redistribution within the plant. *New Phytol.* 171, 329-341.
- Panda, S., Mishra, A.K., Biswal, U.C. 1986. Manganese-induced modification of membrane lipid peroxidation during aging of isolated wheat chloroplasts. *Photobiochem. Photobiophys* 13, 53-61.

- 
- Papadakis, I., Giannakoula, A., Therios, I., Bosabalidis, A., Moustakas, M., Nastou, A. 2007. Mn-induced changes in leaf structure and chloroplast ultrastructure of *Citrus volkameriana* (L.) plants. *J. Plant Physiol.* 164, 100-103.
- Pendias, K., Pendias, H. 1992. Trace elements in soils and plants. USA, CRR Press, 365 p.
- Pfeffer, P.E., Tu, S., Gerasimowicz, W.V., Cavanaugh, J.R. 1986. In vivo <sup>31</sup>P NMR studies of corn root tissue and its uptake of toxic metals. *Plant Physiol.* 80, 77-84.
- Pittman, J. 2005. Managing the manganese: molecular mechanisms of manganese transport and homeostasis. *New Phytol.* 167, 733-742.
- Polle, A. 2001. Dissecting the superoxide dismutase-ascorbate-glutathione-pathway in chloroplasts by metabolic modeling. Computer simulations as a step towards flux analysis. *Plant Physiol.* 126, 445-462.
- Porter, G., Bajita-Locke, J., Hue, N., Strand, S. 2004. Manganese solubility and phytotoxicity affected by soil moisture, oxygen levels, and green manure additions. *Commun. Soil Sci. Plant. Anal.* 35, 99-116.
- Reddi, A.R., Jensen, L.T., Culotta, V.C. 2009. Manganese homeostasis in *Saccharomyces cerevisiae*. *Chem. Rev.* 109, 4722-4732.
- Rengel, Z. 2000. Uptake and transport of manganese in plants. In: A. Sigel, H. Sigel (eds.). *Metal Ions in Biological Systems*. New York, Marcel Dekker, pp. 57-87.
- Rengel, Z. 2001. Xylem and phloem transport of micronutrients. In: W. J. Horst et al.(eds). *Plant nutrition – Food security and sustainability of agro-ecosystems*. Kluwer Academic Publishers, Netherlands, pp. 628-629.
- Rengel, Z., Marschner, P. 2005. Nutrient availability and management in the rhizosphere: exploiting genotypic differences. *New Phytol.* 168, 305-312.
- Roberts, L., Pierson, A., Panaviene, Z., Walker, E. 2004. Yellow Stripe 1. Expanded roles for the maize iron-phytosiderophore transporter. *Plant Physiol.* 135, 112-120.
- Rosas, A., Rengel, Z., Mora, M. 2007. Manganese supply and pH influence growth, carboxylate exudation and peroxidase activity of ryegrass and white clover. *J. Plant Nutr.* 30, 253-270.
- Ryan, P., Delhaize, E., Jones, D. 2001. Function and mechanism of organic anion exudation from plant roots. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52, 527-560.
- Sarkar, D., Pandey, S., Sud, K., Chanemougasoundharam, A. 2004. In vitro characterization of manganese toxicity in relation to phosphorus nutrition in potato (*Solanum tuberosum* L.). *Plant Sci.* 167, 977-986.

- 
- Shi, Q., Bao, Z., Zhu, Z., He, Y., Qian, Q., Yu, J. 2005. Silicon-mediated alleviation of Mn toxicity in *Cucumis sativus* in relation to activities of superoxide dismutase and ascorbate peroxidase. *Phytochemistry* 66, 1551-1559.
- Shi, Q., Zhu, Z., Xu, M., Qian, Q., Yu, J. 2006. Effect of excess manganese on the antioxidant system in *Cucumis sativus* L. under two light intensities. *Environ. Exp. Bot.* 58, 197-205.
- Shigaki, T., Pittman, J., Hirschi, K. 2003. Manganese specificity determinants in *Arabidopsis* metal/H<sup>+</sup> antiporter CAX2. *J. Biol Chem.* 21, 6610-6617.
- Sparks, D. 1995. *Environmental Soil Chemistry*. San Diego, Academic Press, 352 p.
- Sparrow, L., Uren, N. 1987. Oxidation and reduction of Mn in acidic soils: effect of temperature and soil pH. *Soil Biol. Biochem.* 19, 143-148.
- Takahashi, M., Sugiura, M. 2001. Strategies for uptake of a soil micronutrient, manganese, by plant roots. *RIKEN Review* 35, 76-77.
- Tsuji, M., Mori, H., Yamamoto, T., Tanaka, S., Kang, Y., Sakurai, K., Iwasaki, K. 2006. Manganese toxicity of melon plants growing on an isolated soil bed after steam sterilization. *J. Soil Sci. Plant Nutr.* 77, 257-263.
- Thomine, S., Lelievre, F., Debarbieux, E., Schroeder, J., Barbier-Brygoo, H. 2003. AtNRAMP3, a multispecific vacuolar metal transporter involved in plant responses to iron deficiency. *Plant J.* 34, 685-695.
- Wang, Y., Wisniewski, M., Meilan, R., Uratsu, S., Cui, M., Dandekar, A., Fuchigami, L. 2007. Ectopic expression of Mn-SOD in *Lycopersicon esculentum* leads to enhanced tolerance to salt and oxidative stress. *J. Applied Hortic.* 9, 3-8.
- Wang, Y., Ying, Y., Chen, J., Wang, X. 2004. Transgenic *Arabidopsis* overexpressing Mn-SOD enhanced salt-tolerance. *Plant Sci.* 167, 671-677.
- Watmough, S., Eimer, M., Dillon, P. 2007. Manganese cycling in central Ontario forests: Response to soil acidification. *Appl. Geochem.* 22, 1241-1247.
- Wissemeier, A., Horst, W. 1991. Simplified methods for screening cowpea cultivars for manganese leaf-tissue tolerance. *Crop Sci.* 31, 435-439.
- Wissemeier A., Horst, W. 1992. Effect of light intensity on manganese toxicity symptoms and callose formation in cowpea (*Vigna unguiculata* (L.) Walp.). *Plant Sci.* 143, 299-309.
- Wu, Z., Liang, F., Hong, B., Young, J.C., Sussman, M.R., Harper, J.F., Sze, H. 2002. An endoplasmic reticulum-bound Ca<sup>2+</sup>/Mn<sup>2+</sup> pump, ECA1, supports plant growth and confers tolerance to Mn<sup>2+</sup> stress. *Plant Physiol.* 130, 128-137.

---

Yang, X.E., Baligar, V.C., Foster, J.C., Martens, D.C. 1997. Accumulation and transport of nickel in relation to organic acids in ryegrass and maize grown with different nickel levels. *Plant Soil* 196: 271–276.

Zouni, A., Witt, H., Kern, J., Fromme, P., Kraub, N., Saenger, W., Orth, P. 2001. Crystal structure of photosystem II from *Synechococcus elongatus* at 3.8 Å resolution. *Nature* 409, 739-743.

---

Chapter 3:

Effect of Mn excess on biochemical and physiological properties of Highbush blueberry (*Vaccinium corymbosum* L.) cultivars growing in nutrient solution.

Paper in preparation

---

**Effect of Mn excess on biochemical and physiological properties of Highbush blueberry (*Vaccinium corymbosum* L.) cultivars growing in nutrient solution.**

**Millaleo R.<sup>1</sup>, Alvear M.<sup>2,3</sup>, Aguilera P.<sup>1</sup>, Reyes-Díaz M.<sup>2,3</sup>, Mora M.<sup>2,3</sup>, Alberdi M.<sup>2,3</sup>**

<sup>1</sup>Doctorado en Ciencias de Recursos Naturales, Facultad de Ingeniería, Ciencias y Administración, Universidad de La Frontera, Temuco, Chile.

<sup>2</sup>Departamento de Ciencias Químicas y Recursos Naturales, Facultad de Ingeniería, Ciencias y Administración, Universidad de La Frontera, Temuco-Chile.

<sup>3</sup>Center of Plant, Soil Interaction, and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Casilla 54-D, Temuco, Chile.

---

## Abstract

Manganese (Mn) is an essential micronutrient for plants. However, when it is in excess, it can lead to negative effects on physiological and biochemical characteristics in plants. Besides, plant species had different resistant mechanisms against abiotic stress such as excess of metals. One year old plants of blueberry cultivars were grown in hydroponic solution with 2, 10, 50, 250, 500, and 1000  $\mu\text{M}$  of Mn treatments for 25 days. We compared the manganese (Mn) excess in three highbush blueberry (*Vaccinium corymbosum* L.) cultivars: Legacy, Brigitta and Bluegold. Parameters evaluated were: plant growth as shoot and root biomass, Mn content, photosynthetic performance (photochemical efficiency of PSII,  $\text{CO}_2$  assimilation and stomatal conductance). Lipid peroxidation, radical scavenging activity, superoxide dismutase activity and organic acid anions exudations, were also analysed. Results showed detrimental effects on plant growth, photochemical efficiency and  $\text{CO}_2$  assimilation, as well as in biochemical features. Legacy showed a major Mn content in roots compared with other cultivars, evidencing no changes in photosynthesis and in antioxidant capacity. Bluegold presented lower levels in net photosynthesis and photochemical processes and an increase of lipid peroxidation. In organic acid anions exudation, oxalate and citrate had greater level of exudation in Legacy and Brigitta, which suggest a major Mn-tolerance in front to an excess of this trace element compared with Bluegold. In conclusion, Legacy cultivar showed a higher Mn-tolerance than Brigitta and Bluegold, being this last cultivar the most sensitive.

---

### 3.1 Introduction

Manganese (Mn) is an essential micronutrient for most microorganisms and plants, being indispensable for many physiological and biochemical processes such as growth, photosynthesis, photosynthetic proteins and enzymes among others (Marschner, 1995). Depending on its soil content and the factors controlling its availability as low pH (< 5) and redox conditions Mn can be toxic for plants (Rengel 2000; Millaleo et al 2010). A decreasing pH leads to an increment of available toxic manganese ( $Mn^{2+}$ ) in the soil solution which can be readily transported into the root cells and translocated to the shoots, where it is accumulated (Marschner, 1995).

Manganese toxicity is one of the most important factors limiting plant growth, especially in tropics and subtropics areas (Horst, 1988). Many studies showed that under Mn excess physiological and biochemical plant functions, like growth (Lei et al., 2007), and photosynthetic performance including photosynthetic pigments are impaired (Hauck et al., 2002, 2003, 2006; Macfie y Taylor, 1992). In addition, under Mn toxicity reactive oxygen species (ROS) can be exacerbated conducting to oxidative stress and lipid peroxidation membranes as reported in barley, clover, and perennial ryegrass (Demirevska-Kepova et al., 2004; Rosas et al. 2007, Mora et al. 2009, respectively). To overcome with Mn induced oxidative stress plants have developed some mechanisms like an increase of antioxidant enzymes activity [superoxide dismutase (SOD) and peroxidase (POD)] as reported in cucumber (Shi et al., 2006), perennial ryegrass (Mora et al., 2009) and pea (Gangwar et al., 2010) exposed to a Mn excess. The exudation of organic acid anions (carboxylates) by apical roots is another important mechanism to counteract the deleterious effects of toxic cations. Carboxylates are able to reduce the concentration of toxic cations in the rizosphere or accumulating them in the cytoplasm of plant cells (Ryan et al., 2001). In ryegrass and white clover (Rosas et al., 2007) showed that toxic Mn is excluded from the roots by the exudates (exclusion mechanism) and kept out of it, forming soluble or insoluble complexes that decrease Mn availability (Wang and Mulligan, 2009). In ryegrass seedlings, Mora et al., (2009) found that oxalate and citrate were the most abundant carboxylates exudates induced by Mn excess, mainly in the most Mn resistant cultivars.

In southern Chile, acid soils of volcanic origin are common. In these soils a serious constraint for plant growth are toxic ( $Al^{3+}$ ) and toxic ( $Mn^{2+}$ ). In this region highbush blueberry (*Vaccinium corymbosum*) is one of the main cultivated crops. With exception of the work of Rojas-Lillo et al

---

(2014), who studied some physiological and biochemical responses to toxic Mn and UV-B on this crop, the root exudation of carboxylic acids under Mn excess and their association with antioxidant mechanisms are not available. Therefore, the aim of this work is to clarify the implication of the exudation of carboxylic acids and antioxidant responses under Mn excess in *V. corymbosum*.

### 3.2 Materials and Methods

#### *Plant material and growth conditions.*

Three cultivars of highbush blueberry (*Vaccinium corymbosum* L.): ‘Legacy’, ‘Brigitta’ and ‘Bluegold’, were used in this study. One-year old plants of these genotypes coming from a nursery of Temuco, Chile, growing in a substrate of sawdust:soil:pine needles at a 1:1:1 proportion were established under greenhouse in nutrient solution. For conditioning, seedlings were placed in plastic boxes with 10 L of Hoagland’s nutrient solution (Hoagland and Arnon, 1959) for 2 weeks. Later, blueberry seedlings were placed in a new Hoagland’s nutrient solution in plastic boxes (4 plants per box) and Mn treatments were applied. The time of exposition to Mn excess was 25 days. Blueberry plants were subjected to the following Mn concentrations (added as  $\text{MnCl}_2 \times 4\text{H}_2\text{O}$ ): 2  $\mu\text{M}$  (Control), 10, 50, 250, 500 and 1000  $\mu\text{M}$  Mn. The solutions were daily adjusted to  $\text{pH } 4.8 \pm 0.02$ , where volume was restored to its original level, and were renewed every 1 week. The Hoagland solution used included the following concentrations: 2 mM  $\text{Ca}(\text{NO}_3)_2$ , 3 mM  $\text{KNO}_3$ , 1 mM  $\text{MgSO}_4$ , 0,1 mM  $\text{KH}_2\text{PO}_4$ , 22  $\mu\text{M}$   $\text{H}_3\text{BO}_3$ , 2  $\mu\text{M}$   $\text{MnSO}_4$ , 1 mM  $\text{NH}_4\text{NO}_3$ , 0,07  $\mu\text{M}$   $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$ , 1,6  $\mu\text{M}$   $\text{ZnSO}_4$ , 0,4  $\mu\text{M}$   $\text{CuSO}_4$ , 20  $\mu\text{M}$  Fe- EDTA. Experimental conditions used were under semi-controlled environmental conditions in a naturally lit greenhouse, with a temperature range of 25/20 °C (day/night), relative humidity range of 50-80%, the photoperiod of 16/8 h (day/night), an average PPF (photosynthetic photon flux) of 300  $\mu\text{mol m}^{-2} \text{ s}^{-1}$ .

#### *Dry weight and Mn content measurements*

At the beginning of Mn treatments, the fresh weight of whole plants from each cultivar was determined. From these plants, a group was separated for dry weight determinations. At the end of the experiment (25 d), plants of blueberry cultivars were harvested and separated in roots and shoots, for fresh and dry weight determinations. Shoots and roots samples were dried in a forced-

---

air oven for 48 h at 65°C for dry weight determinations. To determine Mn concentrations, first leaves from the first to the fourth node of shoots and roots were dried at 65°C in a forced-air oven for 48 h. Later, samples were ashed in a muffle furnace at 500°C for 8 h and treated with 2M hydrochloric acid (HCl). Mn was quantified using a simultaneous multielement atomic absorption spectrophotometer (Model 969 atomic absorption spectrometer; Unicam, Cambridge, UK) according to Sadzawka et al. (2004) methodology. To obtain the Mn contents, the Mn concentration was multiplied with the respective dry weight for leaves and roots.

#### *Photosynthetic performance*

It was analyzed by measuring the photochemical efficiency of the PSII and net photosynthesis ( $P_n$ ). For the photochemical efficiency of PSII, the leaf chlorophyll fluorescence from the second to fourth node of shoots (previously dark adapted for 20 min) was measured using a portable pulse-amplitude modulated fluorimeter (FMS 2; Hansatech Instruments, King's Lynn, UK) according to (Reyes-Díaz et al. 2009). The maximal photochemical efficiency of PSII ( $F_v/F_m$ ), effective photochemical efficiency of PSII ( $\Phi_{PSII}$ ) as indicator of the effective quantum yield of PSII and non-photochemical quenching (NPQ) were estimated as described by Maxwell and Johnson (2000). The net photosynthesis were performed in attached leaves with similar localization as those used for the determination of fluorescence parameters. The measurement of net CO<sub>2</sub> assimilation was made using a portable CO<sub>2</sub> infrared gas analyzer (Licor LR6400). This analyzer was equipped with a fluorescence leaf cuvette, which controlled the light source (300  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ), temperature, humidity and CO<sub>2</sub> external air obtaining a concentration reference of 360 ppm, with a flow rate of 200 mL min<sup>-1</sup> and 80% external relative humidity. The temperature inside the leaf chamber was maintained at 20°C. Stomatal conductance ( $g_s$ ) was also given by the  $P_n$  determination.

#### *Lipid peroxidation and Radical scavenging activity*

The level of lipid peroxidation (LP) was assessed in fresh samples by monitoring the thiobarbituric acid reacting substances (TBARS) as an index of oxidative damage in plant cells. The absorbance was measured at 532, 600 and 440 nm in order to correct the interference generated by TBARS-sugar complexes according to the modified method (Du and Bramlage, 1992). This method determines the occurrence of malondialdehyde (MDA) as secondary

---

byproduct of the oxidation of polyunsaturated fatty acids that can react with thiobarbituric acid (TBA) (Hodges et al. 1999) The unit for LP was determined as equivalents of malondialdehyde (MDA) contents ( $\text{nmol g}^{-1}$  FW). The radical scavenging activity (RSA) of roots and leaves was measured at the end of Mn treatments (25d) using the method of free radical 1,1-diphenyl-2-picrylhydrazyl (DPPH) scavenging, as described by Chinnici et al. (2004), with some modifications. The absorbance was measured at 515 nm using Trolox as the standard. The unit used was Trolox equivalent per gram of fresh weight (TE [ $\text{mg g}^{-1}$  FW]).

*Superoxide dismutase (SOD) activity.* For determination of the superoxide dismutase (SOD, EC 1.15.1.1) enzyme activity, samples of fresh leaves and roots were frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until extraction of SOD. The extraction procedure was performed as described by Mora et al. (2009). SOD activity was assayed by monitoring the photochemical inhibition of nitroblue tetrazolium (NBT) as described by Giannopolitis and Ries (1977) with minor modifications (Mora et al., 2009). Enzymatic activity of SOD values were standardized by the total protein content, according to Bradford (1976). One unit of SOD activity ( $\text{U g}^{-1}$ ) was defined as the amount of enzyme required to cause 50% inhibition of the reduction of NBT as monitored at 560 nm.

#### *Analysis of root exudates.*

Root exudates were collected 25 d after the beginning of Mn treatments, according to Rosas et al. (2007). Intact blueberry plants coming from Mn treatments assays were rinsed carefully with deionized water and transferred to pots containing deionized water (250 mL). For collection of exudates from the whole root system, roots were submerged during 2 h into aerated deionized water and then, subsamples of control and Mn-treated plants were collected, stored and frozen at  $-20^{\circ}\text{C}$ . The organic acid anions (oxalate, malate, citrate and succinate) exudated from roots were quantified by chromatographic analysis. Root exudate subsamples were concentrated by lyophilisation, the residue re-dissolved in 600–800  $\mu\text{L}$  of deionized-sterile water for HPLC injection and filtered (0.22  $\mu\text{m}$ ). The chromatographic separation was realized in an HPLC system (Merck-Hitachi model L-4200) equipped with a UV–visible detector and a Sphere Column Heater (Phenomenex Terma model TS-130) according to Meier et al (2012). The flow rate was  $1 \text{ mL min}^{-1}$  and injected samples were detected at wavelength of 210 nm. Identification

---

of organic acids anions was determined comparing the retention times and by addition of standards for each organic acid anion.

*Experimental design and statistical analyses.*

The experimental design is a split-plot design with 3 cultivars x 6 Mn treatments x 4 replicates for the physiological and biochemical determinations. Data were tested in their normality (Shapiro-Wilk test) and equal variance by Kolmogorov-Smirnoff test and then data were analyzed by two-way ANOVA. Significant differences among means values were established by using the multiple comparisons test of Tukey's ( $P < 0.05$ ). All analyses were performed with Sigma Stat 2.0 software (SPSS, Chicago), where differences between the values were considered significant at  $P \leq 0.05$ .

### **3.3 Results**

*Shoots and roots dry weight*

The lowest values in dry weight of shoots under increasing Mn treatments, were exhibited in Bluegold, where a consistent decrease (~50%) was observed until 50  $\mu\text{M}$  Mn, where values were constant up the highest dose of Mn (1000  $\mu\text{M}$ ) in comparison to control ( $P < 0.005$ ) (Fig. 1A). Legacy and Brigitta manifested also a parallel tendency to decrease under Mn treatments. In roots, no significant differences in dry weight of all cultivars subjected to Mn treatments were exhibited (Fig.1B).

*Mn contents in leaves and roots.* The Mn content in leaves and roots showed a statistically significant interaction between cultivars and Mn treatments ( $P < 0.001$ ). In leaves, Legacy and Brigitta showed a gradual increment, mainly with the highest Mn treatment (68 and 72%, respectively) compared with control. Instead, Bluegold had significant increment in leaves Mn content only with the highest Mn treatment (1000  $\mu\text{M}$ ) with 52% respect to control (Fig. 2). Brigitta had the major Mn content with the highest Mn dose compared with other cultivars. In roots, the most noticeable increment in Mn contents was found in Legacy, where a significant increase from 50 to 1000  $\mu\text{M}$  Mn was evidenced, and where the highest root Mn acquisition of this cultivar was obtained at 500 and 1000  $\mu\text{M}$  Mn (83 and 89%, respectively, compared to control). Brigitta had significant increase in Mn contents with the highest Mn treatment compared

---

with the control (57%), while Bluegold showed significant differences from 250 to 1000  $\mu\text{M}$  Mn treatment, with a 58% and 74% of increase compared with the control, respectively.

### ***Photosynthetic performance***

The maximal photochemical efficiency of PSII (Fv/Fm) and the effective photochemical efficiency of PSII ( $\Phi\text{PSII}$ ), showed a significant decrease in Brigitta and Bluegold but not in Legacy, specially at the higher Mn doses compared with their controls ( $P > 0.005$ , Table 1). Regardless to  $\Phi\text{PSII}$ , Bluegold and Brigitta have lower values than Legacy at higher Mn doses (500 and 1000  $\mu\text{M}$  Mn). Values of NPQ in Legacy were lower than those of Brigitta and Bluegold, and significantly it decreased (53%) at 500  $\mu\text{M}$  Mn compared with control. Bluegold diminished NPQ values significantly from 10  $\mu\text{M}$  Mn to 250  $\mu\text{M}$  Mn and at the highest Mn doses. Moreover, Brigitta exhibited no significant differences in NPQ between Mn treatments. Indeed, Legacy achieved the lowest values of this parameter concomitant with generally higher  $\Phi\text{PSII}$  values.

Exposure of plants to Mn excess brought to a decrease in the net photosynthesis ( $P_n$ ) in Legacy and Bluegold (Fig. 3). Legacy had a significant reduction from the 50  $\mu\text{M}$  Mn to the highest Mn treatment, with percentages of reduction around 56% (50  $\mu\text{M}$ ), 41% (250  $\mu\text{M}$ ), 48% (500  $\mu\text{M}$ ) and 37% (1000  $\mu\text{M}$ ), when they were compared with the controls. Bluegold also exhibited a decrease of  $P_n$  which ended with a notorious fall at the highest Mn treatment (~80% decrease) ( $P \leq 0.005$ ) respect to control. Significant reductions in stomatal conductance ( $g_s$ ) values were found in Bluegold and Brigitta particularly with the highest Mn dose. Differently, Legacy decreased only in the intermediate Mn treatments (250 to 500  $\mu\text{M}$  Mn) without change respect to the control at the higher Mn dose.

### ***Lipid peroxidation and radical scavenging activity***

Analysis of leaves and roots lipid peroxidation (LP), which indicates a damage of cell membranes were performed. In leaves, Brigitta and Bluegold showed an increasing trend of LP by incrementing Mn doses, mainly at the highest Mn dose compared with the control, while Legacy did not show significant changes (Fig. 4). In roots, both Legacy and Brigitta did not have a significant increment of MDA values in all Mn treatments, nevertheless Bluegold showed an increase of MDA values with higher Mn doses.

---

In addition, antioxidant activity, measured as radical scavenging activity (RSA) in leaves of Legacy and Bluegold exhibited an increase with increasing Mn treatments. However, measurements on Bluegold had significant high values from 50 to 1000  $\mu\text{M}$  Mn compared with the control (35 to 46%). Instead, Legacy only had a significant increment at the highest Mn dose (1000  $\mu\text{M}$ ) respect to control (28%, Fig. 5). It is noteworthy that in roots, cultivars had lower values of RSA than leaves.

Regarding to the antioxidant capacity, the SOD activity in leaves of Legacy remained stable whereas Brigitta had statistically significant decreases in the SOD activity from 50  $\mu\text{M}$  to the latest Mn supplies tested (Fig. 6). In roots, only Bluegold showed a significant gradual reduction in SOD activity, mainly at the 1000  $\mu\text{M}$  Mn compared with the control.

### ***Root exudation of carboxylates***

Roots showed a significant increase of oxalate and citrate exudates under Mn supply mostly in Legacy and Brigitta (Fig. 7). It should be noted that, oxalate was the highest organic acid anion exudate, with values around 0.5 to 2  $\mu\text{mol g}^{-1} \text{ h}^{-1}$ . Besides, Legacy exhibited a significant increment of this acid anion from 500  $\mu\text{M}$  (85% up compared with the control). Bluegold showed no significant increment in oxalate exudation. In contrast, malate exudation in Bluegold and Brigitta generally exhibited a significant decrease at incrementing Mn-treatments.

## **3.4 Discussion**

Mn excess can be an important factor of toxicity in acid soils, affecting the physiological and biochemical features of crop plants (Marschner, 1995), generating different resistance responses even in the same species, genotypes or cultivars (Foy et al. 1995; Fecht-Christoffers et al. 2003). In our research, significant DW reductions were highlighted only in Bluegold shoots, even at the lowest Mn treatment (10  $\mu\text{M}$ ; Fig. 1). Reductions of dry weight (DW) due to a Mn excess have been reported in rice (Horiguchi, 1988), ryegrass (Mora et al., 2009), lucerne (Gherardi and Rengel, 2003), pummel (Li et al., 2010), *Cucumis* species (Shi et al., 2005; 2006), maize (Stoyanova et al., 2009) and barley (Alam et al., 2006). Detrimental effects of Mn excess in ryegrass and clover, where DW reductions were higher for shoots than roots were mainly associated with deficient or low P supply (Rosas et al., 2007; Rosas et al., 2011). A major injury in shoots than roots were associated to a Mn toxicity in Mn-sensitive maize cultivar, differing

---

with other metals behavior (i.e. copper and aluminum), which reduces significantly DW at root level (Doncheva et al., 2009). Additionally, no significant changes were found in DW roots in all blueberry cultivars (Fig. 1), which could support the fact that effects of Mn excess are mainly a reduction in shoots more than roots in this woody plant. With respect to Mn contents, shoots and roots increased significantly and gradually with Mn supplies in all cultivars (Fig. 2). Similar results were obtained by Mora et al. (2009), where a large translocation from roots to shoots was shown by Mn-sensitive ryegrass cultivars, whereas the tolerant ones accumulate more Mn content in roots than shoots. These results suggest an internal mechanism of control on Mn content and lower Mn translocation to shoots in tolerant ryegrass cultivar than in Mn-sensitive cultivar, which had a great translocation to shoot (Mora et al. 2009). Similar tendency was showed in *Citrus grandis* plants subjected to Mn supply where the majority of Mn was retained in the roots with a subsequent low Mn content in aerial tissues implying that a Mn-excess might contribute to a tolerance for Mn toxicity (Li et al., 2010). Recently, Yao et al. (2012a) prove that Mn in woody plants, was readily transferred to above-ground plant parts in chestnut rose, plant considered sensitivity to Mn excess, compared to grape plants (Mn tolerant species), which Mn is stored in the root system, which confer a high Mn tolerance. Additionally, these authors suggested that a restriction of xylem loading can be considered as a limiting process for root-to shoot translocation. Considering the above results in other plants, our results showed that Legacy presented a major Mn accumulation tendency in roots, suggesting a Mn tolerance in this cultivar.

Our findings about maximal photochemical efficiency of PSII (Fv/Fm), the effective photochemical efficiency of PSII ( $\Phi$ PSII) in leaves of blueberry cultivars subjected to Mn excess exhibited a significant decrease in Brigitta and Bluegold with higher Mn dose (Table 1). Researches in rice with cadmium (Cd), suggest photoinhibition by loss the PSII efficiency, due to a decrease of Fv/Fm (Pagliano et al., 2006). Excess Mn also decreased significantly  $\Phi$ PSII in cucumber (*Cucumis sativus* L.) seedlings, suggesting that lead to a decline of carbon metabolism capacity and/or a low consumption of ATP and NADPH in the Calvin cycle (Feng et al., 2009; Kitao et al., 1997). NPQ values of blueberry cultivars showed a significant decrease in Bluegold from the lower Mn doses treatment, although there was not change in higher Mn doses, remaining almost constant along the time in the others cultivars (Table 1). Doncheva et al. (2009) found a decrease in Fv/Fm and an increase in NPQ values in the Mn-sensitive cultivar of maize, suggesting a photoinhibition by Mn excess. Although high NPQ values reflect heat dissipation in

---

PSII (Maxwell and Johnson, 2000), sometimes it is not enough to protect efficiently the photosynthetic organelles, as thylakoids and chloroplast (Doncheva et al., 2009). In this sense, net photosynthesis ( $P_n$ ) and stomatal conductance ( $G_s$ ) of Bluegold were negatively affected by excess Mn, mainly with the highest Mn treatment. Legacy had also a significant reduction in  $P_n$ , but without changes in  $G_s$  values. On contrary,  $P_n$  in Brigitta remained stable, but with a significant decline of  $G_s$  (Fig. 3). Stoyanova et al. (2009) observed a negative relation between Mn leaf concentration and net photosynthesis and stomatal conductance in Mn-sensitive maize cultivars. However, these authors found that in Mn-tolerant maize cultivars, the Mn leaf concentrations increased along with transpiration rate, stomatal conductance and photosynthesis. These results suggest that Mn leaf accumulation could be performed by the transpiration stream; leading to that Mn can be transported by xylem from roots. Early experiments in tobacco young leaves grown in nutrient solution with Mn excess showed a decrease in net photosynthesis (Nable et al., 1988), although this decline is not a consequence of decreased leaf  $\text{CO}_2$  conductance in leaves. These authors concluded that photosynthesis inhibition is an early indicator of Mn accumulation excess in tobacco leaves (Nable et al., 1988).

It is known that ROS production can modify metabolic pathways and increase protein denaturation, membranes, lipids injury and DNA mutation, even destroy cellular structures (Hegedüs et al., 2001; Najeeb et al., 2009; Desikan et al., 2005). In our results, Legacy did not show changes of lipid peroxidation. In contrast, higher increment of lipid peroxidation had both Brigitta and Bluegold, mainly in leaves with increasing Mn treatments (Fig. 4). It has been demonstrated that a Mn excess can increase lipid peroxidation in cucumber leaves (Shi et al., 2005; Shi and Zu, 2008) and wetland plants (Najeeb et al., 2009), indicating that Mn excess can cause oxidative stress. In addition, Mora et al., (2009) showed that in Mn-tolerant ryegrass cultivars subjected to Mn toxicity, lipid peroxidation was lower than in Mn-sensitive cultivars. On the other hand, these authors also found that Mn-sensitive cultivars incremented shoot Mn concentration concomitant with an augment in lipid peroxidation.

To prevent and reduce damage induced by ROS, plants have developed antioxidant defense mechanisms, which include enzymatic and non-enzymatic systems (Foyer and Noctor, 2003; Apel and Hirt, 2004). Radical scavenging activity (RSA) in blueberry cultivars showed higher differences in leaves than roots with Mn excess. Bluegold increased significantly this activity

---

starting from 50  $\mu\text{M}$  Mn to the highest Mn-dose in leaves (Fig. 5) and it was proportional to its LP values, reaching maximums at the highest Mn treatment. Interestingly, Legacy had an increase of RSA at the highest Mn treatment, even though lipid peroxidation remained unchanged (Fig. 4). Results with the same blueberry cultivars, but subjected to metal stress (aluminum, Al toxicity) showed that Bluegold is the Al-sensitive cultivar, with the greatest values of lipid peroxidation and lower RSA values (Reyes-Diaz et al., 2010). The authors suggest that Al-tolerant cultivar Legacy can activate defense mechanisms through RSA which contribute to decrease lipid peroxidation values. Likewise, others studies have shown that an enhanced level of antioxidative components can be correlated with increased stress tolerance (Shi et al., 2005). In the same way, it has been suggested that is an increase in superoxide dismutase (SOD) enzyme activity under Mn toxicity, to prevent and mitigate the oxidative damage in plant cells (Srivastava and Dubey, 2011). In our studies, Bluegold had a tendency to decline its SOD activity with the increasing Mn treatment compared with control of leaves and roots. Differently occurs in Legacy, where it was unchanged (Fig. 6). Researches carried out with blueberries cultivars subjected to Al toxicity at short time (hours), showed that Brigitta was considered as an Al-resistant cultivar, which reflected an amelioration of SOD activity under these metal stress. This indicates a bigger capacity for controlling oxidative stress, compared with Bluegold (Inostroza-Blancheteau et al., 2011). Others studies also confirmed detoxification of Al in barley plants, where SOD activity is involved (Simonovicova et al. 2004).

Organic acid anion exudations in plants are recognized for their chelating capacities of heavy metals decreasing the bioavailability of these metals in soils (Najeeb et al. 2009). Exudation of organic acids has different functions, as the uptake of some nutrients as P (Hoffland, 1992; Johnson et al. 1996) and metal detoxification (Jones, 1998). Organic acid anion exudates associated to Mn detoxification are oxalate, citrate, malate and succinate (Horst et al. 1999). Interesting results were observed in roots exudates of organic acids in our experiments (Fig. 7). Of the four organic acid anions analyzed, oxalate presented a higher amount of exudates, compared with others. Similar results were found in ryegrass and white clover grown under Mn excess and acidic conditions, where oxalate and citrate accounted for the most of the carboxylates exudates (Rosas et al., 2007). In our research, Legacy and Brigitta have similar increasing exudation of oxalate, with incrementing Mn treatments. Contrary, in Bluegold there were no

---

changes in this organic acid exudation. Interestingly, this blueberry cultivar showed the highest values of malate in the control, decreasing drastically with Mn treatments.

Mora et al., (2009) found that Mn-tolerant ryegrass had the highest oxalate exudation rate at increasing excess supply of Mn. Those responses suggest that oxalate can act on internal Mn-tolerance mechanisms. Citrate exudation was found in Mn-tolerant ryegrass, associated to the high affinity of citrate for  $Mn^{2+}$  in the rizosphere and high citrate exudation. In our study, Legacy and Brigitta showed an increase in citrate level exudations, compared with Bluegold. Likewise, in researches with wetland plants, Najeeb et al. (2009) showed that citric acid and other organic acids can provided both protons as electrons for reduction of metals in the rizosphere, acting as chelators. Others heavy metals toxicity as lead (Pb) and Cu are associated to Mn tolerance by a release of oxalate in *Arabidopsis* plants and to citrate efflux in rice cultivars (Ryan et al. 2001).

In summary, our results demonstrated that Mn excess affects differently the physiological and biochemical features of blueberry cultivars. Legacy was the most Mn-tolerant cultivar compared to Brigitta and Bluegold, being the last cultivar the most Mn-sensitive. It was due to Legacy cultivar accumulated a high Mn content in roots, maintaining its growth and photosynthesis. Besides, this cultivar showed a higher antioxidant capacity without changes in lipid peroxidation at increasing Mn treatments. Oxalate and citrate were the most important organic acid anions exudated, incrementing its concentration gradually with the enhancement of Mn doses in Legacy.

### **Acknowledgements**

We thank to Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) N°1080372 for the support of this work and Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) Doctoral Fellowship of the Government of Chile. Beca Apoyo Tesis Doctoral, CONICYT, Chile.

---

### 3.5 References

- Alam S, Kodama R, Akiha F, Kamei S, Kawai S.** 2006. Alleviation of manganese phytotoxicity in barley with calcium. *Journal of Plant Nutrition* 29: 59-74.
- Apel K, Hirt H.** 2004. Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annual Review Plant Biology* 55: 373-379.
- Bradford MM.** 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.
- Chinnici F, Bendini A, Gaiani A, Riponi C.** 2004. Radical scavenging activities of peels and pulps from cv. Golden delicious apples as related to their phenolic composition. *Journal of Agricultural and Food Chemistry* 52:4684-4689.
- Demirevska-Kepova K, Simova-Stoilova L, Stoyanova Z, Holzer R, Feller U.** 2004. Biochemical changes in barley plants after excessive supply of copper and manganese. *Environmental Experimental Botanical* 52: 253-266.
- Desikan R, Hancock JT, Bright J, Harrison J, Weir I, Hooley R, Neill SJ.** 2005. A role for ETR1 in hydrogen peroxide signaling in stomatal guard cells. *Plant Physiology* 137: 831–834.
- Doncheva S, Poschenrieder C, Stoyanova Z, Georgieva K, Velichkova M, Barceló J.** 2009. Silicon amelioration of manganese toxicity in Mn-sensitive and Mn-tolerant maize varieties. *Environmental and Experimental Botany* 65: 189–197.
- Du Z, Bramlage W.** 1992. Modified thiobarbituric acid assay for measuring lipid oxidation in sugar-rich plant tissue extracts. *Journal of Agricultural and Food Chemistry* 40: 1556-1570.
- Fecht-Christoffers MM, Maier P, Horst W.** 2003. Apoplastic peroxidase and ascorbate are involved in manganese toxicity and tolerance of *Vigna unguiculata*. *Physiologia Plantarum* 117: 237-244.

- 
- Feng JP, Shi QH, Wang XF.** 2009. Effects of exogenous silicon on photosynthetic capacity and antioxidant enzyme activities in chloroplast of cucumber seedlings under excess manganese. *Agricultural Science in China* 8: 40–50.
- Foy CD, Weil RR, Coradetti CA.** 1995. Differential manganese tolerances of cotton genotypes in nutrient solution. *Journal of Plant Nutrition* 18: 685-706.
- Foyer CD, Noctor G.** 2003. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiologia Plantarum* 119: 355-364.
- Gangwar S, Singh VP, Prasad SM, Maurya JN.** 2010. Modulation of manganese toxicity in *Pisum sativum* L. seedlings by kinetin. *Scientia Horticulturae* 126: 467–474
- Gherardi M, Rengel, Z.** 2003. Genotypes of lucerne (*Medicago sativa* L.) show differential tolerance to manganese deficiency and toxicity when grown in bauxite residue sand. *Plant Soil* 249: 287-296.
- Giannopolitis CN, Ries SK.** 1977. Superoxide dismutases occurrence in higher plants. *Plant Physiology* 59: 309-314.
- Hauck M, Paul A, Mulack C, Fritz E, Runge M.** 2002. Effects of manganese on the viability of vegetative diaspore of the epiphytic lichen *Hypogymnia physodes*. *Environmental Experimental Botanical* 47: 127-142.
- Hauck M, Paul A, Gross S, Raubuch M.** 2003. Manganese toxicity in epiphytic lichens: chlorophyll degradation and interaction with iron and phosphorus. *Environmental and Experimental Botany* 49: 181-191.
- Hauck M, Paul A, Spribille T.** 2006. Uptake and toxicity of manganese in epiphytic cyanolichens. *Environmental and Experimental Botany* 56: 216–224.
- Hegedüs A, Erdei S, Horváth G.** 2001. Comparative studies of H<sub>2</sub>O<sub>2</sub> detoxifying enzymes in green and greening barley seedlings under cadmium stress. *Plant Science* 160:1085–1093.
- Hoagland DR, Arnon DI.**1959. The water culture method for growing plants without soil. *California Agricultural Experimental Station* 347:1–32.


- 
- Hodges DM, DeLong JM, Forney CF, Prange RK.** 1999. Improving the thiobarbituric acid-reactive substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* 207: 604–611.
- Hoffland E.** 1992. Quantitative evaluation of the role of organic acid exudation in the mobilization of rock phosphate by rape. *Plant Soil* 140: 279-289.
- Horiguchi T.** 1988. Mechanism of manganese toxicity and tolerance of plants. VII. Effect of light intensity on manganese-induced chlorosis. *Journal of Plant Nutrition* 11: 235–245.
- Horst W, Fecht M, Naumann A, Wissemeier A, Maier P.** 1999. Physiology of manganese toxicity and tolerance in *Vigna unguiculata* (L.) Walp. *Journal of Plant Nutrition and Soil Science* 162: 263-274.
- Horst WJ.** 1988. The physiology of manganese toxicity. *In* Manganese in Soils and Plants. Eds. R D Graham, R J Hannam and N C Uren. pp 175-188. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Inostroza-Blancheteau C, Reyes-Díaz M, Aquea F, Nunes-Nesi A, Alberdi M, Arce-Johnson P.** 2011. Biochemical and molecular changes in response to aluminium-stress in highbush blueberry (*Vaccinium corymbosum* L.). *Plant Physiology and Biochemistry* 49: 1005-1012.
- Johnson JF, Allan D, Vance CP, Weiblen C.** 1996. Root carbon dioxide fixation by phosphorus-deficient *Lupinus albus*. Contribution to organic acid exudation by proteoid roots. *Plant Physiology* 112: 19-30.
- Jones DL.** 1998. Organic acids in the rhizosphere – a critical review. *Plant and Soil* 205: 25-44.
- Kitao M, Lei TT, Koike T.** 1997. Effects of manganese toxicity on photosynthesis of white birch (*Betula platyphylla* var. *japonica*) seedlings. *Physiologia Plantarum* 101: 249-256.
- Lei Y, Korpelainen H, Li C.** 2007. Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. *Chemosphere* 68: 686-694.

- 
- Li Q, Chen LS, Jiang HX, Tang N, Yang LT, Lin ZH, Li Y, Yang GH.** 2010. Effects of manganese-excess on CO<sub>2</sub> assimilation ribulose-1,5-bisphosphate carboxylase/oxygenase, carbohydrates and photosynthetic electron transport of leaves, and antioxidant systems of leaves and roots in *Citrus grandis* seedlings. *BMC Plant Biology* 10: 42.
- Macfie S, Taylor G.** 1992. The effects of excess manganese on photosynthetic rate and concentration of chlorophyll in *Triticum aestivum* grown in solution culture. *Physiologia Plantarum* 85:467-475.
- Marschner H.** 1995. Mineral nutrition of higher plants. San Diego: Academic Press.
- Maxwell K, Johnson G.** 2000. Chlorophyll fluorescence. A practical guide. *Journal of Experimental Botany* 51: 659-668.
- Meier S, Alvear M, Borie F, Aguilera P, Ginocchio R, Cornejo P.** 2012. Influence of copper on root exudate patterns in some metallophytes and agricultural plants. *Ecotoxicology and Environmental Safety* 75: 8–15.
- Millaleo R, Reyes-Diaz M, Ivanov AG, Mora ML, Alberdi M.** 2010. Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms. *Journal of soil science and plant nutrition* 10: 470-481.
- Mora ML, Rosas A, Ribera A, Rengel R.** 2009. Differential tolerance to Mn toxicity in perennial ryegrass genotypes: involvement of antioxidative enzymes and root exudation of carboxylates. *Plant and Soil* 320: 79–89.
- Nable RO, Houtz RL, Cheniae GM.** 1988. Early inhibition of photosynthesis during development of Mn toxicity in tobacco. *Plant Physiology* 86: 1136–1142.
- Najeeb U, Xu L, Ali S, Jilani G, Gongga HJ, Shenc WQ, Zhou WJ.** 2009. Citric acid enhances the phytoextraction of manganese and plant growth by alleviating the ultrastructural damages in *Juncus effusus* L. *Journal of Hazardous Materials* 170:1156–1163.


- 
- Pagliano C, Raviolo M, Vecchia FD, Gabbrielli R, Gonnelli C, Rascio N, Barbato R, La Rocca N.** 2006. Evidence for PSII donor-side damage and photoinhibition induced by cadmium treatment on rice (*Oryza sativa* L.). *Journal of Photochemistry and Photobiology B: Biology* 84: 70–78.
- Rengel Z.** 2000. Manganese uptake and transport in plants. *Metal ions in biological systems* 37: 57-87.
- Reyes-Díaz M, Alberdi M, Mora ML.** 2009. Short-term aluminum stress differentially affects the photochemical efficiency of photosystem II in highbush blueberry genotypes. *Journal of American Society Horticultural Science* 134:14–21.
- Reyes-Díaz M, Inostroza-Blancheteau C, Millaleo R, Cruces E, Wulff-Zottele C, Alberdi M, Mora ML.** 2010. Long-term aluminum exposure effects on physiological and biochemical features of highbush blueberry cultivars. *Journal of American Society Horticultural Science* 3: 212-222.
- Rojas-Lillo Y, Alberdi M, Acevedo P, Inostroza-Blancheteau C, Rengel Z, Mora ML, Reyes-Díaz M.** 2014. Manganese toxicity and UV-B radiation differentially influence the physiology and biochemistry of highbush blueberry (*Vaccinium corymbosum*) cultivars. *Functional Plant Biology* 41: 156-167.
- Rosas A, Rengel Z, Mora ML.** 2007. Manganese supply and pH influence growth, carboxylate exudation and peroxidase activity of ryegrass and white clover. *Journal of Plant Nutrition* 30: 253-270.
- Rosas A, Rengel Z, Ribera A, Mora ML.** 2011. Phosphorus nutrition alleviates manganese toxicity in *Lolium perenne* and *Trifolium repens*. *Journal of Plant Nutrition and Soil Science* 174: 210–219.
- Ryan PR, Delhaize E, Jones DL.** 2001. Function and mechanism of organic anion exudation from plant roots. *Annual Review of Plant Physiology and Plant Molecular Biology* 52: 527–60.

- 
- Sadzawka A, Carrasco M, Grez R, Mora M.** 2004. Métodos de análisis de tejidos vegetales. Comisión de normalización y acreditación. Sociedad Chilena de la Ciencia del Suelo (SCCS).
- Shi Q, Bao Z, Zhu Z, He Y, Qian Q, Yu J.** 2005. Silicon-mediated alleviation of Mn toxicity in *Cucumis sativus* in relation to activities of superoxide dismutase and ascorbate peroxidase. *Phytochemistry* 66:1551-1559.
- Shi Q, Zhu Z, Xu M, Qian Q, Yu J.** 2006. Effect of excess manganese on the antioxidant system in *Cucumis sativus* L. under two light intensities. *Environmental and Experimental Botany* 58: 197-205.
- Shi Q, Zhu Z.** 2008. Effects of exogenous salicylic acid on manganese toxicity, element contents and antioxidative system in cucumber. *Environmental and Experimental Botany* 63: 317–326.
- Simonovicova M, Tamás L, Huttová J, Mistrik I.** 2004. Effect of aluminum on oxidative stress related enzymes activities in barley roots. *Biologia Plantarum* 48: 261-266.
- Srivastava S, Dubey RS.** 2011. Manganese-excess induces oxidative stress, lowers the pool of antioxidants and elevates activities of key antioxidative enzymes in rice seedlings. *Plant Growth Regulation* 64:1–16.
- Stoyanova Z, Poschenrieder C, Tzvetkova N, Doncheva S.** 2009. Characterization of the tolerance to excess manganese in four maize varieties. *Soil Science and Plant Nutrition* 55:747–753.
- Wang S, Mulligan CN.** 2009. Effect of natural organic matter on arsenic mobilization from mine tailings. *Journal of Hazardous Materials* 168: 721–726.
- Yao YA, Mou D, Xu G, Lutts S, Achal V, Ma J.** 2012a. Contrasting performance and different tolerance of chestnut rose and grape to excess manganese. *Journal of Plant Growth Regulation* 31: 416-426.

Tables and Figures




**Figure 1.** Effect of Mn treatments on dry weight of shoot (leaves + stems) (**A**) and roots dry weight (**B**) in three blueberry cultivars grown in nutrient solution for 25 days. Results are means  $\pm$  standard error (SE), with n=3. Different lower-case letters indicate significant differences between Mn treatments for the same cultivar ( $P \leq 0.05$ , Tukey's Test).




**Figure 2.** Mn contents in leaves and roots of blueberry cultivars with Mn excess-doses at the end of experiment (25 days). Different lower-case letters indicate significant differences between the Mn treatments for the same cultivar, while different upper-case letters indicate significant differences between the cultivars at a same Mn treatment ( $P \leq 0.05$ ).

**Table 1.** Maximal photochemical efficiency of PSII (Fv/Fm), effective photochemical efficiency of PSII ( $\Phi$ PSII) and non-photochemical quenching (NPQ) of three blueberry cultivars subjected to increasing Mn levels (25 days). Different lower-case letters indicate significant differences between Mn treatments at the same cultivar. Different upper-case letters indicate significant differences between cultivars at the same Mn treatment ( $P \leq 0.05$ ).


<b>Cultivar</b>	<b>Mn treatment [<math>\mu</math>M]</b>	<b>Fv/Fm</b>	<b><math>\Phi</math>PSII</b>	<b>NPQ</b>
Legacy	2	0.78 $\pm$ 0.005 <b>aA</b>	0.115 $\pm$ 0.006 <b>aA</b>	1.18 $\pm$ 0.2 <b>aB</b>
	10	0.77 $\pm$ 0.008 <b>aA</b>	0.103 $\pm$ 0.015 <b>aAB</b>	1.07 $\pm$ 0.2 <b>aB</b>
	50	0.78 $\pm$ 0.004 <b>aA</b>	0.102 $\pm$ 0.006 <b>aA</b>	1.11 $\pm$ 0.2 <b>aB</b>
	250	0.78 $\pm$ 0.002 <b>aAB</b>	0.092 $\pm$ 0.006 <b>aA</b>	1.04 $\pm$ 0.1 <b>aB</b>
	500	0.77 $\pm$ 0.009 <b>aA</b>	0.093 $\pm$ 0.013 <b>aA</b>	0.55 $\pm$ 0.1 <b>bB</b>
	1000	0.76 $\pm$ 0.008 <b>aA</b>	0.089 $\pm$ 0.008 <b>aA</b>	1.53 $\pm$ 0.3 <b>aB</b>
Brigitta	2	0.81 $\pm$ 0.004 <b>aA</b>	0.104 $\pm$ 0.008 <b>aA</b>	2.35 $\pm$ 0.1 <b>aA</b>
	10	0.78 $\pm$ 0.012 <b>abA</b>	0.084 $\pm$ 0.007 <b>abB</b>	2.41 $\pm$ 0.1 <b>aA</b>
	50	0.77 $\pm$ 0.012 <b>bA</b>	0.085 $\pm$ 0.001 <b>abA</b>	2.31 $\pm$ 0.2 <b>aA</b>
	250	0.80 $\pm$ 0.004 <b>aA</b>	0.076 $\pm$ 0.004 <b>abAB</b>	2.62 $\pm$ 0.2 <b>aA</b>
	500	0.78 $\pm$ 0.010 <b>abA</b>	0.064 $\pm$ 0.005 <b>bB</b>	2.86 $\pm$ 0.1 <b>aA</b>
	1000	0.74 $\pm$ 0.005 <b>bA</b>	0.069 $\pm$ 0.002 <b>bAB</b>	2.77 $\pm$ 0.24 <b>aA</b>
Bluegold	2	0.81 $\pm$ 0.016 <b>aA</b>	0.116 $\pm$ 0.006 <b>aA</b>	2.46 $\pm$ 0.2 <b>aA</b>
	10	0.76 $\pm$ 0.013 <b>bA</b>	0.117 $\pm$ 0.001 <b>aA</b>	1.57 $\pm$ 0.2 <b>bB</b>
	50	0.77 $\pm$ 0.005 <b>bA</b>	0.076 $\pm$ 0.010 <b>abA</b>	1.62 $\pm$ 0.2 <b>bAB</b>
	250	0.77 $\pm$ 0.006 <b>abB</b>	0.063 $\pm$ 0.003 <b>bB</b>	1.64 $\pm$ 0.3 <b>bB</b>
	500	0.78 $\pm$ 0.002 <b>abA</b>	0.059 $\pm$ 0.003 <b>bB</b>	2.38 $\pm$ 0.04 <b>aA</b>
	1000	0.76 $\pm$ 0.004 <b>bA</b>	0.056 $\pm$ 0.005 <b>bB</b>	1.87 $\pm$ 0.3 <b>abB</b>




**Figure 3.** Net photosynthesis ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) and stomatal conductance ( $\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$ ) of three highbush blueberry cultivars subjected to increase Mn treatments at the end of experiment (25 days). Values are means of four measurements. Different lower-case letters indicate significant differences between Mn treatments at the same cultivar. Different upper-case letters indicate significant differences between cultivars at the same Mn treatment ( $P \leq 0.05$ ).




**Figure 4.** Lipid peroxidation of leaves and roots of three highbush blueberry cultivars subjected to different Mn treatments for 25 days. It was measured as malondialdehyde (MDA). Values are an average of six replicates each  $\pm$  SE. Different lower-case letters indicate significant differences between Mn treatments at the same cultivar and the same plant organ ( $P \leq 0.05$ ).



**Figure 5.** Radical-scavenging capacity (RSA) of three highbush blueberry cultivars subjected to different Mn treatments at leaves and roots at 25 days, measured as Trolox equivalents (TE). Different lower-case letters indicate statistically significant differences between Mn treatments for the same cultivar ( $P < 0.05$ ).



**Figure 6.** Effect of Mn-excess on the activity of SOD enzyme in leaves and roots of three blueberry cultivars with excess Mn treatments at 25 days of treatment. Values are means of four replicates each  $\pm$  SE. Different lower-case letters indicate statistically significant differences between Mn treatments for the same cultivar ( $P < 0.05$ ).



**Figure 7.** Organic acid anions exuded by roots of blueberry cultivars grown with increasing Mn levels for 25 d. Different lower case letters indicate statistically significant differences between Mn treatments for the same cultivar and different upper case letters show differences between cultivars for the same Mn treatment ( $P < 0.05$ ).

---

Chapter 4:

Excess manganese (Mn) differentially inhibits  
Photosystem I (PSI) versus Photosystem II (PSII) in  
*Arabidopsis thaliana*.

Article published in Journal of Experimental Botany (2013) 64: 343-354.

DOI: 10.1093/jxb/err313

---

## **Excess manganese (Mn) differentially inhibits Photosystem I (PSI) versus Photosystem II (PSII) in *Arabidopsis thaliana***

R. Millaleo<sup>1</sup>, M. Reyes-Díaz<sup>2,3</sup>, M. Alberdi<sup>2,3\*</sup>, A.G. Ivanov<sup>4</sup>, M. Krol<sup>4</sup>, N.P.A. Hüner<sup>4</sup>

<sup>1</sup>Programa de Doctorado en Ciencias de Recursos Naturales, Universidad de La Frontera, Temuco, Chile.

<sup>2</sup>Departamento de Ciencias Químicas y Recursos Naturales; Facultad de Ingeniería, Ciencias y Administración, Universidad de La Frontera, Temuco, Chile.

<sup>3</sup>Center of Plant, Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN-UFRO), Universidad de La Frontera, Temuco, Chile.

<sup>4</sup>Department of Biology and The Biotron Centre for Experimental Climate Change Research, Western University, London, Ontario, Canada, N6A 5B7.

\*Corresponding author: Miren Alberdi Lag

Departamento de Ciencias Químicas y Recursos Naturales,  
Facultad de Ingeniería, Ciencias y Administración,  
Universidad de La Frontera

Casilla 54-D

Temuco (Chile)

Telephone: (56-45) 2325470

Fax: (56-45) 2325050

e-mail: [miren.alberdi@ufrontera.cl](mailto:miren.alberdi@ufrontera.cl)

---

## Abstract

The effects of exposure to increasing Mn concentrations (50 to 1500  $\mu\text{M}$ ) from the start of the experiment on the functional performance of photosystem II (PSII) and photosystem I (PSI) and photosynthetic apparatus composition of *Arabidopsis thaliana* were compared. In agreement with earlier studies excess Mn caused minimal changes in the PSII photochemical efficiency measured as  $F_v/F_m$ , although the characteristic peak temperature of the  $S_{2/3}Q_B^-$  charge recombinations was shifted to lower temperatures at the highest Mn concentration. SDS-PAGE and immunoblot analyses also did not exhibit any significant change in the relative abundance of PSII-associated polypeptides: PSII reaction center protein D1, Lhcb1 (major light harvesting protein of LHCII complex) and PsbO (OEC33 - 33kDa protein of the oxygen-evolving complex). In addition, the abundance of Rubisco also did not change with Mn treatments. However, plants grown under excess Mn exhibited increased susceptibility to PSII photoinhibition. In contrast, *in vivo* measurements of the redox transients of PSI reaction centre (P700) showed a considerable gradual decrease in the extent of P700 photooxidation ( $P700^+$ ) under increased Mn concentrations compared to control. This was accompanied by a slower rate of  $P700^+$  re-reduction indicating a down-regulation of the PSI-dependent cyclic electron flow. The abundance of PSI reaction center polypeptides (PsaA and PsaB) in plants under the highest Mn concentration was also significantly lower compared to the control. The results demonstrate for the first time that PSI is the major target of Mn toxicity within the photosynthetic apparatus of *Arabidopsis* plants. The possible involvement mechanisms of Mn toxicity targeting specifically PSI are discussed.

**Keywords:** chlorophyll fluorescence, Mn toxicity, photosystem I, PSI- PSII-associated proteins, redox state of P700.

---

## 4.1 Introduction

Manganese (Mn) is one of the most abundant metals in the Earth's crust and although it is an important essential micronutrient for all photosynthetic organisms can be also toxic when it is present in excess (Mukhopadhyay and Sharma, 1991; Marschner, 1995). Mn is considered the second most phytotoxic element, after aluminum (Al), affecting negatively the physiological and biochemical properties of plant species (Foy *et al.*, 1978, Foy, 1984; Millaleo *et al.*, 2010). An excess of this metal occurs in acid soils with low pH (<5.5) and/or under reducing conditions (Marschner, 1995; Schaaf *et al.*, 2002), where  $Mn^{2+}$  is the predominant solution species and available ion to plant cells (Bradl, 2004). Thus, a Mn excess results in a sharp decrease in shoot height, biomass accumulation and total leaf area of a woody species (*Populus cathayana*) (Lei *et al.*, 2007), a reduction of the dry weights (DW) of both shoots and roots in ryegrass cultivars (*Lolium perenne*) (Mora *et al.*, 2009) and *Trifolium repens* (Rosas *et al.*, 2007). Furthermore, excess Mn can result in oxidative stress as indicated by the accumulation of  $H_2O_2$  (Demirevska-Kepova *et al.*, 2004; Lei *et al.*, 2007), high levels of apoplastic  $H_2O_2$ -consuming peroxidases (Fecht-Christoffers *et al.*, 2003) and high level of lipid peroxidation (Mora *et al.*, 2009). Mn-stress induced enhancement of antioxidant enzyme activity in leaves of legumes (González *et al.*, 1998), it was also demonstrated in perennial ryegrass (Mora *et al.*, 2009) and woody species (Lei *et al.*, 2007). More recently, a proteomic and transcriptomic studies have demonstrated that chloroplastic proteins important for  $CO_2$  fixation and photosynthesis were of lower abundance upon Mn stress of cowpea (Führs *et al.*, 2008).

Mn has an important role in both the structure and functions of the photosynthetic apparatus (Mukhopadhyay and Sharma, 1991). Mn is a constitutive element associated with the oxygen evolving complex (OEC) of photosystem II (PSII), an important multi-protein pigment complex embedded in the thylakoid membranes (Hankamer *et al.*, 1997; Enami *et al.*, 2008). Therefore, the Mn cluster, together with other ions and extrinsic proteins that constitute OEC are required to oxidize water and reduce P680, the reaction center of PSII (Kern and Renger, 2007; Ferreira *et al.*, 2004; Rutherford and Boussac, 2004). In conjunction with photosystem I (PSI) and linear electron transport, these reducing equivalents (electrons) are used primarily in the conversion of  $CO_2$  into carbohydrate (Ferreira *et al.*, 2004). In addition, Mn is indispensable as a cofactor for various enzymes

---

involved in redox reactions such as Mn-SOD, an essential enzyme involved in protection against oxidative stress in plants (Burnell, 1988; Bowler *et al.*, 1994).

A number of studies have suggested that chloroplasts and photosynthesis are the major targets of Mn toxicity. Indeed, increased amounts of Mn were reported for chloroplasts isolated from Mn-stressed common bean (González and Lynch, 1999) and rice leaves (Lidon *et al.*, 2004). Distinctive ultrastructural changes showing swelling of granal and stromal thylakoids have been also observed in the chloroplasts of *Citrus volkameriana* (Papadakis *et al.*, 2007) and maize plants (Doncheva *et al.*, 2009) under Mn excess. It has been demonstrated that high Mn accumulation is associated with inhibition of the net photosynthesis and carboxylation efficiency in various plant species. The decline of photosynthesis is considered as one of the major mechanisms constituting the toxic effects of excess Mn and is proposed as an early indicator for Mn toxicity in tobacco (Nable *et al.*, 1988), rice (Lidon *et al.*, 2004) and wheat (Macfie and Taylor, 1992). Reduced CO<sub>2</sub> assimilation induced by excess Mn was also reported for common bean (González and Lynch, 1997), deciduous broad leaved trees (Kitao *et al.*, 1997a) and seedlings of *Citrus grandis* (Li *et al.*, 2010). Interestingly, the maximum efficiency of PSII photochemistry ( $F_v/F_m$ ) was not substantially affected by Mn accumulation in various plant species over a wide range of leaf Mn concentrations (Nable *et al.*, 1988; Kitao *et al.*, 1997b; Subrahmanyam and Rathore, 2000; Hajiboland and Hasani, 2007; Doncheva *et al.*, 2009).

The reduction in photosynthesis by excess leaf Mn has been generally attributed to modification of ribulose-1,5- bisphosphate carboxylase/oxygenase (Ohki, 1984; Houtz *et al.*, 1988; McDaniel and Toman, 1994; Kitao *et al.*, 1997b). It has been demonstrated that a high level of Mn affects primarily the activity rather than the amount of RUBISCO (Houtz *et al.*, 1988; Chatterjee *et al.*, 1994) and the presence of excess Mn induces enhanced oxygenase activity (Jordan and Ogren, 1983). In addition, the decline of photosynthesis under Mn stress conditions was also ascribed to peroxidative impairment of photosynthetic enzyme activities caused by polyphenol oxidation products (Panda *et al.*, 1987; Vaughn and Duke, 1984).

In spite of these studies the mechanisms of Mn toxicity causing a decrease in CO<sub>2</sub> assimilation are still not well understood. In addition, very limited information concerning the toxic effect(s) of excess Mn on the polypeptide composition of both PSII and especially

---

PSI is available. Therefore, the objective of this study was to evaluate the role of specific Mn induced changes in the structure and function of PSII and PSI, which could help to understand the mechanisms by which Mn-excess may cause a decrease of CO<sub>2</sub> assimilation in *Arabidopsis thaliana*.

## **4.2 Materials and methods**

**4.2.1 Plant material and growth conditions.** Seeds of *Arabidopsis thaliana* (wild type Columbia) were germinated in a substrate mix (82.5% sphagnum peat moss, 12.5% perlite, 5% vermiculite-Pro-Mix, Premier Tech Horticulture Ltd.) in controlled environment growth cabinets (Model GCW15, Environmental Growth Chambers, Chagrin Falls, OH) with a photosynthetic active radiation (PAR) of 250  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , 20/20°C day/night temperatures, 50% relative humidity and an 8-h photoperiod to prevent flowering. Water was supplied every five days. After 15 days, seedlings were transplanted separately in pots with vermiculite and placed in trays. Each tray containing seven pots (one plant per pot), were supplied with Hoagland nutrient solution for two weeks before applying the Mn treatments (see below).

**4.2.2 Manganese treatments.** Manganese treatments included the final concentrations: 18 (control), 50, 500, 1000 and 1500  $\mu\text{M Mn}$ . Doses applied were prepared according to Delhaize *et al.*, (2007). Manganese was applied as  $\text{MnCl}_2 \times 4\text{H}_2\text{O}$ . Control plants exposed to 18  $\mu\text{M Mn}$  was optimal dose for Mn for *Arabidopsis* (Cailliatte *et al.*, 2010). The five Mn treatments were grown in five labeled trays, with 500 mL of Hoagland's solution. The trays were maintained in controlled environment growth chambers under the same conditions described above. The pH was adjusted to 5.3 with diluted HCl daily and nutrient solution was changed every five days. Plants were subjected to these treatments for 21 days before harvest.

**4.2.3 Plant growth measurements.** Prior to beginning the Mn-treatments, three plant samples were dried in a forced-air oven (70°C, 48 hrs) and weighed to determine dry weight (W1) at 0 day. Similarly, at the end of the experiment, plants were harvested and

---

collected for fresh and dry weight measurements (W2). These data were used to determine mean relative growth rate (RGR) according to Fernando *et al.* (2009):

$$RGR = \frac{(\ln W2 - \ln W1)}{(t2 - t1)}$$

as  $\text{g g}^{-1} \text{d}^{-1}$  DW.

**4.2.4 Manganese concentration.** For Mn chemical analysis, samples of shoot and roots were dry-ashed in a muffle furnace at 500°C for 8 h and digested with 2 M HCl. Manganese was extracted as described by Sadzawka *et al.* (2004), and the Mn concentration was determined using a simultaneous multi-element atomic absorption spectrophotometer (Model 969 atomic absorption spectrometer; Unicam, Cambridge, UK).

**4.2.5 Thylakoid preparation, SDS-PAGE and immunoblotting.** Thylakoid membranes for SDS-PAGE were isolated as described earlier (Krol *et al.*, 1999). Leaf material was ground in cold isolation buffer (50 mM Tricine, 0.4 M sorbitol, 10mM NaCl, 5mM MgCl<sub>2</sub>, pH 7.8) in a mortar on ice, filtered through two layers of miracloth (typical pore size 22-25  $\mu\text{m}$ ; Calbiochem), and centrifuged for 15 min (10,000 g). The supernatant was removed and the pellet was resuspended in cold isolation buffer. Total chlorophyll concentration was measured in 90% (v/v) acetone (Arnon, 1949). For immunodetection of Rubisco total leaf proteins were extracted as described in Rosso *et al.* (2009). Protein content was measured using a BCA protein assay kit (Pierce) by following the absorbance at 562 nm using a Beckman DU-640 spectrophotometer (Beckman Coulter). Proteins were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) according to Laemmli (1970), using 15% (w/v) polyacrylamide gel in the presence of 6 M urea in the separating gel. Chloroplast thylakoids were solubilized with SDS (SDS: chlorophyll, 20:1) and 15  $\mu\text{g}$  of chlorophyll was loaded per lane. All samples for separation of total proteins were loaded on an equal protein basis of 20  $\mu\text{g}$  protein per lane (Rosso *et al.* 2009). Immunoblotting was performed by electrophoretically transferring the proteins from SDS-PAGE gel to nitrocellulose membrane (Bio-Rad) according to the method of Towbin *et al.* (1979). Proteins were probed with antibodies (AgriSera AB, Vanas, Sweden) raised against the reaction center polypeptides of PSI: PsaA, PsaB (1:2000), the major light harvesting

---

protein of PSII complex (LHCII): Lhcb1 protein, (1:5000), the PSII oxygen evolving complex (OEC) extrinsic protein: PsbO (33 kDa) (1:2000), the PSII reaction center protein: D1 and ribulose-1.5-bisphosphate carboxylase/oxygenase protein (Rubisco) (1:5000). As secondary antibodies goat anti-rabbit IgG conjugated with horse-radish peroxidase (Sigma-Aldrich) were used. Polypeptides were detected using enhanced chemiluminescence (ECL) detection kit (Amersham Biosciences) and visualized by exposing the membrane to x-ray film. Densitometric scanning and analysis of X-ray films from each replicate immunoblot was performed with a Hewlett Packard ScanJet 4200C desktop scanner and ImageJ 1.41o densitometry software (Wayne Rosband, National Institute of Health, USA, <http://rsb.info.nih.gov/ij>).

**4.2.6 Measurement of the redox state of P<sub>700</sub>.** The redox state of P<sub>700</sub> was determined *in vivo*, in dark adapted (20 min) *Arabidopsis* leaves under growth temperature and ambient O<sub>2</sub> and CO<sub>2</sub> conditions using a PAM-101 modulated fluorometer equipped with a dual-wavelength emitter-detector ED-P700DW unit and PAM-102 units (Klughammer and Schreiber, 1991) as described in detail by Ivanov *et al.* (1998). Far-red light ( $\lambda_{\text{max}}=715$  nm, 10 W m<sup>-2</sup>, Schott filter RG 715) was provided by an FL-101 light source. The redox state of P700 was evaluated as the absorbance change around 820nm ( $\Delta A_{820-860}$ ) in a custom-designed cuvette. Multiple turnover (MT, 50 ms) and single turnover (ST, half peak 14  $\mu$ s) saturating flashes were applied with XMT-103 and XST-103 (Walz) power/control units, respectively. The relative functional pool size of intersystem electrons on a P<sub>700</sub> reaction center basis was calculated as the complementary area between the oxidation curve of P700 after either ST or MT pulse excitation (ST- and MT-areas) and the stationary level of P700 under FR excitation (Asada *et al.*, 1993; Ivanov *et al.*, 1998). Capacity of PSI cyclic electron (e<sup>-</sup>) flow was determined as the half time for the dark decay of the P700 signal (Ivanov *et al.*, 1998).

**4.2.7 Modulated chlorophyll fluorescence measurements.** Modulated imaging fluorometer (IMAGING-PAM, Heinz Walz GmbH, Efeltrich, Germany) was used for capturing the chlorophyll fluorescence images and estimation of the maximal photochemical efficiency of PSII [ $F_v/F_m = (F_m - F_o)/F_m$ ], quantum yield of PSII

---

photochemistry ( $\Phi_{\text{PSII}}$ ), photochemical (qP) and non-photochemical (NPQ) fluorescence quenching parameters using the nomenclature of van Kooten and Snel (1990) as described earlier (Ivanov *et al.*, 2006a). Control and Mn-treated *Arabidopsis* plants were dark adapted (20 min) and all chlorophyll fluorescence measurements were performed *in vivo* at room temperature. Fluorescence images were captured by a CCD camera (IMAG-K, Allied Vision Technologies) featuring 640x480 pixel CCD chip size and CCTV camera lens (Cosmicar/Pentax F1.2, f=12 mm). Light emitting diode ring array (IMAG-L) consisting of 96 blue LEDs (470 nm) provided standard modulated excitation intensity of  $0.5 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$  (modulation frequency 1-8 Hz) for measuring the basal ( $F_0$ ) chlorophyll fluorescence and a saturation pulse of  $2400 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$  PAR for measuring the maximal chlorophyll fluorescence ( $F_m$ ). The maximal PSII photochemical efficiency ( $F_v/F_m$ ) was determined as  $(F_m - F_0)/F_m$ . The effective photochemical efficiency of PSII ( $\Phi_{\text{PSII}}$ ) was calculated from the expression  $(F'_m - F_s)/F'_m$  (Genty *et al.*, 1989), photochemical quenching (qP) was calculated as:  $(F'_m - F_s)/(F'_m - F'_0)$  (Schreiber *et al.*, 1994), electron transport rate (ETR) was calculated as:  $\text{ETR} = \text{PAR} \times 0.5 \times \Phi_{\text{PSII}} \times 0.84$  (Genty *et al.*, 1989), regulated non-photochemical quenching ( $\Phi_{\text{NPQ}}$ ), constitutive photochemical quenching ( $\Phi_{\text{NO}}$ ) was determined according to Kramer *et al.* (2004). All measurements were performed at 0, 5, 10, 15 and 21 days in plants subjected to the different Mn treatments.

At the end of experiments (21 days), leaves of *Arabidopsis* plants growing under the different Mn treatments and  $250 \mu\text{mol m}^{-2} \text{s}^{-1}$  PAR (normal light), were cut and exposed to high light ( $1000 \mu\text{mol s}^{-1} \text{m}^{-2}$ , PAR) at 1, 2 and 4 hours. During each time of exposition,  $F_v/F_m$  was determined.

**4.2.8 Thermoluminescence measurements.** Thermoluminescence (TL) measurements of control and Mn-treated *Arabidopsis* leaves were performed on a personal-computer-based TL data acquisition and analysis system as described earlier (Ivanov *et al.*, 2001; 2006b). A photomultiplier tube (Hamamatsu R943-02, Hamamatsu Photonics K.K., Shizuoka-ken, Japan) equipped with a photomultiplier power supply (Model PS-302, EG&G Electro Optics) and a preamplifier (Model C1556-03) was used as a radiation measuring set. The heating rate was  $0.6^\circ\text{C s}^{-1}$ . For identifying the  $S_{2/3}Q_B^-$  charge recombination peaks, dark

---

adapted leaf discs were subjected to two consecutive saturating microsecond flashes of white light (1.5  $\mu$ s peak width at 50% of maximum) applied by a xenon-discharge flash lamp (XST103, Heinz Walz GmbH, Effeltrich, Germany). Dark adapted leaves (20 minutes at 20°C) were cooled to 2°C prior to exposing to the flashes. The nomenclature of Sane *et al.* (2012) was used for characterisation of the TL glow peaks.

**4.2.9 Experimental design and statistical analysis.** The experimental design is a randomized split design with one species x-five Mn treatments x-seven replicates for the physiological determinations, and one species x five Mn treatments x three replicates x four times (0, 1, 2 and 4 hours) for the photoinhibition measurements. Data correspond to the means of replicates for each determination as indicated above. Data were tested in their normality and equal variance by the Shapiro-Wilk test and then data were analyzed by a one way ANOVA. Significant differences among means were established by using the multiple comparisons test of Tukey's ( $P < 0.05$ ). All analyses were performed with Sigma Stat 2.0 software (SPSS, Chicago), where differences between the values were considered significant at  $P \leq 0.05$ .

### 4.3 Results

A statistically significant decrease in total biomass of leaves treated with 500 and 1500  $\mu$ M Mn concentrations was found compared to the control plants (18  $\mu$ M Mn), although a decrease of the total leaf biomass was found at all Mn concentrations used. In contrast, roots biomass exhibited significant decrease in all treatments ( $P < 0.05$ ) (Fig. 1). It should be noted, however, that while the decrease of leaf biomass was only 25% lower at the highest Mn concentration used (1500  $\mu$ M), the biomass of roots was more affected and demonstrated 2.5-fold decrease even at the lowest Mn concentration tested (50  $\mu$ M). This is consistent with the relative growth rate (RGR) data, where a considerably higher reduction of root RGR was observed compared to leave across the Mn treatments relative to control plants ( $P < 0.05$ ) (Table 1). In addition, visual symptoms of Mn toxicity (chlorosis) in leaves were observed predominantly in the highest Mn- treatment (data not shown).

Analysis of the total Mn amount in plants exposed to increasing Mn concentrations have demonstrated a gradual increase of Mn in leaves and roots (Fig. 2). However, while Mn

---

concentrations of roots reached values up to  $\sim 8100 \text{ mg kg}^{-1}$  Mn at the highest Mn-treatment, which represent a 40-fold increase of Mn, the increased accumulation of Mn in leaves was much lower (15-fold) compared to roots (Fig. 2). Thus, the differential effects of Mn treatments on the total biomass and RGR in leaves and roots could be explained by the higher accumulation of Mn in roots.

The effects of exposure to increasing Mn concentrations on the photochemical efficiency of PSII measured as  $F_v/F_m$  in *Arabidopsis* leaves are presented in Table 2. A time course measurements did not reveal any statistically significant changes of  $F_v/F_m$  values at the lower Mn doses (50  $\mu\text{M}$  and 500  $\mu\text{M}$ ) for the entire period of treatment compared to control plants (18  $\mu\text{M}$  Mn). At higher Mn concentrations (1000  $\mu\text{M}$  and 1500  $\mu\text{M}$ ) a small reduction (6%) of  $F_v/F_m$  was observed only at a later stage (days 15 and 21) of treatment, the statistically significant ( $P < 0.05$ ) differences being registered at the harvest time (21 days) (Table 2). Moreover, minimal Mn-induced changes were detected in the light response curves of photochemical quenching (qP), effective quantum yield of PSII ( $\Phi_{\text{PSII}}$ ) and PSII electron transport rates (ETR) even after 21 days of Mn-treatments with the highest dose tested (1500  $\mu\text{M}$ ) (data not shown).

In addition, thermoluminescence (TL) measurements were used as an alternative approach for assessing the effects of Mn-excess on the photosynthetic PSII-associated electron transfer reactions especially at its reducing side (Vass and Govindjee, 1996; Sane *et al.*, 2012). Since most of the photosynthetic TL components have been assigned to arise from the reversal of light-driven charge separation in PSII, TL properties of photosynthetic apparatus provide information on the activation energies associated with the back reactions of electron acceptors ( $Q_A$  and  $Q_B$ ) with the electron donors ( $S_2$  and  $S_3$ ) of PSII (Vass and Govindjee, 1996; Sane, 2004; Sane *et al.*, 2012). The temperature maxima ( $T_M$ ) of the TL peaks related to the recombination of these charge pairs reflect the activation energies and hence a measure of the redox potentials of the participating oxidized and reduced donors (de Vault and Govindjee, 1990). Typical TL glow curves representing  $S_2/S_3Q_B^-$  charge recombinations of control and Mn-treated *Arabidopsis* plants obtained following excitation with two consecutive saturating flashes are shown in Fig. 3. The experimental data summarized in Table 3 indicate that treatment with a Mn dose of 500  $\mu\text{M}$  did not exhibit significant differences of the TL peak position, while treatment with the highest

---

concentration (1500  $\mu\text{M}$ ) induced a low temperature shift of the  $T_M$  to 25.8°C compared to control plants. Besides this, the amplitudes and the integrated areas of the TL peaks representing the  $S_2/S_3Q_B^-$  charge recombination used for assessing the PSII photochemistry was not affected at 500  $\mu\text{M}$  Mn-excess, but the overall TL yield was significantly reduced (45%) in plants treated with 1500  $\mu\text{M}$  Mn compared to controls (Table 3).

To further test whether this small but significant Mn-induced effect on PSII photochemistry had physiological implications under additional stress conditions, *Arabidopsis* leaves subjected to different Mn treatments for 21 days were exposed to high light stress (Table 4). Indeed, the photoinhibitory effect on PSII, measured as a decrease in  $F_v/F_m$ , was much stronger (49%) at the highest Mn treatment (1500  $\mu\text{M}$ ) compared to control plants and 500  $\mu\text{M}$  Mn treated plants exhibited only a 15% decrease in  $F_v/F_m$  values after 4 h of exposure to high light (Table 4).

The extent of far-red light-induced absorbance decrease at 820 nm ( $\Delta A_{820-860}$ ) of *Arabidopsis* leaves (Klughammer and Schreiber, 1991; Ivanov *et al.*, 1998; Ivanov *et al.*, 2006a) was used to estimate the potential functional differences of PSI and photosynthetic electron transport pathways between plants exposed to different Mn treatments. Typical traces representing *in vivo* measurements of oxidation-reduction transients of  $P_{700}$  in control and Mn-treated plants are shown in Fig. 4. The relative amount of  $P_{700}^+$ , measured as  $\Delta A_{820-860}$ , gradually decreased with increasing Mn concentrations and was 30% lower in Mn-treated plants at the highest concentration used (1500  $\mu\text{M}$ ) (Fig. 4, Table 5). Concomitantly, kinetic measurements of dark re-reduction of  $P_{700}^+$  after turning off the far-red light, which is thought to reflect the extent and/or capacity for cyclic electron flow around PSI (Maxwell and Biggins, 1976; Ravenel *et al.*, 1994), indicated significantly slower (46%) re-reduction of  $P_{700}^+$  in Mn(1500  $\mu\text{M}$ )-treated compared to control plants (Table 5). In addition, the apparent electron donor pool size to PSI ( $e^-/P_{700}$ ) estimated by measuring single- and multiple-turnover flash-induced  $\Delta A_{820-860}$  under steady-state oxidation of PSI by far-red light (Asada *et al.*, 1993; Ivanov *et al.*, 1998) demonstrated a significant decrease in Mn-treated plants (Table 5). This indicates that the pool size of electrons that can be donated to photo-oxidized  $P_{700}$  ( $P_{700}^+$ ) from the stroma in control plants was 37% higher compared to plants treated with the highest Mn dose (13 electrons per  $P_{700}$ ) (Table 4).

---

The major photosynthetic components within the thylakoid membranes of control and Mn-treated *Arabidopsis* plants were compared by SDS-PAGE and immunodetection to quantify their relative abundance. Immunoblot analyses did not exhibit any significant Mn-stress induced changes in the relative abundance of PSII-associated polypeptides, as revealed by the densitometry analysis of the immunoblot bands for D1 (the PSII reaction center protein), Lhcb1 (major light harvesting protein of LHCII complex) and PsbO (extrinsic protein of the oxygen-evolving complex (OEC)) (Fig. 5A, B). The abundance of Rubisco was only marginally affected by the Mn treatments. In contrast, the abundance of reaction center polypeptides of PSI (PsaA and PsaB) was significantly reduced in Mn-treated compared with control thylakoids (Fig. 5A). The densitometric analysis demonstrated that the relative abundance of PsaA and PsaB in Mn-treated *Arabidopsis* was only about 20% and 60%, respectively of that observed in the control plants (Fig. 5B). Thus, the quantitative analysis of photosynthetic polypeptides clearly indicates that excess Mn has a greater effect on PSI-associated proteins rather than PSII. Consequently, the differences in the extent of P700 oxidation (P700<sup>+</sup>) and the kinetics of P700<sup>+</sup> reduction between control and Mn-treated plants are consistent with the lower levels of PSI-associated immunodetectable PsaA and PsaB protein complexes.

#### 4.4 Discussion

In agreement with a number of previous studies examining the effects of Mn stress on various plant species (Mora *et al.*, 2009; Alam *et al.*, 2006; Lei *et al.*, 2007; Doncheva *et al.*, 2009; Stoyanova *et al.*, 2009; Khabaz-Saberi *et al.*, 2010) *Arabidopsis* plants subjected to increasing Mn concentrations, also exhibited a reduction in dry weight of both shoots and roots at doses between 500 and 1500  $\mu\text{M}$  Mn (Fig. 1). The decline in biomass corresponded with a gradual increase of Mn concentrations in both shoot and roots of *Arabidopsis* subjected to excess Mn supply (Fig. 2). It should be noted that the decline of biomass was more pronounced in roots, where even at the lowest Mn treatment (50  $\mu\text{M}$ ) the dry weight was 2.5 lower compared to leaves. Similar results have been reported by Delhaize *et al.* (2007), where transporter proteins were implicated in the endogenous Mn tolerance of wild-type *Arabidopsis*. More recently, Mora *et al.* (2009) demonstrated that Mn-tolerant ryegrass cultivars accumulated higher Mn concentrations in roots than shoots,

---

while Mn sensitive cultivars exhibited a greater Mn translocation from roots to shoots. These results are also consistent with studies in legumes such as white clover (*Trifolium repens* L.) (Rosas *et al.*, 2007). However, in two contrasting populations of *Populus cathayana*, acclimated to wet and dry climate exposure to excess Mn caused an increase in Mn content of plant tissues especially in leaves and a visual symptoms of Mn toxicity (chlorosis) at high Mn concentrations (Lei *et al.*, 2007). The chlorosis observed in our experiments (data not shown) also correspond to a decreased amounts of both Chl *a* and Chl *b* could be due to a higher Mn accumulation in leaves after exposure to Mn excess, thus suggesting a damage to the photosynthetic apparatus as reported by Demirevska-Kepova *et al.* (2004).

Interestingly, while reduced CO<sub>2</sub> assimilation induced by excess Mn has been reported in many species and is considered one of the major physiological effects of Mn toxicity (Nable *et al.*, 1988; González and Lynch, 1997; Kitao *et al.*, 1997a; Lidon *et al.*, 2004; Macfie and Taylor, 1992; Li *et al.*, 2010), the functional integrity of the photosynthetic apparatus assessed by the maximum efficiency of PSII photochemistry ( $F_v/F_m$ ) did not decline as a result of exposure to excess Mn in various plant species (Nable *et al.*, 1988; Kitao *et al.*, 1997b; Subrahmanyam and Rathore, 2000; Hajiboland and Hasani, 2007; Doncheva *et al.*, 2009). However, some studies have reported a substantial decrease in  $F_v/F_m$  as a result of excess Mn treatment in *Citrus* species (Papadakis *et al.*, 2007; Li *et al.*, 2010), rice (Lidon *et al.*, 2004), Mn-sensitive maize (Doncheva *et al.*, 2009) and cucumber (Feng *et al.*, 2009). Furthermore, Kitao *et al.* (1997b) have shown that the potential maximum efficiency of PSII photochemistry ( $F_v/F_m$ ) is not affected by excess Mn in white birch, although the reduction state of PSII primary electron acceptor ( $Q_A$ ) was increased at high Mn concentrations. These results clearly indicate that the excess Mn- induced decline in CO<sub>2</sub> assimilation may or may not be accompanied by changes in PSII photochemistry and this response is species dependent. The experimental data presented in this study also failed to demonstrate any substantial effects of excess Mn within a wide range of Mn concentrations on the maximum efficiency of PSII photochemistry in *Arabidopsis* plants (Table 2). Assessing the relative abundance of PSII-associated proteins also showed no changes in the immuno-detectable amounts of PSII-reaction center protein D1, the light harvesting chlorophyll-protein complex of PSII (Lhcb1) and the manganese stabilizing

---

33kDa-protein of the water splitting complex of PSII (PsbO) polypeptides (Fig. 5) in plants exposed to high Mn concentrations compared to control *Arabidopsis*. However, the observed low temperature shift of the  $S_{2,3}Q_B^-$  charge recombination and much lower overall TL emission implies lower redox potential of  $Q_B$  (Fig. 3; Table 3) confirms the suggestion of altered reduction state of PSII acceptor side in Mn-stressed plants (Kitao *et al.*, 1997b). Since  $Q_A$  is in quasi-equilibrium with  $Q_B$  and the PQ pool, our results imply that lowering the redox potential of  $Q_B$  will decrease the probability for forward electron transfer between the two quinone acceptors by shifting the redox equilibrium between  $Q_A^-Q_B$  and  $Q_AQ_B^-$  towards  $Q_A^-Q_B$  (Minagawa *et al.*, 1999; Ivanov *et al.*, 2002; 2003) in plants exposed to high Mn concentrations.

In addition to the lack of significant inhibitory effects of excess Mn on PSII photochemistry discussed above, an earlier study reported that the photochemistry of photosystem I (PSI) and the photosynthetic electron transport were not significantly affected during early development of Mn toxicity in tobacco plants (Nable *et al.*, 1988). However, a decreased Hill activity in isolated chloroplasts was found in mungbean leaves exposed to toxic Mn concentrations (Sinha *et al.*, 2002). More recently, Li *et al.* (2010) have suggested that Mn excess can effectively impair the whole photosynthetic electron transport chain, thus restricting the production of reducing equivalents and limiting the rate of  $CO_2$  assimilation in *Citrus grandis* seedlings. Despite these few studies, the potential effect(s) of excess Mn on the functional/structural integrity of PSI remains elusive. To our knowledge, the results presented in this study are the first report of an *in vivo* assessment of high Mn concentrations on PSI photochemistry. In contrast to PSII photochemistry, *in vivo* measurements of the oxidation state of P700(P700<sup>+</sup>) (Klughammer and Schreiber, 1991; Ivanov *et al.*, 1998), the primary donor of PSI demonstrated that the relative amount of oxidizable P700(P700<sup>+</sup>) decreased by 30% in Mn-treated *Arabidopsis* plants at concentrations above 1000  $\mu$ M (Fig. 4, Table 5). The functional impairment of PSI photochemistry by excess Mn was accompanied by a significant reduction in the abundance of PSI reaction center polypeptides (PsaA and PsaB) (Fig. 5). This clearly indicates that the major target of Mn toxicity within the photosynthetic electron transport chain of *Arabidopsis* is PSI rather than PSII related components. The reduced amounts of PSI reaction center polypeptides PsaA and PsaB would imply acceptor side limitations of the

---

photosynthetic electron transport and this could explain the increased reduction state  $Q_A$  in Mn-stressed plants reported earlier (Kitao *et al.*, 1997b). Moreover, decreased expression of another Fe-containing chloroplastic protein precursor, ferredoxin-1 serving as a terminal electron acceptor of the photosynthetic electron transport observed in Mn-treated young rice leaves also supports Mn-induced limitations at the acceptor side of PSI (Führs *et al.*, 2010).

One of the major mechanisms considered for Mn toxicity involves the inhibition of other essential cations including Fe, thus suggesting that a Mn-induced Fe-deficiency may play a key role in the physiological responses to excess Mn (Foy *et al.*, 1978; Foy, 1984; Kohno *et al.*, 1984). More recently, chloroplast alterations in maize plants exposed to excess Mn (Doncheva *et al.*, 2009) and Mn-toxicity in young rice leaves (Führs *et al.*, 2010) have been also ascribed to Mn-induced Fe deficiency rather than to direct Mn-induced oxidative stress. Given that about 80% of the plant Fe is located in the chloroplast (Terry and Abadia, 1986) and the functional photosynthetic apparatus requires 22-23 iron atoms, of which PSI is the most Fe-abundant component (Ferreira and Straus, 1994), it seems reasonable to assume that the observed lower abundance of PSI reaction center polypeptides and the associated decline of PSI photochemistry in Mn-treated *Arabidopsis* plants were consequences of a Mn-induced moderate Fe deficiency.

Although light energy is important for photosynthetic processes in plants, an excess of light can be also harmful because it can result in photoinhibition, which can be exacerbated when it is combined with other stresses (Powles, 1984; Aro *et al.*, 1993; Sonoike, 1996). Photoinhibition is a complex phenomenon that may cause damage to the photosynthetic apparatus reducing the photosynthetic efficiency when light conditions exceed the photon requirements for photosynthesis (Murata *et al.*, 2007). It is considered that PSII is the main site of photoinhibition (Aro *et al.*, 1993; Sonoike, 1996; Takahashi and Murata, 2008), being more unstable than PSI, because the D1 protein, one of the two major heterodimeric polypeptides of the PSII reaction center complex, has a very high, light-dependent turnover rate (Scheller and Haldrup, 2005; Aro *et al.*, 1993; Burnap, 2004; Takahashi and Murata, 2008). Earlier reports have shown that the susceptibility to Mn toxicity is strongly dependent on the light intensity and exposure of Mn-treated plants to high light can exacerbate the toxic effect of Mn. (Nable *et al.*, 1988; Horiguchi, 1988; González *et al.*,

---

1998; Clair and Lynch, 2004; Hajiboland and Hasani, 2007). Our results also demonstrate that *Arabidopsis* plants pre-disposed to high Mn concentrations are more susceptible to photoinhibitory damage of PSII photochemistry in a concentration dependent manner (Table 4).

Apart from the radiation less dissipation of excess excitation energy in the chlorophyll pigment bed of LHCII, associated with the formation of the xanthophylls pigment zeaxanthin (Z), which is considered one of the major protective mechanisms against photoinhibitory damage (Horton *et al.*, 1996; Niyogi, 1999), PSI-dependent cyclic electron transport (CET) has been also suggested to play a significant role in preventing the photoinhibitory damage of the photosynthetic apparatus during exposure of plants to high light conditions (Munekage *et al.*, 2002; Takahashi *et al.*, 2009). Considering the increased re-reduction rate of P700<sup>+</sup> in *Arabidopsis* (Table 5), we believe that the higher susceptibility of plants exposed to excess Mn to photoinhibition (Table 4) might be due to lower capacity of PSI-driven cyclic electron flow under conditions of Mn toxicity.

In summary, the results presented in this research demonstrate for the first time that exposure of *Arabidopsis* plants to excess Mn causes specific negative effects on the abundance of polypeptides comprising the reaction center of PSI, thus resulting in decreased PSI photochemistry and lower capacity for CET, which may be due to a Mn-induced Fe-deficiency and may have critical physiological implications under conditions of Mn toxicity in higher plants.

**Acknowledgements:** This work was supported by Natural Sciences and Engineering Research Council of Canada (N.P.A.H.). CONICYT Internship: “Pasantias Doctorales en el Extranjero 2010”. Department of Biology and the Biotron, University of Western Ontario, London, Ontario, Canada. Fondecyt Project N° 1080372, Government of Chile.

---

#### 4.5 References

**Alam S, Kodama R, Akiha F, Kamei S, Kawai S.** 2006. Alleviation of manganese phytotoxicity in barley with calcium. *Journal of Plant Nutrition* **29**, 59-74.

**Arnon DI.** 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* **24**:1-15.

**Aro E-M, Virgin I, Andersson B.** 1993. Photoinhibition of photosystem II. Inactivation, protein damage and turnover. *Biochimica et Biophysica Acta* **1143**, 113-134.

**Asada K, Heber U, Schreiber U.** 1993. Electron flow to the intersystem chain from stromal components and cyclic electron flow in maize chloroplasts, as determined in intact leaves by monitoring redox change of P700 and chlorophyll fluorescence. *Plant and Cell Physiology* **34**, 39-50.

**Bowler C, VanCamp W, VanMontagu M, Inze D.** 1994. Superoxide dismutase in plants. *Critical Reviews in Plant Sciences* **13**, 199-218.

**Bradl H.** 2004. Adsorption of heavy metal ions on soils and soils constituents. *Journal of Colloids and Interface Science* **277**, 1-18.

**Burnap RL.** 2004. D1 protein processing and Mn cluster assembly in light of the emerging Photosystem II structure. *Physical Chemistry Chemical. Physics* **6**, 4803-4809.

**Burnell JN.** 1988. The biochemistry of manganese in plants. In: Graham RD, Hannam J, Uren NC, eds. *Manganese in Soils and Plants*. Dordrecht, The Netherlands: Kluwer Academic Publishers, 125-133.

**Cailliatte R, Schikora A, Briat JF, Mari S, Curie C.** 2010. High-affinity manganese uptake by the metal transporter NRAMP1 is essential for *Arabidopsis* growth in low manganese conditions. *Plant Cell* **22**: 904–917.

---

**Chatterjee C, Nautiyal N, Agarwala SC.** 1994. Influence of changes in manganese and magnesium supply on some aspects of wheat physiology. *Soil Science and Plant Nutrition* **40**, 191-197.

**Clair SB St, Lynch JP.** 2004. Photosynthetic and antioxidant enzyme responses of sugar maple and red maple seedlings to excess manganese in contrasting light environments. *Functional Plant Biology* **31**, 1005-1014.

**de Vault D, Govindjee.** 1990. Photosynthetic glow peaks and their relationship with the free energy changes. *Photosynthesis Research* **24**, 175-181.

**Delhaize E, Gruber BD, Pittman JK, White RG, Leung H, Miao Y, Jiang L, Ryan PR, Richardson AE.** 2007. A role for the AtMTP11 gene of *Arabidopsis* in manganese transport and tolerance. *Plant Journal* **51**, 198-210.

**Demirevska-Kepova K, Simova-Stoilova L, Stoyanova Z, Holzer R, Feller U.** 2004. Biochemical changes in barley plants after excessive supply of copper and manganese. *Environmental and Experimental Botany* **52**, 253-266.

**Doncheva S, Poschenrieder C, Stoyanova Z, Georgieva K, Velichkova M, Barceló J.** 2009. Silicon amelioration of manganese toxicity in Mn-sensitive and Mn-tolerant maize varieties. *Environmental and Experimental Botany* **65**, 189-197.

**Enami I, Okumura A, Nagao R, Suzuki T, Iwai M, Shen JR.** 2008. Structures and functions of the extrinsic proteins of photosystem II from different species. *Photosynthesis Research* **98**, 349-363.

**Fecht-Christoffers MM, Maier P, Horst W.** 2003. Apoplastic peroxidase and ascorbate are involved in manganese toxicity and tolerance of *Vigna unguiculata*. *Physiologia Plantarum* **117**, 237-244.

---

**Feng J-P, Shi Q-H, Wang X-F.** 2009. Effects of exogenous silicon on photosynthetic capacity and antioxidant enzyme activities in chloroplast of cucumber seedlings under excess manganese. *Agricultural Science in China* **8**, 40-50.

**Fernando DR, Baker AJM, Woodrow IE.** 2009. Physiological responses in *Macadamia integrifolia* on exposure to manganese treatment. *Australian Journal of Botany* **57**, 406-413.

**Ferreira F, Straus NA.** 1994. Iron deprivation in cyanobacteria. *Journal of Applied Phycology* **6**, 199-210.

**Ferreira KN, Iverson TM, Maghlaoui K, Barber J, Iwata S.** 2004. Architecture of the photosynthetic oxygen-evolving center. *Science* **303**, 1831-1838.

**Foy CD, Chaney RL, White MC.** 1978. The physiology of metal toxicity in plants. *Annual Review of Plant Physiology* **29**, 511-566.

**Foy CD.** 1984. Physiological effects of hydrogen, aluminum, and manganese toxicities in acid soils. *Soil Acidity and Liming Madison: American Society of Agronomy* **2**, 57-97.

**Führs H, Hartwig M, Molina LEB, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ.** 2008. Early manganese-toxicity response in *Vigna unguiculata* L. – a proteomic and transcriptomic study. *Proteomics* **8**, 149-159.

**Führs H, Behrens C, Gallien S, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ.** 2010. Physiological and proteomic characterization of manganese sensitivity and tolerance in rice (*Oryza sativa*) in comparison with barley (*Hordeum vulgare*). *Annals of Botany* **105**, 1129-1140.

**Genty B, Briantais J-M, Baker NR.** 1989. The relationship between quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica et Biophysica Acta* **990**, 87-92.

---

**González A, Lynch JP.** 1997. Effects of manganese toxicity on leaf CO<sub>2</sub> assimilation of contrasting common bean genotypes. *Physiologia Plantarum* **101**, 872-880.

**González A, Steffen KL, Lynch JP.** 1998. Light and excess manganese. Implication for oxidative stress in common bean. *Plant Physiology* **18**, 493-504.

**González A, Lynch JP.** 1999. Subcellular and tissue Mn compartmentation in bean leaves under Mn toxicity stress. *Australian Journal of Plant Physiology* **26**, 811-822.

**Hajiboland R, Hasani BD.** 2007. Effect of Cu and Mn toxicity on chlorophyll fluorescence and gas exchange in rice and sunflower under different light intensities. *Journal of Stress Physiology and Biochemistry* **3**, 4-17.

**Hankamer B, Barber J, Boekema EJ.** 1997. Structure and membrane organization of photosystem II in green plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **48**, 641-671.

**Horiguchi T.** 1988. Mechanism of manganese toxicity and tolerance of plants. VII. Effect of light intensity on manganese-induced chlorosis. *Journal of Plant Nutrition* **11**, 235-245.

**Horton P, Ruban AV, Walters RG.** 1996. Regulation of light harvesting in green plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **47**, 655-684.

**Houtz RL, Nable RO, Cheniae GM.** 1988. Evidence for effects on the *in vivo* activity of ribulose-bisphosphate carboxylase/oxygenase during development of Mn toxicity in tobacco. *Plant Physiology* **86**, 1143-1149.

**Ivanov AG, Morgan R, Gray GR, Velitchkova MY, Huner NPA.** 1998. Temperature/light dependent development of selective resistance to photoinhibition of photosystem I. *FEBS Letters* **430**, 288-292.

---

**Ivanov AG, Sane PV, Zeinalov Y, Malmberg G, Gardeström P, Huner NPA, Öquist G.** 2001. Photosynthetic electron transport adjustments in overwintering Scots pine (*Pinus sylvestris* L.). *Planta* **213**, 575-585.

**Ivanov AG, Sane PV, Zeinalov Y, Simidjiev I, Huner NPA, Öquist G.** 2002. Seasonal responses of photosynthetic electron transport in Scots pine (*Pinus sylvestris* L.) studied by thermoluminescence. *Planta* **215**, 457-465.

**Ivanov AG, Sane P, Hurry V, Krol M, Sveshnikov D, Huner NPA, Öquist G.** 2003. Low-temperature modulation of the redox properties of the acceptor side of photosystem II: photoprotection through reaction centre quenching of excess energy. *Physiologia Plantarum* **119**, 376-383.

**Ivanov AG, Hendrickson L, Krol M, Selstam E, Öquist G, Hurry V, Huner NPA.** 2006a. Digalactosyl-diacylglycerol deficiency impairs the capacity for photosynthetic intersystem electron transport and state transitions in *Arabidopsis thaliana* due to photosystem I acceptor-side limitations. *Plant and Cell Physiology* **47**, 1146-1157.

**Ivanov AG, Sane PV, Krol M, Gray GR, Balseris A, Savitch LV, Öquist G, Hüner NPA.** 2006b. Acclimation to temperature and irradiance modulates PSII charge recombination. *FEBS Letters* **580**, 2797-2802.

**Jordan DB, Ogren WL.** 1983. Species variation in kinetic properties of Rubisco. *Archives of Biochemistry and Biophysics* **227**, 425-433.

**Kern J, Renger G.** 2007. Photosystem II: Structure and mechanism of the water:plastoquinone oxidoreductase. *Photosynthesis Research* **94**, 183-202.

**Khabaz-Saberi H, Rengel Z, Wilson R, Setter TL.** 2010. Variation of tolerance to manganese toxicity in Australian hexaploid wheat. *Journal of Plant Nutrition and Soil Science* **173**, 103-112.

---

**Kitao M, Lei TT, Koike T.** 1997a. Comparison of photosynthetic responses to manganese toxicity of deciduous broad leaved trees in northern Japan. *Environmental Pollution* **97**, 113-118.

**Kitao M, Lei TT, Koike T.** 1997b. Effects of manganese toxicity on photosynthesis of white birch (*Betula platyphylla* var. *japonica*) seedlings. *Physiologia Plantarum* **101**, 249-256.

**Klughammer C, Schreiber U.** 1991. Analysis of light-induced absorbancy changes in the near-infrared spectral region. 1. Characterization of various components in isolated chloroplasts. *Zeitschrift für Naturforschung C* **46**, 233-244.

**Kohno Y, Foy CD, Fleming AL, Krizek DT.** 1984. Effect of Mn concentration on the growth and distribution of Mn and Fe in two bush bean cultivars grown in solution culture. *Journal of Plant Nutrition* **7**, 547-566.

**Kramer DM, Johnson G, Kiirats O, Edwards GE.** 2004. New fluorescence parameters for the determination of QA redox state and excitation energy fluxes. *Photosynthesis Research* **79**, 209-218.

**Krol M, Ivanov AG, Jansson S, Kloppstech K, Huner NPA.** 1999. Greening under high light and cold temperature affects the level of xanthophyll-cycle pigments, early light-inducible proteins, and light harvesting polypeptides in wild-type barley and the *Chlorina f2* mutant. *Plant Physiology* **120**, 193-203.

**Laemmli U.** 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.

**Lei Y, Korpelainen H, Li C.** 2007. Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. *Chemosphere* **68**, 686-694.

---

**Li Q, Chen L-S, Jiang H-X, Tang N, Yang L-T, Lin Z-H, Li Y, Yang G-H.** 2010. Effects of manganese-excess on CO<sub>2</sub> assimilation ribulose-1,5-bisphosphate carboxylase/oxygenase, carbohydrates and photosynthetic electron transport of leaves, and antioxidant systems of leaves and roots in *Citrus grandis* seedlings. *BMC Plant Biology* **10**, 42.

**Lidon FC, Barreiro MG, Ramalho JC.** 2004. Manganese accumulation in rice: implications for photosynthetic functioning. *Journal of Plant Physiology* **161**, 1235-1244.

**Macfie S, Taylor G.** 1992. The effects of excess manganese on photosynthetic rate and concentration of chlorophyll in *Triticum aestivum* grown in solution culture. *Physiologia Plantarum* **85**, 467-475.

**McDaniel KL, Toman FR.** 1994. Short-term effects of manganese toxicity on ribulose-1,5-bisphosphate carboxylase in tobacco chloroplasts. *Journal of Plant Nutrition* **17**, 523-536.

**Marschner H.** 1995. *Mineral Nutrition of Higher Plants*. Academic Press, San Diego, 889 p.

**Maxwell PC, Biggins J.** 1976. Role of cyclic electron transport in photosynthesis as measured by turnover of P700 *in vivo*. *Biochemistry* **15**, 3975-3981.

**Millaleo R, Reyes-Díaz M, Ivanov AG, Mora ML, Alberdi M.** 2010. Manganese as essential and toxic element for plants: Transport, accumulation and resistance mechanisms. *Journal of Soil Science and Plant Nutrition* **10**, 476-494.

**Minagawa J, Narusaka Y, Inoue Y, Satoh K.** 1999. Electron transfer between Q<sub>A</sub> and Q<sub>B</sub> in photosystem II is thermodynamically perturbed in phototolerant mutants of *Synechocystis sp.* PCC 6803. *Biochemistry* **38**, 770-775.

---

**Mora M, Rosas A, Ribera A, Rengel R.** 2009. Differential tolerance to Mn toxicity in perennial ryegrass genotypes: involvement of antioxidative enzymes and root exudation of carboxylates. *Plant Soil* **320**, 79-89.

**Mukhopadhyay MJ, Sharma A.** 1991. Manganese in cell metabolism of higher plants. *Botanical Review* **57**, 117-149.

**Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T.** 2002. *PGR5* is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. *Cell* **110**, 361-371.

**Murata N, Takahashi S, Nishiyama Y, Allakhverdiev SI.** 2007. Photoinhibition of photosystem II under environmental stress. *Biochimica et Biophysica Acta* **1767**, 414-421.

**Nable RO, Houtz RL, Cheniae GM.** 1988. Early inhibition of photosynthesis during development of Mn toxicity in tobacco. *Plant Physiology* **86**, 1136-1142.

**Niyogi KK.** 1999. Photoprotection revisited: Genetic and molecular approaches. *Annual Review of Plant Physiology and Plant Molecular Biology* **50**, 333-359.

**Ohki K.** 1984. Manganese deficiency and toxicity effects on growth, development and nutrient composition in wheat. *Agronomy Journal* **76**, 213-218.

**Panda S, Mishra AK, Biswal UC.** 1987. Manganese induced peroxidation of thylakoid lipids and changes in chlorophyll-a fluorescence during aging of cell free chloroplasts in light. *Phytochemistry* **26**, 3217-3219.

**Papadakis IE, Giannakoula A, Therios IN, Bosabalidis AM, Moustakas M, Nastou A.** 2007. Mn-induced changes in leaf structure and chloroplast ultrastructure of *Citrus volkameriana* (L) plants. *Journal of Plant Physiology* **164**, 100-103.

---

**Powles SB.** 1984. Photoinhibition of photosynthesis induced by visible light. Annual Review of Plant Physiology **35**, 15-44.

**Ravenel J, Peltier G, Havaux M.** 1994. The cyclic electron pathways around photosystem I in *Chlamydomonas reinhardtii* as determined *in vivo* by photoacoustic measurements of energy storage. Planta 193, 251-259.

**Rosas A, Rengel Z, Mora M.** 2007. Manganese supply and pH influence growth, carboxylate exudation and peroxidase activity of ryegrass and white clover. Journal of Plant Nutrition **30**, 253-270.

**Rosso D, Bode R, Li W, Krol M, Saccon D, Wang S, Schillaci LA, Rodermeel SR, Maxwell DP, Hüner NPA.** 2009. Photosynthetic redox imbalance governs leaf sectoring in the *Arabidopsis thaliana* variegation mutants *immutans*, *spotty*, *var1*, and *var2*. The Plant Cell **21**, 3473-3492.

**Rutherford AW, Boussac A.** 2004. Water photolysis in biology. Science **303**, 1782-1784.

**Sadzawka A, Carrasco M, Grez R, Mora M.** 2004. Métodos de análisis de tejidos vegetales. Comisión de normalización y acreditación. Sociedad Chilena de la Ciencia del Suelo (SCCS). Available at [http://www.inia.cl/platina/descarga/docs/libros/asadzawka\\_tejevveg\\_2004.pdf](http://www.inia.cl/platina/descarga/docs/libros/asadzawka_tejevveg_2004.pdf).

**Sane PV.** 2004. Thermoluminescence: A technique for probing photosystem II, in: R. Carpentier (Ed.), Photosynthesis Research Protocols, Humana Press, Totowa, NJ, pp. 229-248.

**Sane PV, Ivanov AG, Öquist G, Hüner NPA.** 2012. Thermoluminescence, in: J.J. Eaton-Rye, B.C. Tripathy, T.D. Sharkey (Eds.), Photosynthesis: Plastid Biology, Energy Conversion and Carbon Assimilation, Advances in Photosynthesis and Respiration Vol. 34, Springer Science+Business Media B.V., pp. 445-474.

---

**Schaaf G, Catoni E, Fitz M, Schwake R, Schneider A, von Wirén N, Frommer WB.** 2002. A putative role for vacuolar calcium/manganese proton antiporter *AtCAX2* in heavy metal detoxification. *Plant Biology* **2**, 612-618.

**Scheller H, Haldrup A.** 2005. Photoinhibition of photosystem I. *Planta* **221**, 5-8.

**Schreiber U, Bilger W, Neubauer C.** 1994. Chlorophyll fluorescence as a noninvasive indicator for rapid assessment of in vivo photosynthesis. *In* ED Schulze, MM Caldwell, eds, *Ecophysiology of Photosynthesis*. Springer-Verlag, Berlin, pp 49-70.

**Sinha S, Mukherji S, Dutta J.** 2002. Effect of manganese toxicity on pigment content, Hill activity and photosynthetic rate of *Vigna radiata* L. Wilczek seedlings. *Journal of Environmental Biology* **23**, 253-257.

**Sonoike K.** 1996. Photoinhibition of photosystem I: its physiological significance in the chilling sensitivity of plants, *Plant and Cell Physiology* **37**, 239-247.

**Stoyanova Z, Poschenrieder C, Tzvetkova N, Doncheva S.** 2009. Characterization of the tolerance to excess manganese in four maize varieties. *Soil Science and Plant Nutrition* **55**, 747-753.

**Subrahmanyam D, Rathore VS.** 2000. Influence of manganese toxicity on photosynthesis in ricebean (*Vigna umbellata*) seedlings. *Photosynthetica* **38**, 449-453.

**Takahashi S, Murata N.** 2008. How do environmental stresses accelerate photoinhibition? *Trends in Plant Science* **13**, 178-182.

**Takahashi S, Milward SE, Fan DY, Chow WS, Badger MR.** 2009. How does cyclic electron flow alleviate photoinhibition in *Arabidopsis*? *Plant Physiology* **149**, 1560-1567.

---

**Terry N, Abadia J.** 1986. Function of iron in chloroplasts. *Journal of Plant Nutrition* **9**, 609-646.

**Towbin H., Staehelin T., Gordon J.** 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Science USA* **76**, 4350-4354.

**Vaughn KC, Duke SO.** 1984. Function of polyphenol oxidase in higher plants. *Physiologia Plantarum* **60**, 106-112.

**van Kooten O, Snel JFH.** 1990. The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosynthesis Research* **25**, 147-150.

**Vass I, Govindjee** 1996. Thermoluminescence from the photosynthetic apparatus. *Photosynthesis Research* **48**, 117-126.

---

**Table 1.** Relative growth rates (RGR) of *Arabidopsis thaliana* (leaves and roots) growing at increased Mn treatments under acidic conditions (pH 5.3) for 21 d in comparison with the control. Different lower-case letters indicate statistically significant differences between the Mn treatments ( $P < 0.05$ ).

---

<b>Relative growth rate (RGR)</b>		
<b>(g g<sup>-1</sup> d<sup>-1</sup> DW)</b>		
<b>Mn treatments</b>	<b>Leaves</b>	<b>Roots</b>
C	0.069 ± 0.001 a	0.206 ± 0.006 a
50 µM	0.068 ± 0.003 a	0.071 ± 0.005 b
500 µM	0.046 ± 0.006 b	0.053 ± 0.006 b
1000 µM	0.059 ± 0.003 ab	0.064 ± 0.007 b
1500 µM	0.040 ± 0.004 b	0.057 ± 0.004 b

---

**Table 2.** Maximal photochemical efficiency of PSII (Fv/Fm) in leaves of *Arabidopsis thaliana* measured at 0, 5, 10, 15 and 21 days subjected to increasing Mn treatments ( $\mu\text{M}$ ). Results are means of 5 repetitions of two plants each  $\pm$  SE. Different lower case indicate significant differences between the Mn treatments at the same time, while different upper-case letters indicate significant differences between the times at a same Mn treatment ( $P < 0.05$ ).

<b>Fv/Fm</b>						
<b>Mn treatments</b>	<b>Time (days)</b>					
	<b>(<math>\mu\text{M}</math>)</b>	0	5	10	15	21
<b>C</b>		0.803 aA	0.803 aA	0.803 aA	0.803 aA	0.803 aA
<b>50</b>		0.806 aA	0.799 aA	0.800 aA	0.800 aA	0.800 aA
<b>500</b>		0.803 aA	0.801 aA	0.804 aA	0.787 bA	0.791 aA
<b>1000</b>		0.797 aA	0.805 aA	0.800 aA	0.778 bB	0.773 bB
<b>1500</b>		0.802 aA	0.798 aA	0.775 bB	0.769 bcBC	0.756 cC

---

**Table 3.** Characteristic thermoluminescence (TL) peak emission temperatures ( $T_M$ ) and the overall TL emission area (A) of S<sub>2</sub>/S<sub>3</sub>Q<sub>B</sub>- glow peaks of control and Mn-treated (21 days) *Arabidopsis* plants. The samples (leaf disks) were dark adapted for 30 min then cooled to 2°C and subsequently illuminated with two single turnover flashes of white light. The peak areas are presented as a percentage of the total TL light emission in control leaves. Mean values ± SE were calculated from 4 independent experiments.

---

<b>Mn treatments</b>		
<b>(<math>\mu</math>M)</b>	<b><math>T_M</math> (<math>^{\circ}</math>C)</b>	<b>A (%)</b>
<b>Control</b>	28.7 ± 1.1	100.0
<b>500 <math>\mu</math>M</b>	29.1 ± 1.3	106.2 ± 7.6
<b>1500 <math>\mu</math>M</b>	25.8 ± 0.9	54.5 ± 7.9

---

**Table 4.** Effect of high light treatments ( $1000 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ , PFD, for 1, 2 and 4 hours) on the maximal photochemical efficiency of PSII measured as  $F_v/F_m$  in control *Arabidopsis thaliana* leaves and plants exposed for 21 days to different Mn doses. Results are means of 5 repetitions in three leaves of two plants each  $\pm$  SE. Time 0 was measured in plants subjected to  $250 \mu\text{mol m}^{-2} \text{s}^{-1}$  PFD. Different upper-case letters indicate significant differences between the times of exposure at the same Mn treatment ( $P < 0.05$ ).

Irradiance exposition time (h)	$F_v/F_m$		
	Control	500 $\mu\text{M}$ Mn	1500 $\mu\text{M}$ Mn
0	$0.815 \pm 0.003^A$	$0.812 \pm 0.004^A$	$0.780 \pm 0.014^A$
1	$0.761 \pm 0.012^A$	$0.750 \pm 0.002^A$	$0.677 \pm 0.014^B$
2	$0.752 \pm 0.007^A$	$0.731 \pm 0.002^A$	$0.619 \pm 0.028^B$
4	$0.685 \pm 0.036^A$	$0.689 \pm 0.004^A$	$0.388 \pm 0.042^C$

**Table 5.** Effects of Mn treatments on far-red light induced steady state oxidation of P700 ( $\Delta A_{820-860}$ , P700<sup>+</sup>), the relative intersystem electron donor pool size to PSI ( $e^-$  /P700) and half times for P700<sup>+</sup> reduction ( $t_{1/2}$ ) of *Arabidopsis* leaves at 21 days of treatment. Different lower-case letters indicate significant differences between the Mn treatments ( $P < 0.05$ ).

Mn Treatments ( $\mu\text{M}$ )	$\Delta A_{820-860}$ (P700 <sup>+</sup> )	$e^-$ /P700 ( $\text{MT}_{\text{AREA}}/\text{ST}_{\text{AREA}}$ )	$t_{1/2}$ (s)
Control	423.2 $\pm$ 7.9 a	20.7 $\pm$ 0.9 a	0.670 $\pm$ 0.05 b
50	416.4 $\pm$ 8.3 a	15.7 $\pm$ 0.6 b	0.812 $\pm$ 0.06 ab
500	368.4 $\pm$ 8.3 b	16.4 $\pm$ 0.6 b	0.938 $\pm$ 0.08 ab
1000	293.4 $\pm$ 8.4 c	15.6 $\pm$ 0.5 b	0.908 $\pm$ 0.07 ab
1500	292.9 $\pm$ 4.2 c	13.0 $\pm$ 0.3 c	0.984 $\pm$ 0.08 a

---

**Figure legends:**


**Figure 1.** Total biomass (leaves and roots, g DW) of *Arabidopsis thaliana* plants subjected to Mn treatments ( $\mu\text{M}$ ) at the end of experiment (21 days). Different lower-case letters indicate significant differences among the Mn treatments ( $P \leq 0.05$ ).

**Figure 2.** Mn concentration in leaves and roots of *Arabidopsis thaliana* plants subjected to Mn treatments ( $\mu\text{M}$ ) at the end of experiment (21 days). Different lower-case letters indicate statistically significant differences among the Mn treatments ( $P \leq 0.05$ ).


**Figure 3.** Thermoluminescence (TL) glow curves of  $S_2/S_3Q_B^-$  charge recombinations in control (solid lines) and Mn-treated ( $500 \mu\text{M Mn}^{2+}$  - dashed lines,  $1500 \mu\text{M Mn}^{2+}$  - dotted lines) *Arabidopsis* leaves after illumination with two single turnover flashes. TL glow curves were recorded immediately after illumination. The presented glow curves are averages from 4 independent measurements.

**Figure 4.** Typical traces of *in vivo* measurements of P700 oxidation by far-red light (FR) in control (A,  $18 \mu\text{M Mn}$ ) and Mn-treated (B,  $1500 \mu\text{M Mn}$ ) *Arabidopsis* plants. After reaching a steady state level of P700<sup>+</sup> by FR light, single turnover (ST) and multiple turnover (MT) pulses of white light were applied. Arrows indicate application of ST, MT, FR light sources. The measurements were performed at the growth temperature of  $20^\circ\text{C}$ .

**Figure 5.** A - Representative immunoblots of SDS-PAGE separated proteins from thylakoid membranes probed with antibodies raised against PSI (PsaA and PsaB), PSII (D1, PsbO, Lhcb1)-associated polypeptides and Rubisco (PbcL) in control (C,  $18 \mu\text{M Mn}$ ) and *Arabidopsis* plants treated with  $500$  and  $1500 \mu\text{M Mn}$ . B - The relative polypeptide abundance was quantified by densitometric analysis (area x intensity bands) of the immunoblots and the presented data were normalized to the relative abundance of PsaA, PsaB, D1, PsbO, Lhcb1 and PbcL in control *Arabidopsis* plants. Mean values  $\pm$  SE were calculated from 3 independent experiments. Different lower-case letters indicate significant differences between the Mn treatments ( $P < 0.05$ ).



**Figure 1**



**Figure 2**




Figure 3




Figure 4




Figure 5.



Chapter 5:  
General discussion

---

## 5.1 General discussion

In this thesis we analyzed biochemical and physiological responses of plants subjected to increasing Mn treatments growing in nutrient solution. In Chapter 3, three cultivars of highbush blueberry (*Vaccinium corymbosum* L.): ‘Legacy’, ‘Brigitta’ and ‘Bluegold’ were established under greenhouse conditions in nutrient solution with increasing Mn treatments: 2  $\mu$ M (Control), 10, 50, 250, 500 and 1000  $\mu$ M Mn. The pH used was 4.8. Results showed that shoots dry weight (DW) of Legacy and Brigitta were unchanged, whereas Bluegold had a significant and gradual reduction (~50%) of shoot DW with increasing Mn treatments (See Model; Fig. 5.1). Decreases in DW due to Mn excess are also observed in other plant species as rice (Horiguchi, 1988), ryegrass (Mora et al., 2009), lucerne (Gherardi and Rengel, 2004), pummel (Li et al., 2010), *Cucumis* species (Shi et al., 2005; 2006), maize (Stoyanova et al., 2009) and barley (Alam et al., 2006). According Rosas et al. (2007, 2011), detrimental effects on biomass are observed in prairie species subjected to Mn excess with stronger decrease in shoots than in roots, with visual symptoms on leaves. Similarly in maize, the DW of Mn-sensitive maize cultivar was affected mainly in shoots than roots. This DW reduction is a typical injury caused by Mn excess. Thus, this symptom is different to other metals behavior (i.e. copper and aluminum), which tissues DW are reduced significantly at root level (Doncheva et al., 2009). In our study, Legacy showed a high Mn content in roots compared with other cultivars, suggesting a major tendency to accumulate Mn in roots (See model; Fig. 5.1). Similarly with our results, Mora et al. (2009) observed that tolerant ryegrass accumulated higher Mn concentrations in roots than in shoots. These results suggest an internal mechanism of control on Mn concentration and a lower Mn translocation to shoots in Mn-tolerant than Mn-sensitive ryegrass cultivar, which had a great translocation to shoot. In other Mn-tolerant plant species, also is relatively low the Mn translocation to aerial parts (Li et al., 2010). Thus, it can be considered that Legacy is a Mn-tolerant genotype with a low translocation of Mn to shoots, whereas Bluegold is Mn-sensitive genotype, with a major Mn translocation (See Model; Fig. 5.1).

The fluorescence parameters of the PSII (Fv/Fm,  $\Phi$ PSII) showed a decrease in leaves of blueberry cultivars with the increasing Mn doses mainly in Bluegold and Brigitta (Table 1). Researches made with other metal, suggest a photoinhibition by loss the PSII efficiency, when a decrease in Fv/Fm was observed in plants subjected to cadmium (Cd) (Pagliano et

---

al., 2006). Excess Mn decreased significantly  $\Phi$ PSII in cucumber seedlings, suggesting a decline of carbon metabolism capacity and/or a low consumption of ATP and NADPH in Calvin cycle (Feng et al., 2009; Kitao et al., 1997). Doncheva et al. (2009) found a decrease in Fv/Fm and an increase in NPQ values in the Mn-sensitive cultivar of maize, suggesting a photoinhibition by Mn excess. On the other hand, net photosynthesis and stomatal conductance of blueberry cultivars were negatively affected by increasing Mn doses in Bluegold, as was demonstrate in Mn-sensitive maize cultivars (Stoyanova et al., 2009) and cucumber plants (Feng et al., 2009). Nonetheless, early experiments in tobacco young leaves grown under Mn excess showed a decrease in net photosynthesis (Nable et al., 1988), but not associated with decreased leaf CO<sub>2</sub> conductance. These authors concluded that photosynthesis inhibition is an early indicator of Mn accumulation in tobacco leaves (Nable et al., 1988). With those previous researches, it could be considered to Bluegold as Mn-sensitive genotype, suggesting a major translocation to aerial part (See model, Fig. 5.1).

Abiotic stress, including Mn excess, can produce oxidative damage on plant cells leading to oxidative stress (González et al., 1998; Demirevska-Kepova et al., 2004). In our results, Legacy showed a major lipid peroxidation compared with the other cultivars; however, all the lipid peroxidation levels remained constant from the beginning until the end of Mn experiment. Different results have been reported in cucumber and wetland plants subjected to Mn excess where Mn doses increased lipid peroxidation in leaves (Shi et al., 2005; Shi and Zu, 2008; Najeeb et al., 2009). In ryegrass studies with Mn excess, lipid peroxidation was lower in Mn-tolerant ryegrass cultivars than the Mn-sensitive ones, indicating that when Mn-sensitive cutivars incremented shoot Mn concentration, a proportional augment in lipid peroxidation were observed (Mora et al., 2009). However, to prevent and reduce damage effects induced by ROS, plant has developed antioxidant defense mechanisms, which include enzymatic and non-enzymatic systems (Foyer and Noctor, 2003; Apel and Hirt, 2004). In the present study, radical scavenging activity (RSA) in blueberry cultivars with Mn excess had differences in leaves and roots, being higher in leaves than in roots in all cultivars. Interestingly, Legacy had an increase of RSA, even though TBARS amount remained unchanged, while in Bluegold the RSA was proportional to its TBARS values, reaching maximum at the highest Mn treatment (See model; Fig. 5.1). It is noteworthy that

---

the same cultivars used in this work and subjected to Al toxicity showed that Bluegold is the Al-sensitive cultivar, with the greatest values of TBARS without changes in RSA (Reyes-Diaz et al., 2010). The authors suggest that Legacy is Al-tolerant cultivar due to the possibility to activate defense mechanism, which contribute to decrease lipid peroxidation. Specifically, a decrease in enzyme activities as SOD was observed in Bluegold subjected to Mn excess.

Organic acid exudation in plants is plant-species specific and depends on the magnitude of stress (Jones, 1998). It has different functions as the uptake of some nutrients as P (Hoffland, 1992; Johnson et al. 1996) and the metal detoxification, where the responses depend highly of the stress- and plant-species (Jones, 1998). Organic acids associated to Mn detoxification are oxalate, citrate, malate and succinate (Horst et al. 1999). In our study, four organic acid exudates were analyzed. From these, oxalate presented a mayor amount of exudates in all cultivars. Mora et al. (2009) found that Mn-tolerant ryegrass had the highest oxalate exudation rate at increasing Mn excess. Those responses suggest that oxalate can present an internal Mn-tolerance mechanism. High citrate exudation was found also in Mn-tolerant ryegrass, associated to the high affinity of citrate for  $Mn^{2+}$  in the rizosphere. In general, researches indicate that these organic anions exudation are released under heavy metals toxicity (lead and Cu), protecting the roots by chelating cations forming nontoxic complexes and enhanced the tolerance (Ryan et al. 2001).


For a better understanding of the results obtained in this chapter, we have designed a model that summarizes the main mechanisms involved in Mn toxicity in highbush blueberry (Fig. 5.1). In this model it is observed that Mn available is more abundant in roots of the Mn-tolerant cultivar than the sensitive ones, appearing a lower translocation from roots to leaves in Mn-tolerant cultivar. Probably, the excess of  $Mn^{2+}$  in roots, are inactivated by organic acid anions via chelation by mainly oxalate in the Mn-tolerant cultivar. In Mn-sensitive cultivar, exudation of organic acid anions did not change. With respect to SOD activity in leaves, it was not affected in Mn tolerant cultivar; whereas, in the Mn-sensitive cultivar there is a tendency to strongly decrease of the SOD activity when compared with the control at the highest Mn treatment.

---

In Chapter 4, we analyze how a Mn excess inhibits differentially the photosystems in *Arabidopsis thaliana*. Interestingly, while reduced CO<sub>2</sub> assimilation induced by Mn excess has been reported in many species, it is considered one of the major physiological effects of Mn toxicity on plants (Nable et al., 1988; González and Lynch, 1997; Kitao et al., 1997a; Lidon et al., 2004; Macfie and Taylor, 1992; Li et al., 2010). The functional integrity of the photosynthetic apparatus assessed by the maximal efficiency of PSII photochemistry ( $F_v/F_m$ ) did not decline as a result of exposure to excess Mn in several plant species (Nable et al., 1988; Kitao et al., 1997b; Subrahmanyam and Rathore, 2000; Hajiboland and Hasani, 2007; Doncheva et al., 2009). However, some studies have reported a substantial decrease in  $F_v/F_m$  as a result of excess of Mn treatment in *Citrus* species (Papadakis et al., 2007; Li et al., 2010), rice (Lidon et al., 2004), Mn-sensitive maize (Doncheva et al., 2009) and cucumber (Feng et al., 2009). These results clearly indicate that the excess Mn- induced decline in CO<sub>2</sub> assimilation may or may not be accompanied by changes in PSII photochemistry and that response is species dependent. The experimental data presented in our study also failed to demonstrate any substantial effect of excess Mn within a wide range of Mn concentrations on the maximum efficiency of PSII photochemistry in *Arabidopsis* plants. Assessing the relative abundance of PSII-associated proteins also showed no changes in the immuno-detectable amounts of PSII-reaction center protein D1, the light harvesting chlorophyll-protein complex of PSII (Lhcb1) and the manganese stabilizing 33kDa-protein of the water splitting complex of PSII (PsbO) polypeptides in plants exposed to high Mn concentrations compared to control. In addition to the lack of significant inhibitory effects of excess Mn on PSII photochemistry, an earlier study reported that the photochemistry of photosystem I (PSI) and the photosynthetic electron transport were not significantly affected during early development of Mn toxicity in tobacco plants (Nable et al., 1988). However, a decreased Hill activity in isolated chloroplasts was found in mungbean leaves exposed to toxic Mn concentrations (Sinha et al., 2002). More recently, Li et al. (2010) have suggested that Mn excess can effectively impair the whole photosynthetic electron transport chain, thus restricting the production of reducing equivalents and limiting the rate of CO<sub>2</sub> assimilation in *Citrus grandis* seedlings.

---

Despite these few reports, the potential effect(s) of excess Mn on the functional/structural integrity of PSI remains elusive. To our knowledge, the results presented in our study are the first report of an *in vivo* assessment of high Mn concentrations on PSI photochemistry. In contrast to PSII photochemistry, *in vivo* measurements of the oxidation state of P700 (P700<sup>+</sup>) (Klughammer and Schreiber, 1991; Ivanov et al., 1998), the primary donor of PSI demonstrated that the relative amount of oxidizable P700 (P700<sup>+</sup>) decreased by 30% in Mn-treated *Arabidopsis* plants at concentrations above 1000  $\mu$ M. The functional impairment of PSI photochemistry by excess Mn was accompanied by a significant reduction in the abundance of PSI reaction center polypeptides (PsaA and PsaB). This clearly indicates that the major target of Mn toxicity within the photosynthetic electron transport chain of *Arabidopsis* is PSI rather than PSII related components. The reduced amounts of PSI reaction center polypeptides PsaA and PsaB would imply acceptor side limitations of the photosynthetic electron transport and this could explain the increased reduction state Q<sub>A</sub> in Mn-stressed plants reported earlier (Kitao et al., 1997b).



**Figure 5.1. Theoretical model showing effect of Mn excess on physiological and biochemical features in leaves and roots of Mn- tolerant and Mn-sensitive blueberry cultivars.** The left side of the model represents the behaviour in leaves and roots of the Mn-sensitive blueberry cultivar, whereas in the right side the changes of Mn-tolerant blueberry cultivar in leaves and roots are shown. Note that the size of leaves of the tolerant cultivar is greater than the sensitive ones. Black arrows represent the possible direction of transport Mn ions, (grey circles) into the plants. White arrows showed an enhancement or decrease of specific parameters. Blue circles indicate organic acid anions exudates in roots (*oxal*: oxalate; *citr*: citrate). Other abbreviations of parameters are: *Fv/Fm*: maximal photochemical efficiency of PSII;  $\Phi$ PSII: effective photochemical efficiency of PSII; *Pn*: net photosynthesis; *g<sub>s</sub>*: stomatal conductance; SOD: superoxide dismutase enzyme activity;

---

RSA: radical scavenging activity; LP: lipid peroxidation. In the Mn tolerant cultivar, parameters which did not change or with a little variation were not represented in the model. It is important to note for a better comprehensive that the present model was performed having into account the comparison between the control and Mn treated plants with the higher or the highest Mn treatment(s).



Chapter 6:

General conclusions

---

## 6.1 General conclusions

According to main results in this thesis, it is concluded that:

- Mn excess affects differently the physiological and biochemical features of blueberry cultivars. Legacy was the most Mn-tolerant cultivar compared with Brigitta and Bluegold, being the last cultivar the most Mn-sensitive. This assumption is based on the major Mn accumulation in Legacy roots, maintaining its growth and photosynthesis. Likewise, this cultivar showed an enhancement of antioxidant capacity, maintaining lipid peroxidation at increasing Mn treatments. In addition, oxalate and citrate were the most important organic acid anions exudated, incrementing its concentration gradually with the augment of Mn doses in Legacy.
- Additionally, results of Mn excess on photosynthetic performance in *A. thaliana* indicated that the exposure of *Arabidopsis* plants to excess Mn causes specific negative effects on the abundance of polypeptides comprising the reaction center of PSI, thus resulting in decreased PSI photochemistry and lower capacity for PSI-dependent cyclic electron transport, which may be due to a Mn-induced Fe-deficiency and may have critical physiological implications under conditions of Mn toxicity in higher plants.

## 6.2 Futures perspectives

The knowledge presented here will contribute to understand the mechanism of Mn resistant in plants, which is very important because this element is an essential micronutrient and on the other hand, it is toxic when, is in excess depending on the acidity condition of the soil. Likewise, the understanding of this mechanism will also contribute to improving the yield and quality of cultivated plants in acid soils. Future efforts for developing Mn-tolerant plants should take all the physiological and biochemical processes into account, helping the farmers to improve the nutrition quality of plants.

---

### 6.3 General references

- Aguilera SM, Borie G, Peirano P, Galindo G.** 1997. Organic matter in volcanic soils in Chile. Chemical and biochemical characterization. *Communication in Soil Science and Plant Analysis* 28: 899-912.
- Aguilera M, Mora M, Borie G, Peirano P, Zunino H.** 2002. Balance and distribution of sulphur in volcanic ash-derived soils in Chile. *Soil Biology and Biochemistry* 34: 1355-1361.
- Alam S, Kodama R, Akiha F, Kamei S, Kawai S.** 2006. Alleviation of manganese phytotoxicity in barley with calcium. *Journal of Plant Nutrition*. 29: 59-74.
- Apel K, Hirt H.** 2004. Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annual Review Plant Biology* 55: 373-379.
- Blamey F, Joyce D, Edwards D, Asher C.** 1986. Role of trichomes in sunflower tolerance to manganese toxicity. *Plant Soil*. 91: 171-180.
- Blokhina O, Virolainen E, Fagerstedt KV.** 2003. Antioxidants, oxidative damage and oxygen deprivation stress: a Review. *Annals of Botany* 91: 179-194.
- Boojar M, Goodarzi F.** 2008. Comparative evaluation of oxidative stress status and manganese availability in plants growing on manganese mine. *Ecotoxicology and Environmental Safety* 71: 692–699.
- Bowler C, VanCamp W, VanMontagu M, Inze D.** 1994. Superoxide dismutase in plants. *Critical Reviews in Plant Sciences* 13: 199-218.
- Burnell JN.** 1988. The biochemistry of manganese in plants. In: Graham RD, Hannam J, Uren NC, eds. *Manganese in Soils and Plants*. Dordrecht, The Netherlands: Kluwer Academic Publishers, 125-133.
- Dakora FD, Phillips DA.** 2002. Root exudates as mediators of mineral acquisition in low-nutrient environments. *Plant Soil* 245:35–47.

- 
- Delhaize E, Kataoka T, Hebb DM, White RG, Ryan PR.** 2003. Genes encoding proteins of the cation diffusion facilitator family that confer manganese tolerance. *Plant Cell* 15: 1131–1142.
- Delhaize E, Gruber BD, Ryan PR.** 2007a. The roles of organic anion permeases in aluminium resistance and mineral nutrition. *FEBS Letters* 581: 2255–2262.
- Delhaize E, Gruber BD, Pittman JK, White RG, Leung H, Miao Y, Jiang L, Ryan PR, Richardson AE.** 2007b. A role for the AtMTP11 gene of *Arabidopsis* in manganese transport and tolerance. *The Plant Journal* 51: 198–210.
- Demirevska-Kepova K, Simova-Stoilova L, Stoyanova Z, Holzer R, Feller U.** 2004. Biochemical changes in barley plants after excessive supply of copper and manganese. *Environmental Experimental Botanical* 52: 253-266.
- Doncheva S, Poschenrieder C, Stoyanova Z, Georgieva K, Velichkova M, Barceló J.** 2009. Silicon amelioration of manganese toxicity in Mn-sensitive and Mn-tolerant maize varieties. *Environmental and Experimental Botany* 65: 189–197.
- Ducic T, Polle A.** 2005. Transport and detoxification of manganese and copper in plants. *Brazilian Journal of Plant Physiology* 17: 103-112.
- Enami I, Okumura A, Nagao R, Suzuki T, Iwai M, Shen JR.** 2008. Structures and functions of the extrinsic proteins of photosystem II from different species. *Photosynthesis Research* 98: 349-363.
- Farasova A, Beinrohr E.** 1998. Metal-metal interaction in accumulation of V, Ni, Mo, Mn, and Cu in under and above ground parts of *Sinapis alba*. *Chemosphere* 36: 1305-1317.
- Fecht-Christoffers M, Maier P, Horst W.** 2003. Apoplastic peroxidases and ascorbate are involved in manganese toxicity and tolerance of *Vigna unguiculata*. *Physiologia Plantarum* 117: 237-244.

- 
- Feng JP, Shi QH, Wang XF.** 2009. Effects of exogenous silicon on photosynthetic capacity and antioxidant enzyme activities in chloroplast of cucumber seedlings under excess manganese. *Agricultural Science in China* 8: 40–50.
- Ferreira KN, Iverson TM, Maghlaoui K, Barber J, Iwata S.** 2004. Architecture of the photosynthetic oxygen-evolving center. *Science* 303: 1831-1838.
- Foy C.** 1984. Physiological effects of hydrogen, aluminum, and manganese toxicities in acid soils. In *Soil Acidity and Liming*, ed. F Adams, pp. 57–97. Madison, WI: Agron. Soc. Am.
- Foy C.** 1978. The physiology of metal toxicity in plants. *Annual Review Plant Physiology* 29: 511-566.
- Foyer C, Noctor G.** 2003. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiologia Plantarum* 119: 355-364.
- Gherardi M, Rengel Z.** 2004. The effect of manganese supply on exudation of carboxylates by roots of lucerne (*Medicago sativa*). *Plant and Soil* 260: 271-282.
- González A, Steffen KL, Lynch JP.** 1998. Light and excess manganese. Implication for oxidative stress in common bean. *Plant Physiology* 118: 493-504.
- González A, Lynch JP.** 1997. Effects of manganese toxicity on leaf CO<sub>2</sub> assimilation of contrasting common bean genotypes. *Physiologia Plantarum* 101:872-880.
- Goussias C, Boussac A, Rutherford W.** 2002. Photosystem II and photosynthetic oxidation of water: an overview. *Phil. Trans. R. Soc. Lond. B* 357: 1369-1381.
- Guo T, Zhang G, Zhang Y.** 2007. Physiological changes in barley plants under combined toxicity of aluminum, copper and cadmium. *Colloids and Surfaces B: Biointerfaces* 57: 182-188.
- Hajiboland R, Hasani BD.** 2007. Effect of Cu and Mn toxicity on chlorophyll fluorescence and gas exchange in rice and sunflower under different light intensities. *Journal of Stress Physiology and Biochemistry* 3: 4–17.

- 
- Hankamer B, Barber J, Boekema EJ.** 1997. Structure and membrane organization of photosystem II in green plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 48: 641-671.
- Hegedüs A, Erdei S, Horvath G.** 2001. Comparative studies of H<sub>2</sub>O<sub>2</sub> detoxifying enzymes in green and greening barley seedlings under cadmium stress. *Plant Science* 160: 1085-1093.
- Hirschi K., Korenkov V, Wilganowski N, Wagner GJ.** 2000. Expression of *Arabidopsis* CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. *Plant Physiology* 124: 125-133.
- Hoffland E, Van den Boogaard R, Nelemans JA, Findenegg GR.** 1992. Biosynthesis and root exudation of citric and malic acids in phosphate-starved rape plants. *New Phytologist* 122: 675-680.
- Horiguchi T.** 1988. Mechanism of manganese toxicity and tolerance of plants. VII. Effect of light intensity on manganese-induced chlorosis. *Journal of Plant Nutrition* 11: 235-245.
- Horst WJ, Fecht M, Naumann A, Wissemeier AH, Maier P.** 1999. Physiology of manganese toxicity and tolerance in *Vigna unguiculata* (L.) Walp. *Journal of Plant Nutrition and Soil Science* 162: 263-274.
- Horst WJ.** 1988. The physiology of manganese toxicity. In *Manganese in Soils and Plants*, R.D. Graham, J. Hannam, and N.C. Uren, eds. (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp.175-188.
- Howard LR, Clark JR, Brownmiller C.** 2003. Antioxidant capacity and phenolic content in blueberries as affected by genotype and growing season. *Journal of the Science of Food and Agriculture* 83:1238-1247.
- Humphries J, Stangoulis J, Graham R.** 2007. Manganese. In: A. Barker, D. Pilbeam (eds). *Handbook of Plant Nutrition*, Taylor and Francis, USA, pp. 351-366.

- 
- Hue N.** 1988. A possible mechanism for manganese toxicity in Hawaii soils amended with a low-Mn sewage sludge. *Journal Environmental Quality* 14: 473-479.
- Ireland G, Wilk P.** 2006. Blueberry production in northern NSW. *Primefact* 195. 8p.
- Ivanov AG, Morgan R, Gray GR, Velitchkova MY, Hüner NPA.** 1998. Temperature/light dependent development of selective resistance to photoinhibition of photosystem I. *FEBS Letters* 430: 288–292.
- Jarvis S.** 1987. Laboratory studies of effects of lime on soluble Al, Ca, Mg and Mn in a pelostagnogley soil. *Journal of Soil Science* 38: 443-451.
- Johnson JF, Vance CP, Allan DL.** 1996. Phosphorus deficiency in *Lupinus albus*: altered lateral root development and enhanced expression of phosphoenolpyruvate carboxylase. *Plant Physiology* 112: 31–41.
- Jones DL.** 1998. Organic acids in the rhizosphere – a critical review. *Plant and Soil* 205: 25–44.
- Kern J, Renger G.** 2007. Photosystem II: Structure and mechanism of the water:plastoquinone oxidoreductase. *Photosynthesis Research* 94: 183-202.
- Kitao M, Lei TT, Koike T.** 1997a. Comparison of photosynthetic responses to manganese toxicity of deciduous broad leaved trees in northern Japan. *Environmental Pollution* 97: 113–118.
- Kitao M, Lei TT, Koike T.** 1997b. Effects of manganese toxicity on photosynthesis of white birch (*Betula platyphylla* var. *japonica*) seedlings. *Physiologia Plantarum* 101: 249–256.
- Klughammer C, Schreiber U.** 1991. Analysis of light-induced absorbancy changes in the near-infrared spectral region. 1. Characterization of various components in isolated chloroplasts. *Zeitschrift für Naturforschung C* 46: 233–244.

- 
- Kochian L, Hoekenga O, Piñeros M.** 2004. How do crop plants tolerate acid soils? Mechanisms of aluminum tolerance and phosphorous efficiency. *Annual Review Plant Biology* 55: 459-493.
- Lei Y, Korpelainen H, Li C.** 2007. Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. *Chemosphere* 68: 686-694.
- Li Q, Chen L-S, Jiang H-X, Tang N, Yang L-T, Lin Z-H, Li Y, Yang G-H.** 2010. Effects of manganese-excess on CO<sub>2</sub> assimilation ribulose-1,5-bisphosphate carboxylase/oxygenase, carbohydrates and photosynthetic electron transport of leaves, and antioxidant systems of leaves and roots in *Citrus grandis* seedlings. *BMC Plant Biology* 10: 42.
- Lidon F, Barreiro MG, Ramalho JC.** 2004. Manganese accumulation in rice: implications for photosynthetic functioning. *Journal of Plant Physiology* 161: 1235-1244.
- Lidon F, M Teixeira.** 2000. Rice tolerance to excess Mn: Implications in the chloroplast lamellae and synthesis of a novel Mn protein. *Plant Physiology Biochemistry* 38: 969-978.
- Macfie S, Taylor G.** 1992. The effects of excess manganese on photosynthetic rate and concentration of chlorophyll in *Triticum aestivum* grown in solution culture. *Physiologia Plantarum* 85: 467-475.
- Marschner H.** 1995. *Mineral Nutrition of Higher Plants*. 2<sup>nd</sup> Edition. Academic Press. San Diego, USA. 889 p.
- Marschner H.** 1991. Mechanisms of adaptation of plants to acid soils. In: R. Wright, V. Baligar, R. Murrmann (Eds). *Development in Plant and Soil Sciences*. Plant-soil interactions at low pH. Kluwer Academic Publisher, West Virginia, pp. 1-20.

- 
- Mora M, Rosas A, Ribera A, Rengel R.** 2009. Differential tolerance to Mn toxicity in perennial ryegrass genotypes: involvement of antioxidative enzymes and root exudation of carboxylates. *Plant and Soil* 320: 79–89.
- Mora M.** 1993. Nivel de fertilidad de los suelos de la IX Región y su relación con la acidificación. *Frontera Agrícola* 1: 5-12.
- Mora M, García J, Santander J, Demanet R.** 1999. Rol de los fertilizantes nitrogenados y fosfatados en los procesos de acidificación de los suelos. *Frontera Agrícola* 5: 59-81.
- Mora M., Alfaro M, Jarvis S, Demanet R, Cartes P.** 2006. Soil aluminium availability in Andisols of southern Chile and its effect on forage production and animal metabolism. *Soil Use and Management* 22: 95-101.
- Mukhopadhyay MJ, Sharma A.** 1991. Manganese in cell metabolism of higher plants. *Botanical Review* 57: 117–149.
- Nable RO, Houtz RL, Cheniae GM.** 1988. Early inhibition of photosynthesis during development of Mn toxicity in tobacco. *Plant Physiology* 86: 1136-1142.
- Najeeb U, Xu L, Ali S, Jilani G, Gonga HJ, Shenc WQ, Zhou WJ.** 2009. Citric acid enhances the phytoextraction of manganese and plant growth by alleviating the ultrastructural damages in *Juncus effusus* L. *Journal of Hazardous Materials* 170:1156–1163.
- ODEPA,** 2012. Boletín Frutícola. Ministerio de Agricultura. Oficina de Estudios y Políticas Agrarias, Gobierno de Chile. 32 p.
- ODEPA, CIREN.** 2012. Catastro frutícola. Principales resultados Región de La Araucanía. Oficina de Estudios y Políticas Agrarias y Centro de Información de Recursos Naturales. Santiago Chile, 42 p.
- Pagliano C, M Raviolo, FD Vecchia, R Gabbrielli, C Gonnelli, N Rascio, R Barbato, N La Rocca.** 2006. Evidence for PSII donor-side damage and photoinhibition induced by cadmium treatment on rice (*Oryza sativa* L.). *Journal of Photochemistry and Photobiology B: Biology* 84: 70–78.

- 
- Papadakis IE, Giannakoula A, Therios IN, Bosabalidis AM, Moustakas M, Nastou A.** 2007. Mn-induced changes in leaf structure and chloroplast ultrastructure of *Citrus volkameriana* (L) plants. *Journal of Plant Physiology* 164: 100–103.
- Peñaloza E, Martínez J, Montenegro A, Corcuera L.** 2004. Respuesta de dos especies de lupino a aluminio fitotóxico. *Agricultura Técnica* 64: 127-138.
- Pittman J.** 2005. Managing the manganese: molecular mechanisms of manganese transport and homeostasis. *New Phytologist* 167: 733-742.
- Porter GS, Bajita-Locke JB, Hue NV, Strand D.** 2004. Manganese solubility and phytotoxicity affected by soil moisture, oxygen levels, and green manure additions. *Communications in Soil Science and Plant Analysis* 35: 99–116.
- Prior RL, Cao G, Martin A, Sofic E, McEwen J, O'Brien C, Lischner N, Ehlenfeldt M, Kalt W, Krewer G, Mainland CM.** 1998. Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of *Vaccinium* species. *Journal of Agricultural and Food Chemistry* 46: 2686-2693.
- Reyes-Díaz M, Inostroza-Blancheteau C, Millaleo R, Cruces E, Wulff-Zottele C, Alberdi M, Mora ML.** 2010. Long-term aluminum exposure effects on physiological and biochemical features of highbush blueberry cultivars. *Journal of American Society Horticultural Science* 3: 212-222.
- Rosas A, Rengel Z, Mora M.** 2007. Manganese supply and pH influence growth, carboxylate exudation and peroxidase activity of ryegrass and white clover. *Journal of Plant Nutrition* 30: 253-270.
- Rosas A, Rengel Z, Ribera A, Mora ML.** 2011. Phosphorus nutrition alleviates manganese toxicity in *Lolium perenne* and *Trifolium repens*. *Journal of Plant Nutrition and Soil Science* 174: 210–219.
- Ryan PR, Raman H, Gupta S, Horst WJ, Delhaize E.** 2009. A second mechanism for aluminum resistance in wheat relies on the constitutive efflux of citrate from roots. *Plant Physiology* 149: 340–351.

- 
- Ryan PR, Delhaize E, Jones DL.** 2001. Function and mechanism of organic anion exudation from plant roots. *Annual Review of Plant Physiology and Plant Molecular Biology* 52: 527–60.
- Rutherford AW, Boussac A.** 2004. Water photolysis in biology. *Science* 303: 1782-1784.
- Sasaki A, Yamaji N, Xia J, Ma JF.** 2011. OsYSL6 is involved in the detoxification of excess Manganese in rice. *Plant Physiology* 157: 1832-1840.
- Shi Q, Bao Z, Zhu Z, He Y, Qian Q, Yu J.** 2005. Silicon-mediated alleviation of Mn toxicity in *Cucumis sativus* in relation to activities of superoxide dismutase and ascorbate peroxidase. *Phytochemistry* 66: 1551-1559.
- Shi Q, Zhu Z, Xu M, Qian Q, Yu J.** 2006. Effect of excess manganese on the antioxidant system in *Cucumis sativus* L. under two light intensities. *Environmental and Experimental Botany* 58: 197-205.
- Shi Q, Zhu Z.** 2008 Effects of exogenous salicylic acid on manganese toxicity, element contents and antioxidative system in cucumber. *Environmental and Experimental Botany* 63: 317–326.
- Sinha S, Mukherji S, Dutta J.** 2002. Effect of manganese toxicity on pigment content, Hill activity and photosynthetic rate of *Vigna radiata* L. Wilczek seedlings. *Journal of Environmental Biology* 23: 253-257.
- Schützendübel A, Polle A.** 2002. Plant responses to abiotic stresses: heavy metal-induced oxidative stress and protection by mycorrhization. *Journal of Experimental Botany* 53: 1351-1365.
- Sparrow L, Uren N.** 1987. Oxidation and reduction of Mn in acidic soils: effect of temperature and soil pH. *Soil Biology and Biochemistry* 19: 143-148.
- Sposito G.** 1989. *The chemistry of soils*. New York. 277 p.

- 
- Srivastava S, Dubey RS.** 2011. Manganese-excess induces oxidative stress, lowers the pool of antioxidants and elevates activities of key antioxidative enzymes in rice seedlings. *Plant Growth Regulation* 64:1–16.
- Stoyanova Z, Poschenrieder C, Tzvetkova N, Doncheva S.** 2009. Characterization of the tolerance to excess manganese in four maize varieties. *Soil Science and Plant Nutrition* 55:747–753.
- Subrahmanyam D, Rathore VS.** 2000. Influence of manganese toxicity on photosynthesis in ricebean (*Vigna umbellata*) seedlings. *Photosynthetica* 38: 449–453.
- Taylor GJ.** 1991. Current views of the aluminum stress response: the physiological basis of tolerance. *Current Topics in Plant Biochemistry and Physiology* 10:57-93.
- Watmough S, Eimers M, Dillon P.** 2007. Manganese cycling in central Ontario forests: Response to soil acidification. *Applied Geochemistry* 22: 1241-1247.
- Whitehead D.** 2000. Nutrient elements in grassland. *Soil-Plant-Animal Relationships*. CABI publishing, UK. P: 220-252.
- Wissemeyer A, Horst W.** 1992. Effect of light intensity on manganese toxicity symptoms and callose formation in cowpea (*Vigna unguiculata* (L.) Walp.). *Plant Soil* 143: 299-309.



**APPENDIX**

## **MANGANESE AS ESSENTIAL AND TOXIC ELEMENT FOR PLANTS: TRANSPORT, ACCUMULATION AND RESISTANCE MECHANISMS**

**R. Millaleo<sup>1</sup>, M. Reyes-Díaz<sup>2</sup>, A.G. Ivanov<sup>3</sup>, M.L. Mora<sup>4</sup>, and M. Alberdi<sup>4\*</sup>**

<sup>1</sup>Programa de Doctorado en Ciencias de Recursos Naturales, Universidad de La Frontera, Casilla 54-D, Temuco, Chile. <sup>2</sup>Center of Plant-Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Casilla 54-D, Temuco, Chile. <sup>3</sup>Department of Biology and the Biotron, University of Western Ontario, London, Ontario, N6A 5B7 Canada. <sup>4</sup>Departamento de Ciencias Químicas, Center of Plant-Soil Interaction and Natural Resources Biotechnology - Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Casilla 54-D, Temuco, Chile.

\*Corresponding author: [malberdi@ufro.cl](mailto:malberdi@ufro.cl)

### **ABSTRACT**

Manganese is an essential element for plants, intervening in several metabolic processes, mainly in photosynthesis and as an enzyme antioxidant-cofactor. Nevertheless, an excess of this micronutrient is toxic for plants. Mn phytotoxicity is manifested in a reduction of biomass and photosynthesis, and biochemical disorders such as oxidative stress. Some studies on Mn toxicity and Mn translocation from soil to plant cells in Mn<sup>2+</sup> form have demonstrated their importance under low pH and redox potential conditions in the soil. When Mn is inside the cells, mechanisms that can tolerate this toxicity are also observed, being important the compartmentalization of this metal in different organelles of shoot and leaf plant cells. A key role of antioxidative systems in plants in relation to high Mn amounts has also been reported as a defense mechanism. The purpose of this review is to show the role of Mn as an essential micronutrient and as a toxic element to higher plants as well as to their transport and tolerance mechanisms. The forms and dynamics of this element in soils and the importance of the acidity for this dynamic and availability for plants are also given.

**Keywords:** Manganese, Mn toxicity, resistance mechanisms

### **INTRODUCTION**

Manganese (Mn) is an essential micronutrient in most organisms. In plants, it participates in the structure of photosynthetic proteins and enzymes. Its deficit is dangerous for chloroplasts because it affects the water-splitting system of photosystem II (PSII), which provides the necessary electrons for photosynthesis (Buchanan, 2000). However, its excess seems also to be particularly damaging to the

photosynthetic apparatus (Mukhopadhyay and Sharma, 1991). Thus, Mn has two roles in the plant metabolic processes: as an essential micronutrient and as a toxic element when it is in excess (Kochian *et al.*, 2004; Ducic and Polle, 2005). Mn toxicity is favored in acid soils (Pendias and Pendias, 1992). With decreasing pH, the amount of exchangeable manganese – mainly Mn<sup>2+</sup> form – increases in the soil solution. This Mn form is available for

plants and can be readily transported into the root cells and translocated to the shoots, where it is finally accumulated (Marschner, 1995). In contrast, other forms of Mn predominate at higher pH values, such as Mn (III) and Mn (IV), which are not available and cannot be accumulated in plants (Rengel, 2000).

Excessive Mn concentrations in plant tissues can alter various processes, such as enzyme activity, absorption, translocation and utilization of other mineral elements (Ca, Mg, Fe and P), causing oxidative stress (Ducic and Polle, 2005; Lei *et al.*, 2007). The threshold of Mn injury as well as the tolerance to an excess of this metal is highly dependent on the plant species and cultivars or genotypes within a species (Foy *et al.*, 1988, Horst, 1988).

The purpose of this review is to illustrate the most current understanding about Mn role as an essential micronutrient and as a toxic element to higher plants, the long distance and cellular transport in plants as well as the mechanisms or strategies involved for to resist an overload of this metal. The forms and dynamics of this element in soils and the importance of the acidity for this dynamic and availability to plants are also given.

#### **MANGANESE FORMS AND DYNAMICS IN SOILS**

Manganese biogeochemistry in soils is complex, because it is present in several oxidation states (0, II, III, IV, VI and VII), while in biological systems it occurs preferably as II, III and IV. Divalent manganese (Mn II) is the most soluble species of Mn in soil, whereas the solubility of Mn III and Mn IV are very low (Guest *et al.*, 2002). Mn oxides can form co-precipitates with iron (Fe) oxides, exhibiting amphoteric behavior. In

addition, Mn interacts both with cations and anions in oxidation-reduction reactions involving Mn. These reactions are influenced by a variety of physical, chemical and microbiological processes (Bradl, 2004).

Both pH and redox conditions influence Mn bioavailability in soils (Marschner, 1995; Porter *et al.*, 2004). In most acid soils at low pH (<5.5) and an increased redox potential of Mn, oxides can be easily reduced in the soil exchange sites (Kogelmann and Sharpe, 2006), increasing the concentration of soluble  $Mn^{2+}$  (Watmough *et al.*, 2007), which is the predominant Mn form in the soil solution (Adriano, 2001) and the most available Mn form for plants (Marschner, 1995). At higher soil pH (up to pH 8), chemical  $Mn^{2+}$  auto-oxidation is favored over  $MnO_2$ ,  $Mn_2O_3$ ,  $Mn_3O_4$  and even  $Mn_2O_7$ , which are not normally available to plants (Ducic and Polle, 2005; Humpries *et al.*, 2007; Gherardi and Rengel, 2004). Furthermore, high pH allows Mn adsorption into soil particles, decreasing their availability (Fageria *et al.*, 2002). Nevertheless, some reports have suggested that an excess of available Mn is produced under reduced soil conditions, even at high soil pH values (Hue, 1988). A reducing environment can be produced when there is an excess of water, poor drainage or applications of organic material (Hue, 1988; El-Jaoual and Cox, 1998). Different organic molecules can dissolve solid Mn oxides through transfer of electrons, transforming them into an available Mn form for plants (Laha and Luthy, 1990). Soil acidification is also accentuated by abundant pluviometry during winter, causing the main cations to leak from the soil (Mora *et al.*, 2006). On the other hand, lime application is a key factor in decreasing soluble Mn in acid soils with a high Mn content, given that it can increase soil pH (Hue and Mai, 2002).

The total Mn content in soils is variable. Sparks (1995) reported small amounts of Mn in soils, fluctuating from 20 to 10,000 mg kg<sup>-1</sup> soil, whereas other authors have registered total Mn contents between 450 and ~ 4,000 mg Mn kg<sup>-1</sup> soil (Adriano, 2001). In addition, the total Mn soil content was from 15 to 17 mg kg<sup>-1</sup> in acid soils without liming (pH about 4.4) (Hue and Mai, 2002). In liming acid soils, the interchangeable Mn concentration varied from 14 to 96 mg kg<sup>-1</sup> soil in one year, with higher concentrations under high moisture and temperature conditions (Conyers *et al.*, 1997). In Chilean volcanic soils, so-called Andisols, Mn concentrations fluctuate between 4.5 and 80 mg kg<sup>-1</sup> depending on the agronomic management. Moreover, the Mn amount is higher in pasture soils (up to 400 mg kg<sup>-1</sup>) mainly in winter (Data from Laboratorio de Análisis de Suelo y Planta, Universidad de La Frontera, Temuco, Chile).

Environmental conditions also affect Mn soil contents. The highest concentrations of soluble and exchangeable Mn are found after hot, dry summers and under warm waterlogged conditions in acid soils. This is probably due to the inhibition of Mn-oxidizing organisms, thereby allowing the chemical reduction of Mn oxides in these soils (Sparrow and Uren, 1987; Conyers *et al.*, 1997).


#### **Manganese dynamics in the rhizosphere**

Rhizosphere, which is the narrow zone of soil immediately surrounding the root system, is of great importance for mineral plant nutrition. In this zone, both the mobilization and immobilization of nutrients occur (Marschner, 1995). As shown in Figure 1, a mobilization of Mn<sup>2+</sup> is produced by the rhizosphere acidification due to the release of H<sup>+</sup> or low molecular weight organic acids

(LMWOA) from plants (Rengel and Marschner, 2005). Organic acids released in anion forms from roots can chelate Mn<sup>2+</sup> released from the MnO<sub>x</sub> (Mn oxides) (Ryan *et al.*, 2001). Neumann and Römheld (2001) reported that mobilization of micronutrients (including Mn) into the rhizosphere is due mainly by its acidification and complexation with the organic acids (citrate) in various plant species. It has been reported that organic amendments (chip compost and pine bark) applied to melon plants released organic compounds such as arabinose and malic acid that can dissolve MnO<sub>x</sub> (Tsuji *et al.*, 2006). Soil microorganisms can also help Mn mobilization and immobilization, depending on soil conditions (Marschner, 1995). In aerated soils, microorganisms may mobilize Mn through MnO<sub>x</sub> reduction favored by H<sup>+</sup> root excretion. In contrast, Mn-oxidizing bacteria can decrease Mn availability in aerated and calcareous soils or in poorly aerated and/or submerged soils. Another key factor in the Mn dynamics in soil is organic matter (OM). Given that OM is negatively charged, it has a great Mn adsorption capacity, forming Mn complexes which decrease the amount of exchangeable Mn. However, the Mn adsorbed by OM can be exchanged by the H<sup>+</sup> released from the roots (Bradl, 2004).

#### **MANGANESE TRANSPORT AND ACCUMULATION IN PLANTS**

As mentioned above, reduced Mn (Mn<sup>2+</sup>) form is the only available metal form for plants. It can be taken up via an active transport system in epidermal root cells and transported as divalent cation Mn<sup>2+</sup> into the plants (Marschner, 1995; Gherardi and Rengel, 2003; Pittman, 2005). Manganese uptake by roots is characterized as a biphasic process. The initial and rapid uptake phase is reversible and non-metabolic, with Mn<sup>2+</sup> and Ca<sup>2+</sup> or



**Figure 1.** The role of soil rhizosphere in the mobilization and immobilization of Mn in soils.

other cations being freely exchanged in the rhizosphere. In this phase,  $Mn^{2+}$  appears to be adsorbed by the negatively charged cell wall constituents of the root-cell apoplastic spaces (Humphries *et al.*, 2007; Clarkson, 1988). The second phase is slow, with  $Mn^{2+}$  being less readily exchanged. Its uptake into the symplast is dependent on plant metabolism (Maas and Moore, 1968), although the exact mechanisms are not clear (Humphries *et al.*, 2007). It has been shown that in transgenic tobacco transformed with a tomato root protein with a metal binding site at its N-terminus (LeGlp1), Mn binds to this protein. This strongly suggests the involvement of LeGlp1 in Mn uptake from the soil (Takahashi and Sugiura, 2001). Kinetic measurements have demonstrated 100 to 1,000 times higher rates of Mn transport than the estimated plant requirement for this element (Clarkson, 1988). These transport rates

are explained by the high capacity of ion carriers and channels in the Mn ion transportation through the plasma membrane at a speed of several hundred to several million ions per second per protein molecule (Humphries *et al.*, 2007). According to these authors, Mn distribution from root cells within the whole plant involves primary transport in the xylem, transference from the xylem to the phloem and re-translocation into the phloem. Xylem transport from roots to the above-ground parts of plants is performed by the transpiration stream, whereas phloem transport is more selective, taking place from sources to sinks (Marschner, 1995). Nonetheless, a low mobility in phloem has been reported for Mn, and its redistribution may depend on the plant species and stages of development (Herren and Feller, 1994). In fact, it has been reported that Mn transport from roots to grains is frequently insufficient at

a mature stage of wheat. The relatively poor Mn mobility in the phloem emphasizes the importance of xylem in the transport of this element, even in wheat grain discharge (Rengel, 2001).

Manganese generally tends to accumulate predominantly in the plant shoots than in the roots, as demonstrated in Mn labeling experiments with  $^{54}\text{Mn}$  at an early stage of wheat (*Triticum aestivum* cv. Arina) development, where a fast Mn transport from roots to shoots was visualized in the xylem and was essentially immobile in the phloem (Page and Feller, 2005). Similar effects on Mn translocation have been shown by the same technique in young (28 days) white lupine plants (*Lupinus albus*) (Page *et al.*, 2006). Nevertheless, Mn was present in a large amount in the root system, hypocotyls and stem in older lupine plants, immediately after the labeling phase (day 0). Seven days later (day 7) almost all  $^{54}\text{Mn}$  had moved to the youngest fully expanded leaves and only a small fraction to the other leaves. Mn accumulation was observed in the periphery of the oldest leaves. These authors reported that Mn was rapidly released from the roots into the xylem, reaching photosynthetically active leaves via the transpiration stream. Furthermore, the low mobility of this element via phloem in the shoot may be due to a restricted loading of soluble Mn into the phloem or by insolubilization in the leaves, although the issue remains to be clarified (Page *et al.*, 2006). Page and Feller (2005) emphasized that little is known about the mechanisms involved in the loading of Mn into the phloem and the chemical transport forms.

In addition to the long distance transport of Mn, short distance transport mechanisms are important for the translocation of this metal into the cell and cell organelles. These mechanisms involve Mn translocation throughout the

plasma membrane and the biomembranes of organelles (Ducic and Polle, 2005; Pittman, 2005). Possible mechanisms of homeostasis and Mn transport, based on studies performed in yeast (*Saccharomyces cerevisiae*) cells and in *Arabidopsis thaliana* plants, have been discussed (Delhaize *et al.*, 2007; Reddi *et al.*, 2009). They pointed out that transport proteins play an important role for the maintenance of adequate Mn concentrations in the cytoplasm. Moreover, a variety of metal transporter family proteins with a broad-specificity such as  $\text{Fe}^{2+}$  and  $\text{Ca}^{2+}$  transporters have also the ability to transport Mn into the plant cells (Pittman, 2005). Migocka and Klobus (2007) demonstrated the activation of an antiport system in *Cucumis sativus* with different affinities for Pb, Mn, Ni and Cd in the root plasma membrane. This antiport system participates as part of the general defense mechanism activated under heavy metal stress. Using tonoplast-enriched vesicles, Shigaki *et al.*, (2003) suggested that a Cd/H antiporter might also be involved in Mn accumulation in vacuoles.

In Table 1 we have summarized some of the metal transporter proteins, their families, cellular localization and the transported metals. Transporter proteins are described according to their localization. In the plasma membrane, four  $\text{Mn}^{2+}$  uptake transporters are identified: AtIRT1, Nramp, AtYSL (Ducic and Polle, 2005; Pittman, 2005) and PHO84 (Ducic and Polle, 2005). IRT1 can transport Mn when expressed in yeast (Korshunova *et al.*, 1999). Nramp is also considered a metal transporter protein that can transport Mn away from other cations. This protein may be localized in the tonoplast rather than in the plasma membranes (Thomine *et al.*, 2003). AtYSL belongs to the yellow stripe-like (YSL) proteins, also involved in metal-complex transport in the plasma

membrane (Roberts *et al.*, 2004). This complex can be formed by nicotianamine (NA), which is a strong chelator of metals including  $Mn^{2+}$  (Pittman, 2005). PHO84 is a transporter protein identified in *S. cerevisiae* and it has a high-affinity phosphate uptake (Mitsukawa *et al.*, 1997). Luk *et al.* (2003) reported a new form in Mn transport as  $MnHPO_4$ . However, there is no evidence for  $Mn^{2+}$  or  $MnHPO_4$  accumulation by a plant phosphate transporter (Pittman, 2005).

Some transport proteins have been related to  $Mn^{2+}$  transport and accumulation into the intracellular compartments, such as the vacuole. It has been suggested that a metal transporter (specifically antiporter CAX2, calcium exchanger 2) originally identified as a  $Ca^{2+}$  transporter (which can also transport  $Cd^{2+}$ ) located in the cytosol. It has also the ability to transport Mn to the vacuole in tobacco plants (*Nicotiana tabacum*) and yeast (Hirschi *et al.*, 2000; Pittman, 2005) (Table 1). ATP-binding cassette (ABC) protein transporters are considered to be involved in detoxification processes (Martinoia *et al.*, 2002). Studies on cyanobacteria also suggested the putative role of these proteins in  $Mn^{2+}$  transport (Bartsevich and Pakrasi, 1996).

Another protein, considered indirectly as a metal transport protein is the ShMTP1, which is able to sequester metal ions within cells or efflux them out of the cells (Delhaize *et al.*, 2003). Therefore, these authors considered it a metal-tolerant protein. Ducic and Polle, (2005) and Pittman (2005) highlighted that, despite the available information about Mn transport across membranes in plant cells, the Mn transport and efflux strategies into the mitochondria, chloroplasts and Golgi are not completely understood. Nonetheless, Mills *et al.* (2008) have recently identified a Ca-ATPase that also transports  $Mn^{2+}$  into Golgi apparatus (Table 1).

## **MANGANESE AS AN ESSENTIAL ELEMENT IN PLANT METABOLISM**

The main Mn role in photosynthesis is its involvement in the water-splitting system of photosystem II (PSII), which provides electrons necessary for photosynthetic electron transport. In water photolysis, a group of four Mn atoms (Mn cluster) is associated with the oxygen evolving complex (OEC) bound to the reaction center protein (D1) of PSII (Goussias *et al.*, 2002). The Mn cluster in PSII accumulates four positive charges, which oxidize two water molecules, releasing one  $O_2$  molecule and four protons. Therefore, this metal cluster is considered a catalyst compound of water oxidation (Zouni *et al.*, 2001), where Mn ions are close to a redox-active tyrosine residue (Z and D) (Goussias *et al.*, 2002).

Manganese also plays a role in ATP synthesis (Pfeffer *et al.*, 1986), in RuBP carboxylase reactions (Houtz *et al.*, 1988) and the biosynthesis of fatty acids, acyl lipids and proteins (Ness and Woolhouse, 1980). In addition, Mn plays a primary role in the activation and as cofactor of various enzymes in plants (~35) (Burnell, 1988), such as: Mn-superoxide dismutase, Mn-catalase, pyruvate carboxylase and phospho-enolpyruvate carboxykinase (Ducic and Polle, 2005). Manganese is also essential for the biosynthesis of chlorophyll (through the activation of specific enzymes), aromatic amino acids (tyrosine), secondary products, like lignin and flavonoids (Lidon *et al.*, 2004). It also participates in the biosynthetic pathway of isoprenoids (Lidon *et al.*, 2004) and assimilation of nitrate (Ducic and Polle, 2005). Hence, Mn is involved in metabolic processes such as respiration, photosynthesis, synthesis of aminoacids and hormone activation (indol acetic acid, IAA) throughout the IAA-oxidases (Burnell, 1988).

**Table 1.** Some transporter proteins implicated in Mn<sup>2+</sup> and other cations transport and their cellular localization. (Summarized from Ducic and Polle, 2005; Pittman, 2005 and Mills *et al.*, 2008). Abbreviations as follows: At= *Arabidopsis thaliana*; Sh= *Stylosanthes hamata*).

Transporter proteins	Protein cellular localization and protein family transporters	Transported ions
AtIRT1	Plasma membrane protein (ZIP, zinc-regulated transporter/iron regulated transporter (ZRT/IRT1) related protein) family transporter	Mn <sup>2+</sup> and Fe <sup>2+</sup> , Zn <sup>2+</sup> and Cd <sup>2+</sup> under Fe-deficiency conditions.
AtECA1	Endoplasmic reticulum (ER) Ca <sup>2+</sup> - and Mn <sup>2+</sup> -transporting P-type ATPase	Ca <sup>2+</sup> and Mn <sup>2+</sup>
AtCAX2	Vacuolar cation/H <sup>+</sup> antiporter CAX (the cation exchanger)	Mn <sup>2+</sup> , Ca <sup>2+</sup> and Cd <sup>2+</sup>
AtNramp3	Vacuolar Nramp transporter. Also, possible plasma membrane localization (Nramp?)	Mn <sup>2+</sup> , Fe <sup>2+</sup> and Cd <sup>2+</sup> in Fe-deficiency conditions
ShMTP1	Vacuolar-localized cation diffusion facilitator (CDF) family transporter	Related to the <i>Stylosanthes hamata</i> Mn <sup>2+</sup> transporter
ABC	Vacuolar- localized ATP binding cassette transporter families	Related to the cyanobacterium <i>Synechocystis</i> Mn <sup>2+</sup> transporter
AtOPT3	Probably located in the plasma membrane (AtOPT3?). It is an oligopeptide transporter-like protein (OPT).	Possible transport of Cu <sup>2+</sup> and Fe <sup>2+</sup> and Mn <sup>2+</sup>
AtYSL	Probably located in the plasma membrane. Yellow stripe-like transporter with equivalent function to rice OsYSL2, a Mn <sup>2+</sup> -nicotianamine (NA) and Fe <sup>2+</sup> -NA transporter	Mn <sup>2+</sup> and Fe <sup>2+</sup>
AtECA3	Ca <sup>2+</sup> transporters (Ca-ATPases) in Golgi.	Mn <sup>2+</sup> and Ca <sup>2+</sup>
PHO84	Probably located in the plasma membrane. It is MnHPO <sub>4</sub> transporter.	Mn binding to phosphate.

As a cofactor of superoxide dismutase (SOD), manganese participates in the plant's defense against oxidative stress, produced by elevated levels of activated forms of oxygen and free radicals (reactive oxygen species, ROS), which are harmful to plants. It has been proposed that Mn can act as a scavenger of superoxide ( $O_2^{\cdot-}$ ) and hydrogen peroxide ( $H_2O_2$ ). However, this mechanism is still unclear (Ducic and Polle, 2005).

Manganese SOD (MnSOD) belongs to the group of metal-containing SOD enzymes which are classified according to their metal cofactor: iron SOD (Fe-SOD), localized in the chloroplast; copper-zinc SOD (Cu/Zn-SOD), located in the chloroplast, cytosol, and possibly in the extracellular space. Manganese SOD is found mainly in mitochondria (Clemens *et al.*, 2002) and peroxisomes (Alscher *et al.*, 2002). MnSOD may play an important role in the adaptive responses of plant cells under environmental stresses such as salt stress, enhancing their tolerance. This tolerance has been demonstrated in transgenic plants of *Arabidopsis*, where Mn-SOD was overexpressed (Wang *et al.*, 2004). Similarly, in tomato transgenic plants which overexpressed MnSOD, an improvement in tolerance to NaCl stress in seedlings was concomitant with an improvement in seed germination and root development (Wang *et al.*, 2007).

#### **MANGANESE PHYTOTOXICITY AND INJURY SYMPTOMS**

As an essential micronutrient, low Mn levels are absolutely necessary for normal nutrition and development of plants. Normal Mn contents of leaves differ greatly between species (30-500 mg kg<sup>-1</sup> Mn dry mass, Clarkson, 1988). Nonetheless, when it is present in

excessive amounts, it is extremely toxic to plant cells (Migocka and Klobus, 2007). The injury extent of Mn toxicity is approximately proportionate to the concentration of accumulated Mn excess. However, there is considerable inter- and intra-specific variation among Mn levels that induce toxicity as well as the symptoms of this toxicity in plant species (Foy *et al.*, 1988).

In addition to a decrease in growth rate, symptoms of Mn toxicity such as chlorosis (interveinal and marginal) and necrotic leaf spots are very common and have been reported in the whole plants of canola (Moroni *et al.*, 2003), clover (Rosas *et al.*, 2007), ryegrass (Mora *et al.*, 2009) as well as in leaves of barley and cowpea (Demirevska-Kepova *et al.*, 2004; Führes *et al.*, 2008) (Table 2). Necrotic brown spots and chlorotic leaves are frequently reliable indicators of the severity of Mn toxicity in plants (Wissemeier and Horst, 1991). The interveinal chlorosis due to Mn toxicity can have an appearance similar to that observed under Fe deficiency (Sarkar *et al.*, 2004). Moreover, the Mn toxicity is intensified when other available elements such as Ca, Mg, K, Fe and Si are in a low quantity (Abou *et al.*, 2002). However, a decrease in productivity by Mn toxicity without the appearance of leaf visual symptoms is sometimes observed (Miner and Sims, 1983). It is important to know that all these symptoms induced by Mn toxicity are preceded by an alteration of the photosynthetic apparatus and the photosynthetic performance of plants.

Other studies have shown that in rice (*Oryza sativa* cv. Safari) exposed to Mn excess in a nutrient solution, Mn was predominantly accumulated in leaves compared with roots (Lidon, 2001), whereas in *Sinapis alba* Mn mostly accumulated in the shoots (Farasova and Beinrohr, 1998).

**Table 2.** Symptoms of Mn toxicity and Mn concentrations in organs of some plant species subjected to toxic Mn concentrations according references from the last decade (earlier references in the text). Mn treatments were performed in nutrient solutions with MnCl<sub>2</sub> or MnSO<sub>4</sub>.

Species	Mn treatment	Mn concentrations in different plant organs	Symptoms of Mn toxicity	References
Rice ( <i>Oryza sativa</i> L.)	583 µM	Shoots: 2020 µg g <sup>-1</sup> dw	Decrease of shoot growth rate.	Lidon and Teixeira (2000a)
Barley ( <i>Hordeum vulgare</i> L.)	1830 µM 18300 µM	Leaves: 656 mg g <sup>-1</sup> dw Leaves: 1615 mg g <sup>-1</sup> dw	Dark-brown necrotic spots, individually or in groups.	Demirevska-Kepova et al. (2004)
Ryegrass ( <i>Lolium perenne</i> L.)	355 µM	Shoot: 2357 mg kg <sup>-1</sup> Roots: 2408 mg kg <sup>-1</sup>	Chlorotic leaves. Decrease of dry weight in roots.	Rosas et al. (2007)
Ryegrass ( <i>L. perenne</i> L.)	150 µM	Shoot: 902 mg kg <sup>-1</sup> Roots: 1342 mg kg <sup>-1</sup>	Dry weight reduction. Dry weight reduction.	Mora et al. (2009)
Clover ( <i>Trifolium repens</i> L.)	355 µM	Shoot: 2050 mg kg <sup>-1</sup> Root: 7481 mg kg <sup>-1</sup>	Reddish borders on leaves. Decrease of dry weight.	Rosas et al. (2007)
Soybean ( <i>Glycine max</i> L.)	200 µM	Leaves: 806 mg kg <sup>-1</sup> Roots: 502 mg kg <sup>-1</sup>	Chlorotic leaves. No visual symptoms, increase in root diameter.	Lavres Jr et al. (2009)
Cowpea ( <i>Vigna unguiculata</i> L.)	50 µM	Leaves: ~ 25 µmol g <sup>-1</sup> dw	Brown spots on leaves.	Fühns et al. (2008, 2009)
Canola ( <i>Brassica napus</i> L.)	200 µM	Shoot: ~ 3500 µg g <sup>-1</sup> dw	Necrotic leaf spots, chlorosis in leaf margin.	Moroni et al. (2003)
<i>Juncus effuses</i> L. (wetland plant)	500 µM	176 mg kg <sup>-1</sup>	Reduction in plant dry biomass and height, no phytotoxic visual symptoms.	Najeeb et al. (2009)
<i>Populus cathayana</i>	1000 µM	Leaves: 713 mg kg <sup>-1</sup> dw	Decrease in shoot height, total biomass, and total leaf area.	Lei et al. (2007)

Despite the importance of Mn excess in the photosynthetic performance of plants, only a few studies about this issue are available. A reduction in photosynthesis, in chlorophyll *a* and *b* contents and their biosynthesis, as well as a reduction in carotenoids is frequently found in plants and also in algae under Mn excess (Macfie and Taylor, 1992; Hauck *et al.*, 2003). In rice cultivated at different Mn concentrations (from 2.3 to 583  $\mu\text{M}$ ), a significant decrease in chlorophyll *a* content has been reported at the highest Mn concentration (Lidon and Teixeira, 2000a). Nable *et al.* (1988) reported an early inhibition of photosynthesis concomitant with a high Mn accumulation in the leaves in *Nicotiana tabacum* cultivated in nutrient solutions with a Mn excess (1,000  $\mu\text{M}$ ). The authors concluded that the inhibition of photosynthesis is an early indicator for Mn toxicity in tobacco leaves. Lidon *et al.*, (2004) also observed a decline in net photosynthesis ( $\mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$ ) and photosynthetic capacity ( $\mu\text{mol O}_2 \text{ m}^{-2}\text{s}^{-1}$ ) in rice plants subjected to 9.1 and 36.4  $\mu\text{M}$  Mn treatment, respectively, with no changes in the levels of the ratio between variable (Fv) and maximum (Fm) chlorophyll fluorescence (Fv/Fm). However, an increase in the photochemical quenching and the quantum yield of non-cyclic electron transport was found up to 36.4  $\mu\text{M}$  Mn. Since Mn is accumulated in thylakoids, this element may interfere with thylakoid stacking, decreasing net photosynthesis (Lidon and Teixeira, 2000a).

Similarly to Mn toxicity, Mn deficiency also depressed leaf photosynthetic capacity in plants of *Carya illinoensis*, primarily by reducing the number of PSII units per unit leaf area, but these PSII units maintained abilities similar to those of the control plants (Henriques, 2003). Increased Mn concentrations (0 – 1,000 and 10,000  $\mu\text{M}$

Mn) in wheat plants inhibited the biosynthesis of chlorophyll and carotenoids, inducing a decrease in photosynthetic electron transport rates and therefore a decrease in the rate of photosynthesis (Macfie and Taylor, 1992). Higher Mn concentrations are also involved in a shortening of root and shoot and in a decreased chlorophyll concentration in the Mn accumulator plant, *Alyssum murale* (Abou *et al.*, 2002). Recently, Amao and Ohashi (2008) suggested that high Mn amounts in spinach leaves inhibited the activity of oxygen evolved complex of PSII.

### **Manganese toxicity and oxidative stress**

Mn toxicity can also trigger oxidative stress in plant cells (Demirevska-Kepova *et al.*, 2004). As a toxic metal, Mn can cause metabolic alterations and macromolecular damage that disrupt the cell homeostasis (Hegedüs *et al.*, 2001; Polle, 2001). According to Lynch and St. Clair (2004), Mn toxicity in plants generates reactive oxygen species (ROS), mainly OH, the most reactive oxidant and harmful species in cells (Lidon and Henriques, 1993). With respect to the oxidative stress responses to Mn excess, Lidon and Teixeira (2000b) reported that at the first growth stages of rice, two kinetic phases can be distinguished: in the first one, there is an increase in Mn accumulation in the thylakoid lamellae (McCain and Markley, 1989), which inhibits electron leakage from the Hill and Mehler reactions, limiting ROS formation. In the second kinetic phase, higher Mn amounts inhibit the non-cyclic photophosphorylation process, promoting an increase in ROS production that parallels an injury increase in the thylakoid peroxidase system (Lidon and Teixeira, 2000b). These authors concluded that Mn excess increases the disorganization in chloroplast lamellae,

but elevated activity of superoxide dismutase (SOD) still limits cell damage.

It has also been reported that in *Cucumis sativus* plants both Mn excess and optimum light intensity determine an enhancement in oxidative stress by increased Mn content in the tissues concomitant with an inhibition of plant growth (Shi *et al.*, 2006). Investigations performed by González *et al.*, (1998) showed that lipid peroxidation was not induced by Mn-toxicity stress in the mature leaves of *Phaseolus vulgaris*, although other studies have shown that lipid peroxidation occurred in isolated chloroplast of wheat (Panda *et al.*, 1986). However, González *et al.* (1998) mentioned that this damage process could be related to the development stage of leaves, with the damage being more intense in immature than in mature leaves.

The geographical origin of the species and climatic conditions also affect the degree of Mn toxicity in plant species or populations as shown in two populations of *Populus cathayana*, coming from a wet and dry climate cultivated in an acid solution with increasing Mn concentrations (Lei *et al.*, 2007). The results showed that the wet climate population accumulated more Mn in plant tissues especially in leaves, decreasing their growth, chlorophyll contents, and activities of antioxidant enzymes than the dry climate population.

#### **TOLERANCE MECHANISMS TO MANGANESE TOXICITY**

The ability of plant for to grow and survive in a metal-contaminated environment, commonly called resistance, can be achieved through different mechanisms: avoidance and/or tolerance. The former involves a protective role that prevents the metal ions from entering the cytoplasm of plant cells (Blamey *et al.*,

1986; Marschner, 1991). The latter strategy (tolerance) implies a detoxification of metal ions after they have crossed the plasma cell membrane or internal organelle biomembranes (Macfie *et al.*, 1994). However, as shown in the following paragraphs, the differentiation of these mechanisms in the pertinent literature is very confusing.

The sequestering of Mn in the apoplast is considered an avoidance mechanism. However, some researchers have included this feature as a tolerance mechanism. For example, Horst *et al.*, (1999) suggested that tolerance to Mn excess in *Vigna unguiculata* is performed by the reduction of Mn<sup>2+</sup> activity in the apoplast throughout complexation by organic acids. In this species, symptoms such as brown leaf cell spots are also identified as oxidized Mn, and phenolic compounds present in the cell walls are considered a Mn tolerance mechanism (Wissemeier and Horst, 1992). On the other hand, the Mn<sup>2+</sup> oxidation by peroxidases in the cell walls of roots is considered by Marschner (1991) as an avoidance mechanism, although the existence of such a mechanism was not considered in the study of Horiguchi (1987). In this study, it is suggested that oxidized Mn deposition in plant tissues corresponds to a tolerance mechanism to Mn toxicity, with *Cucumis sativus* being more tolerant to high Mn deposition in tissues than melon (*Cucumis melo*). Blamey *et al.*, (1986) reported the accumulation and secretion of Mn<sup>2+</sup> in and around the trichomes of sunflower plants (*Helianthus annuus*) as a Mn tolerance mechanism. Another strategy that plants use to prevent the toxic effects of heavy metals as well as of Mn can be the efflux from the cell. In this process, the Mn cell is delivered into the Golgi apparatus and finally exported from the cell via secretory pathway vesicles that carry the metal to the cell surface (Ducic and Polle, 2005).

Summarizing the available literature regarding tolerance mechanisms to Mn, it appears that the main Mn tolerance mechanism is the sequestration by organic compounds in metabolically less active cells or organelles. The vacuole is considered the biggest and most important compartment, because it can store many toxic compounds (Pittman, 2005). Hence, an increase in phenolic compounds was found in the hydrophyte (*Trapa natans*) leaves exposed to high Mn levels (130  $\mu\text{M}$ ) (Baldissserotto *et al.*, 2004). These compounds chelate Mn inside the vacuole, segregating the metal ion in the protoplasm and thus reducing the damage (Davis *et al.*, 2001). A similar key role has recently been assigned to oxalic acid in Mn internal sequestration by chelating specifically the Mn excess in vacuoles of Mn hyperaccumulator plants (*Phytolacca americana*) (Dou *et al.*, 2008). Furthermore, it has been observed that a Mn excess can accumulate dark material in the vacuoles, probably for deposition of Mn oxides or an increase in polyphenol oxidase activity in *Citrus volkameriana* plants (Papadakis *et al.*, 2007). Similarly, studies about the effect of Mn excess in varieties of the conifer *Pseudotsuga menziesii* showed dark deposits of Mn-complexes in plant root vacuoles, which were associated with phosphate, establishing “free”  $\text{Mn}^{2+}$  to form insoluble complexes, giving a greater tolerance (Ducic and Polle, 2007). Additionally, these studies showed that both, root elongation and biomass production were inhibited by Mn treatment above 2,500  $\mu\text{M}$ , mainly in *P. menziesii* var. *glauca*, confirming that it is a species with lower tolerance than *P. menziesii* var. *viridis*.

Another strategy to confer Mn tolerance inside plant cells is associated with several metal transporter proteins identified in the Mn transport mechanisms described in section 4. Metal transporter proteins located in the tonoplast (CAX2 in

tobacco plants, ShMTP1 in *Arabidopsis* plants) conferred greater tolerance on elevated  $\text{Mn}^{2+}$  levels due to the internal sequestering of this element (Hirschi *et al.*, 2000; Delhaize *et al.*, 2003). Other transporter proteins (ECA1) can maintain low cytosolic Mn, since it moves into the endoplasmic reticulum (Wu *et al.*, 2002). In this research, ECA1 was able to reduce cytosolic  $\text{Mn}^{2+}$ , preventing an interference with the internal distribution of other ions ( $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$  or  $\text{Ca}^{2+}$ ).

Unlike other metal stresses, the accumulation of Mn excess does not have a single cell target. Depending on the plant species, different organelles can serve as stores for this accumulation (Lidon *et al.*, 2004). Under high Mn levels apart from vacuoles, chloroplasts are important sinks of this metal in *Citrus volkameriana*. This feature, together with the larger size of this organelle, is considered to be an adaptive response of this plant to Mn excess (Papadakis *et al.*, 2007). In general, cultivated plants like rice are considered tolerant to Mn toxicity (Lidon, 2001) because their leaf tissues can accumulate from 5 to 10 times more Mn than other grasses (Foy *et al.*, 1978). The Mn tolerance mechanism in this species included the inhibition of apoplastic influx from the cortex toward the stele and symplastic Mn assimilation in the shoot protoplast, where the chloroplast is the main target (Lidon, 2001).

The distribution of Mn excess in both roots and shoots is dependent on plant species and genotype. Early research associated Mn tolerance in some plants with a greater retention of Mn excess in the roots, as mentioned by Andrew and Hegarty (1969) in regard to tropical and temperate legume species. The root retention of heavy metals has been attributed to the formation of metal complexes in roots (Foy *et al.*, 1978). Metals with high electro-negativity

accumulate in roots in larger amounts than metals with low electro-negativity. In the latter instance, Mn and Zn metals can easily be translocated to the tops (Chino, 1981).

In proteomics studies, a comparison of Mn-sensitive and Mn-tolerant cultivars of cowpea (*Vigna unguiculata*.) has shown relevant features of leaf apoplast in the expression of Mn toxicity: formation of brown spots, induction of callose formation and an enhanced release of phenols and peroxidases into the apoplast (Fecht-Christoffers *et al.*, 2003; Fecht-Christoffers *et al.*, 2006). Specific proteins involved in the regulation processes, such as CO<sub>2</sub> fixation, stabilization of the Mn cluster of the photosystem II, pathogenesis-response reactions and protein degradation, were affected at low or high Mn levels, mainly in the Mn-sensitive cowpea cultivar (Führs *et al.*, 2008). Chloroplastic proteins, which are important for CO<sub>2</sub> fixation and photosynthesis, were of lower abundance upon Mn-induced stress, suggesting the scavenging of metabolic energy for a specific stress response. Führs *et al.*, (2008) concluded that a coordinated interplay of apoplastic and symplastic reactions seems to be important during the Mn-stress response in plants.

To alleviate metal toxicity in plants, the antioxidant systems are also considered an important tolerance mechanism. The antioxidant systems include antioxidant enzyme “scavengers” such as superoxide dismutase (SOD), catalase (CAT), peroxidases (phenol peroxidase, POX, ascorbate peroxidase, APX, guaiacol peroxidase, GPX) and the non-enzymatic antioxidant molecules: ascorbate,  $\alpha$ -tocopherol, carotenoids, flavonoids and glutathione (Foyer and Noctor, 2003; Apel and Hirt, 2004). These constitute other mechanisms against ROS, produced by Mn toxicity.

Higher activities of antioxidant enzymes are found in response to a Mn excess in woody plants (Lei *et al.*, 2007), in herbs such as white clover (*Trifolium repens*.) and in ryegrass (*Lolium perenne*), which suggest a lower oxidative stress (Rosas *et al.*, 2007; Mora *et al.*, 2009). Another plant tolerance mechanism to metal toxicity is associated with lower metal uptake and translocation to other organs (Hall, 2002). The exudation of organic acid anions (carboxylates) to the rhizosphere may minimize the absorption by roots of such metals as aluminium and nickel (Ma *et al.*, 2001; Yang *et al.*, 1997). There are few reports regarding the release of organic acid anions by roots in the event of Mn toxicity (González and Lynch, 1999). Recently, Mora *et al.* (2009) reported that the root exudates of oxalate and citrate may decrease Mn availability in the rhizosphere, enhancing their Mn tolerance in ryegrass subjected to Mn toxicity.

Other nutrient applications can help minimize the effects of Mn toxicity. Thus, it has been reported that silica (Si) addition significantly decreases lipid peroxidation caused by an Mn excess decreasing the symptoms of Mn phytotoxicity and improving plant growth in some plants (Iwasaki *et al.*, 2002; Shi *et al.*, 2005). There is also an association between Mn toxicity and the decrease in Ca concentration in barley plants (*Hordeum vulgare*), indicating a competition and a specific interaction during the absorption and/or translocation of these elements (Alam *et al.*, 2001). Another study, in which Ca is applied to reduce the Mn toxicity, showed that Ca additions inhibited Mn translocation from roots to shoots in barley plants, but did not affect the Mn absorption in roots (Alam *et al.*, 2006). This suggested that Ca could avoid Mn accumulation in shoots, protecting the photosynthetic apparatus from the dangerous effect of an

excess of Mn. In barley, Mn toxicity could also be repressed by high K contents, which inhibit both the absorption and the translocation of Mn (Alam *et al.*, 2005).

## CONCLUSIONS AND PERSPECTIVES

Manganese is considered an essential micronutrient for the metabolic process in plants. Nevertheless, both deficiency and excess alter these processes. Acid soils make excessive Mn amounts toxic for the plants. Mn toxicity is a world-wide problem in areas with acid soils. This toxicity alters physiological, biochemical and molecular processes at the cell level. It is crucial to know the limitations of these soils for the purpose of soil-plant interaction management, especially in relation to the presence and Mn excess. Thus, the knowledge of Mn uptake, translocation, accumulation and resistance mechanisms in crop plants under Mn excess and toxicity is of great importance to crop improvement.

Most molecular and physiological approaches to Mn transport inside plant cells have recently been analyzed, as these are useful tools for understanding resistance mechanisms. However, the question as to which is the best candidate gene related to Mn toxicity in acid soil continues to elude plant scientists due to Mn toxicity in plants being a complex trait and involving multiple physiological and biochemical mechanisms and a wide array of genes. The understanding of these mechanisms will contribute to improving the yield and quality of cultivated plants in acid soils. Future efforts for developing Mn-tolerant plants should take all these aspects into account.

## ACKNOWLEDGMENTS

Financial support for this review was provided by Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT N° 1080372) and Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) Doctoral Fellowship of the Government of Chile.

## REFERENCES

- Abou, M., Symeonidis, L., Hatzistavrou, E., Yupsanis, T. 2002.** Nucleolytic activities and appearance of a new DNase in relation to nickel and manganese accumulation in *Alyssum murale*. *J. Plant Physiol.* 159, 1087-1095.
- Adriano, D.C. 2001.** Trace elements in terrestrial environments. Biogeochemistry, bioavailability and risks of metals. Springer-Verlag, New York, 867 p.
- Alam, S., Kamei, S., Kawai, S. 2001.** Amelioration of manganese toxicity in barley with iron. *J. Plant Nutr.* 28, 1421-1433.
- Alam, S., Akiha, F., Kamei, S., Huq, S., Kawai, S. 2005.** Mechanism of potassium alleviation of manganese phytotoxicity in barley. *J. Plant Nutr.* 28, 889-901.
- Alam, S., Kodama, R., Akiha, F., Kamei, S., Kawai, S. 2006.** Alleviation of manganese phytotoxicity in barley with calcium. *J. Plant Nutr.* 29, 59-74.
- Alscher, R., Erturk, N., Heath, L. 2002.** Role of superoxide dismutase (SODs) in controlling oxidative stress in plants. *J. Exp. Bot.* 53, 1331-1341.
- Andrew, C.S., Hegarty, M.P. 1969.** Comparative responses to manganese excess of eight tropical and four temperate pasture legume species. *Aust. J. Agric. Res.* 20, 687-696.
- Amao, Y., Ohashi, A. 2008.** Effect of Mn ion on the visible light induced water oxidation activity of photosynthetic organ grana from spinach. *Catal. Commun.* 10, 217-220.

- Apel, K., Hirt, H. 2004.** Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373-379.
- Baldisserotto, C., Ferroni, L., Medici, V., Pagnoni, A., Pellizzari, M., Fasulo, M.P., Fagioli, F., Bonora, A., Pancaldi, S. 2004.** Specific intra-tissue responses to manganese in the floating lamina of *Trapa natans* L. *Plant Biol.* 6, 578-589.
- Bartsevich, V., Pakrasi, H.B. 1996.** Manganese transport in the cyanobacterium *Synechocystis* sp PCC 6803. *J. Biol. Chem.* 271, 26057-26061.
- Blamey, F., Joyce, D., Edwards, D., Asher, C. 1986.** Role of trichomes in sunflower tolerance to manganese toxicity. *Plant Soil.* 91, 171-180.
- Bradl, H. 2004.** Adsorption of heavy metal ions on soils and soils constituents. *J. Colloid. Interf. Sci.* 277, 1-18.
- Buchanan, B., Grusen, W., Jones, R. 2000.** Biochemistry and molecular biology of plants. American Society. of Plant Physiologists Maryland. 1367 p.
- Burnell, J. 1988.** The biochemistry of manganese in plants. In: R.D. Graham, R.J. Hannam, N.J. Uren. (eds). Manganese in Soil and Plants. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 125-137.
- Chino, M. 1981.** Metal stress in rice plants. In: K. Kitagisi, I. Yamane (eds). Heavy metal pollution in soils of Japan. Japan Scientific Societies Press, Tokyo, pp. 65-80.
- Clarkson, D.T. 1988.** The uptake and translocation of manganese by plant roots. In: R.D. Graham, R.J. Hannam, N.J. Uren. (eds). Manganese in Soil and Plants. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 101-111.
- Clemens, K.L., Force, D.A., Britt, R.D. 2002.** Acetate binding at the photosystem II oxygen evolving complex: an S2-State multiline signal ESEEM study. *J. Am. Chem. Soc.* 124, 10921-10933.
- Conyers, M., Uren, N., Helyar, K., Poile, G., Cullis, B. 1997.** Temporal variation in soil acidity. *Aust. J. Soil Res.* 35, 1115-1129.
- Davis, M.A., Pritchard, S.G., Boyd, R.S., Prior, S.A. 2001.** Developmental and induced responses of nickel-based and organic defenses of the nickel-hyperaccumulating shrub, *Psychotria douarrei*. *New. Phytol.* 150, 49-58.
- Delhaize, E., Kataoka, T., Hebb, D.M., White, R.G., Ryan, P.R. 2003.** Genes encoding proteins of the cation diffusion facilitator family that confer manganese tolerance. *Plant Cell* 15, 1131-1142.
- Delhaize, E., Gruber, B.D., Pittman, J.K., White, R.G., Leung, H., Miao, Y., Jiang, L., Ryan, P.R., Richardson, A.E. 2007.** A role for the AtMTP11 gene of *Arabidopsis* in manganese transport and tolerance. *Plant J.* 51, 198-210.
- Demirevska-Kepova, K., Simova-Stoilova, L., Stoyanova, Z., Holzer, R., Feller, U. 2004.** Biochemical changes in barley plants after excessive supply of copper and manganese. *Environ. Exp. Bot.* 52, 253-266.
- Dou, C., Fu, X, Chen, X, Shi, J., Chen, Y. 2008.** Accumulation and detoxification of manganese in hyperaccumulator *Phytolacca americana*. *Plant Biol.* 11, 664-670.
- Ducic, T., Polle, A. 2005.** Transport and detoxification of manganese and copper in plants. *Braz. J. Plant Physiol.* 17, 103-112.
- Ducic, T., Polle, A. 2007.** Manganese toxicity in two varieties of Douglas fir (*Pseudotsuga menziesii* var. *viridis* and *glauca*) seedlings as affected by phosphorus supply. *Funct. Plant Biol.* 34, 31-40.
- El-Jaoual, T., Cox, D. 1998.** Manganese toxicity in plants. *J. Plant Nutr.* 21, 353-386.
- Fageria, N., Baligar, V., Clark, R. 2002.** Micronutrients in crop production. *Adv. Agron.* 77, 185-268.
- Farasova, A., Beinrohr, E. 1998.** Metal-metal interaction in accumulation of V, Ni, Mo, Mn, and Cu in under and above ground parts of *Sinapis alba*. *Chemosphere* 36, 1305-1317.
- Fecht-Christoffers, M., Braun, H., Lemaitre-Guillier, C., VanDorsselaer, A., Horst, W. 2003.** Effect of manganese toxicity on the proteome of the leaf apoplast in cowpea. *Plant Physiol.* 133, 1935-1946.
- Fecht-Christoffers, M., Führs, H., Braun, H., Horst, W. 2006.** The role of hydrogen peroxide-producing and hydrogen peroxide-consuming peroxidases in the leaf apoplast of cowpea in manganese tolerance. *Plant Physiol.* 140, 1451-1463.
- Foy, C., Chaney, R., White, M. 1978.** The physiology of metal toxicity in plants. *Annu. Rev. Plant Physiol.* 29, 511-566.

- Foy, C., Scott, B., Fisher, J. 1988.** Genetic differences in plant tolerance to manganese toxicity. In: R.D. Graham, R.J. Hannam, N.J. Uren. (eds). *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 293–307.
- Foyer, C., Noctor, G. 2003.** Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiol. Plant.* 119, 355-364.
- Führs, H., Hartwig, M., Buitrago, L., Heintz, D., Van Dorsselaer, A., Braun, H., Horst, W. 2008.** Early manganese-toxicity response in *Vigna unguiculata* L. – a proteomic and transcriptomic study. *Proteomics* 8, 149-159.
- Führs, H., Götze, S., Specht, A., Erban, A., Gallien, S., Heintz, D., Van Dorsselaer, A., Kopka, J., Braun, H.P., Horst, W.J. 2009.** Characterization of leaf apoplastic peroxidases and metabolites in *Vigna unguiculata* in response to toxic manganese supply and silicon. *J. Exp. Bot.* 60, 1663–1678.
- Gherardi M., Rengel, Z. 2003.** Genotypes of lucerne (*Medicago sativa* L.) show differential tolerance to manganese deficiency and toxicity when grown in bauxite residue sand. *Plant Soil* 249, 287-296.
- Gherardi M., Rengel, Z. 2004.** The effect of manganese supply on exudation of carboxylates by roots of lucerne (*Medicago sativa*). *Plant Soil* 260, 271-282.
- González, A., Steffen, K., Lynch, J. 1998.** Light and excess manganese. *Plant Physiol.* 118, 493-504.
- González, A., Lynch, J. 1999.** Subcellular and tissue Mn compartmentation in bean leaf under Mn toxicity stress. *Aust. J. Plant Physiol.* 26, 811-822.
- Goussias, C., Boussac, A., Rutherford, W. 2002.** Photosystem II and photosynthetic oxidation of water: an overview. *Phil. Trans. R. Soc. Lond. B* 357, 1369-1381.
- Guest, C., Schulze, D., Thompson, I., Huber, D. 2002.** Correlating manganese X-ray absorption near-edge structure spectra with extractable soil manganese. *Soil Sci. Soc. Am. J.* 66, 1172-1181.
- Hall, J.L. 2002.** Cellular mechanisms for heavy metal detoxification and tolerance. *J. Exp. Bot.* 53, 1-11.
- Hauck, M., Paul, A., Gross, S., Raubuch, M. 2003.** Manganese toxicity in epiphytic lichens: chlorophyll degradation and interaction with iron and phosphorus. *Environ. Exp. Bot.* 49, 181-191.
- Hegedüs, A., Erdei, S., Horváth, G. 2001.** Comparative studies of H<sub>2</sub>O<sub>2</sub> detoxifying enzymes in green and greening barley seedlings under cadmium stress. *Plant Sci.* 160, 1085-1093.
- Henriques, F. 2003.** Gas exchange, chlorophyll a fluorescence kinetics and lipid peroxidation of pecan leaves with varying manganese concentrations. *Plant Sci.* 165, 239-244.
- Herren, T., Feller U. 1994.** Transfer of zinc from xylem to phloem in the peduncle of wheat. *J. Plant Nutr.* 17, 1587-1598.
- Hirschi, K., Korenkov, V., Wilganowski, N., Wagner, G.J. 2000.** Expression of *Arabidopsis* CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. *Plant Physiol.* 124, 125-133.
- Horiguchi, T. 1987.** Mechanism of manganese toxicity and tolerance of plants. II. Deposition of oxidized manganese in plant tissues. *J. Soil Sci. Plant Nutr.* 33, 595-606.
- Horst, W.J. 1988.** The physiology of manganese toxicity. In: R.D. Graham, R.J. Hannam, N.J. Uren (eds). *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 175–188.
- Horst, W., Fecht, M., Naumann, A., Wissemeier, A., Maier, P. 1999.** Physiology of manganese toxicity and tolerance in *Vigna unguiculata* (L.) Walp. *J. Plant Nutr. Soil Sci.* 162, 263-274.
- Houtz, R.L., Nable, R.O., Cheniae, G.M. 1988.** Evidence for effects on the in vivo activity of ribulose-biphosphate carboxylase/oxygenase during development of Mn toxicity in tobacco. *Plant Physiol.* 86, 1143-1149.
- Hue, N. 1988.** A possible mechanism for manganese toxicity in Hawaii soils amended with a low-Mn sewage sludge. *J. Environ. Qual.* 17, 473-479.
- Hue, N., Mai, Y. 2002.** Manganese toxicity in watermelon as affected by lime and compost amended to a Hawaiian acid Oxisol. *Hortscience* 37, 656-661.

- Humphries, J., Stangoulis, J., Graham, R. 2007.** Manganese. In: A. Barker, D. Pilbeam (eds). Handbook of Plant Nutrition, Taylor and Francis, USA, pp. 351-366.
- Iwasaki, K., Maier, P., Fecht, M., Horst, W. 2002.** Effects of silicon supply on apoplastic manganese concentrations in leaves and their relation to manganese tolerance in cowpea (*Vigna unguiculata* (L.)Walp). Plant Soil 238, 281-288.
- Kochian, L., Hoekenga, O., Piñeros, M. 2004.** How do crops plants tolerate acid soils? Mechanisms of aluminum tolerance and phosphorus efficiency. Annu. Rev. Plant Biolog. 55, 459-93.
- Kogelmann, W., Sharpe, W. 2006.** Soil acidity and manganese in declining and nondeclining sugar maple stands in Pennsylvania. J. Environ. Qual. 35, 433-441.
- Korshunova, Y., Eide, D., Clark, W., Guerinot, M., Pakrasi, H. 1999.** The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range. Plant Mol. Biol. 40, 37-44.
- Laha, S., Luthy, R. 1990.** Oxidation of aniline and other primary aromatic amines by manganese dioxide. Environ. Sci. Technol. 24, 363-373.
- Lavres J., Malavolta, E., Nogueira, N.L., Moraes, M.F., Rodrigues, A., Lanzoni, M., Pereira, C. 2009.** Changes in anatomy and root cell ultrastructure of Soybean genotypes under manganese stress. R. Bras. Ci. Solo 33,395-403.
- Lei, Y., Korpelainen, H., Li, C. 2007.** Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. Chemosphere 68, 686-694.
- Lidon, F. 2001.** Tolerance of rice to excess manganese in the early stages of vegetative growth. Characterization of manganese accumulation. J. Plant Physiol. 158, 1341-1348.
- Lidon, F.C., Barreiro, M., Ramalho, J. 2004.** Manganese accumulation in rice: implications for photosynthetic functioning. J. Plant Physiol. 161, 1235-1244.
- Lidon, F.C., Teixeira, M.G. 2000a.** Rice tolerance to excess Mn: Implications in the chloroplast lamellae and synthesis of a novel Mn protein. Plant Physiol. Biochem. 38, 969-978.
- Lidon, F.C., Teixeira, M.G. 2000b.** Oxy radicals production and control in the chloroplast of Mn-treated rice. Plant Sci.152, 7-15.
- Lidon, F.C., Henriques, F. 1993.** Oxygen metabolism in higher plant chloroplasts. Photosynthetica 29, 249-279.
- Luk, E., Jensen, L.T., Culotta, V.C. 2003.** The many highways for intracellular trafficking of metals. JBIC 8, 803-809.
- Lynch, J.P., St.Clair, S.B. 2004.** Mineral stress: the missing link in understanding how global climate change will affect plants in real world soils. Field Crop. Res. 90, 101-115.
- Ma, J.F. Ryan, P. R., Delhaize, E. 2001.** Aluminium tolerance in plants and the complexing role of organic acids. Trends Plant Sci. 6, 273-278.
- Maas E., Moore D. 1968.** Manganese absorption by excised barley roots. Plant Physiol. 43, 527-530.
- McCain, D.C., Markley, J:L. 1989.** More manganese accumulates in maple sun leaves than in shade leaves. Plant Physiol. 90: 1417-1421.
- Macfie, S., Taylor, G. 1992.** The effects of excess manganese on photosynthetic rate and concentration of chlorophyll in *Triticum aestivum* grown in solution culture. Physiol. Plant. 85, 467-475.
- Macfie, S., Cossins, E., Taylor, G. 1994.** Effects of excess manganese on production of organic acids in Mn-tolerant and in Mn-sensitive cultivars of *Triticum aestivum* L. (wheat). J. Plant Physiol. 143, 135-144.
- Marschner, H. 1991.** Mechanisms of adaptation of plants to acid soils. In: R. Wright, V. Baligar, R. Murrmann (Eds). Development in Plant and Soil Sciences. Plant-soil interactions at low pH. Kluwer Academic Publisher, West Virginia, pp. 1-20.
- Marschner, H. 1995.** Mineral Nutrition of Higher Plants. Academic Press, San Diego, 889 p.
- Martinoia, E., Klein, M., Geisler, M., Bovet, L., Forestier, C., Kolukisaoglu, H., Mueller-Roeber, B., Schulz, B. 2002.** Multifunctionality of plant ABC transporters: more than just detoxifiers. Planta 214, 345-355.
- Migocka M., Klobus G. 2007.** The properties of the Mn, Ni and Pb transport operating at plasma membranes of cucumber roots. Physiol. Plant. 129, 578-587.
- Mills, R., Doherty, M., López-Marqués, R., Weimar, T., Dupree, P., Palmgren, M.,**

- Pittman, J., Williams, L. 2008.** ECA3, a Golgi-localized P2A-type ATPase, plays a crucial role in manganese nutrition in *Arabidopsis*. *Plant Physiol.* 146, 116-128.
- Miner, G., Sims, J. 1983.** Changing fertilization practices and utilization of added plant nutrient for efficient production of tobacco. *Recent Adv. Tobacco Sci.* 9, 4-76.
- Mitsukawa, N., Okumura, S., Shirano, Y., Sato, S., Kato, T., Harashima, S., Shibata, D. 1997.** Overexpression of an *Arabidopsis thaliana* high-affinity phosphate transporter gene in tobacco cultured cells enhances cell growth under phosphate-limited conditions. *Proc. Natl. Acad. Sci.* 94, 7098-7102.
- Mora, M., Alfaro, M., Jarvis, S., Demanet, R., Cartes, P. 2006.** Soil aluminium availability in Andisols of southern Chile and its effect on forage production and animal metabolism. *Soil Use Manage.* 22, 95-101.
- Mora, M., Rosas, A., Ribera, A., Rengel, R. 2009.** Differential tolerance to Mn toxicity in perennial ryegrass genotypes: involvement of antioxidative enzymes and root exudation of carboxylates. *Plant Soil* 320, 79-89.
- Moroni, J., Scott, B., Wratten, N. 2003.** Differential tolerance of high manganese among rapeseed genotypes. *Plant Soil* 253, 507-519.
- Mukhopadhyay M., Sharma, A. 1991.** Manganese in cell metabolism of higher plants *Bot. Rev.* 57, 117-149.
- Nable, R., Houtz, R. Cheniae, G. 1988.** Early inhibition of photosynthesis during development of Mn toxicity in tobacco. *Plant Physiol.* 86, 1136-1142.
- Najeeb, U., Xu, L., Shafaqat, A., Jilani, G., Gong, H.J., Shen, W.Q., Zhou, W.J. 2009.** Citric acid enhances the phytoextraction of manganese and plant growth by alleviating the ultrastructural damages in *Juncus effusus* L. *J. Hazard. Mater.* 170, 1156-1163.
- Ness, P.J., Woolhouse, H.W. 1980.** RNA synthesis in *Phaseolus* chloroplasts. 1. Ribonucleic acid synthesis and senescing leaves. *J. Exp. Bot.* 31, 223-233.
- Neumann, G., Romheld, V. 2001.** The release of root exudates as affected by the plants physiological status. In: R. Pinto, Z. Varanini, P. Nannipieri (eds). *The rhizosphere: Biochemistry and organic substances at the soil-plant interface*, Marcel Dekker. New York, pp. 41-93.
- Page, V., Feller, U. 2005.** Selective transport of zinc, manganese, nickel, cobalt and cadmium in the root system and transfer to the leaves in young wheat plants. *Ann. Bot.* 96, 425-434.
- Page, V., Weisskopf, L., Feller, U. 2006.** Heavy metals in white lupin: uptake, root-to-shoot transfer and redistribution within the plant. *New Phytol.* 171, 329-341.
- Panda, S., Mishra, A.K., Biswal, U.C. 1986.** Manganese-induced modification of membrane lipid peroxidation during aging of isolated wheat chloroplasts. *Photobiochem. Photobiophys* 13, 53-61.
- Papadakis, I., Giannakoula, A., Therios, I., Bosabalidis, A., Moustakas, M., Nastou, A. 2007.** Mn-induced changes in leaf structure and chloroplast ultrastructure of *Citrus volkameriana* (L.) plants. *J. Plant Physiol.* 164, 100-103.
- Pendias, K., Pendias, H. 1992.** Trace elements in soils and plants. USA, CRR Press, 365 p.
- Pfeffer, P.E., Tu, S., Gerasimowicz, W.V., Cavanaugh, J.R. 1986.** In vivo <sup>31</sup>P NMR studies of corn root tissue and its uptake of toxic metals. *Plant Physiol.* 80, 77-84.
- Pittman, J. 2005.** Managing the manganese: molecular mechanisms of manganese transport and homeostasis. *New Phytol.* 167, 733-742.
- Polle, A. 2001.** Dissecting the superoxide dismutase-ascorbate-glutathione-pathway in chloroplasts by metabolic modeling. Computer simulations as a step towards flux analysis. *Plant Physiol.* 126, 445-462.
- Porter, G., Bajita-Locke, J., Hue, N., Strand, S. 2004.** Manganese solubility and phytotoxicity affected by soil moisture, oxygen levels, and green manure additions. *Comm. Soil Sci. Plant. Anal.* 35, 99-116.
- Reddi, A.R., Jensen, L.T., Culotta, V.C. 2009.** Manganese homeostasis in *Saccharomyces cerevisiae*. *Chem. Rev.* 109, 4722-4732.
- Rengel, Z. 2000.** Uptake and transport of manganese in plants. In: A. Sigel, H. Sigel (eds.). *Metal Ions in Biological Systems*. New York, Marcel Dekker, pp. 57-87.
- Rengel, Z. 2001.** Xylem and phloem transport of micronutrients. In: W. J. Horst et al.(eds). *Plant nutrition – Food security and sustainability of agro-ecosystems*. Kluwer Academic Publishers, Netherlands, pp. 628-629.

- Rengel, Z., Marschner, P. 2005.** Nutrient availability and management in the rhizosphere: exploiting genotypic differences. *New Phytol.* 168, 305–312.
- Roberts, L., Pierson, A., Panaviene, Z., Walker, E. 2004.** Yellow Stripe 1. Expanded roles for the maize iron-phytosiderophore transporter. *Plant Physiol.* 135, 112-120.
- Rosas, A., Rengel, Z., Mora, M. 2007.** Manganese supply and pH influence growth, carboxylate exudation and peroxidase activity of ryegrass and white clover. *J. Plant Nutr.* 30, 253-270.
- Ryan, P., Delhaize, E., Jones, D. 2001.** Function and mechanism of organic anion exudation from plant roots. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52, 527-560.
- Sarkar, D., Pandey, S., Sud, K., Chanemougasoundharam, A. 2004.** In vitro characterization of manganese toxicity in relation to phosphorus nutrition in potato (*Solanum tuberosum* L.). *Plant Sci.* 167, 977-986.
- Shi, Q., Bao, Z., Zhu, Z., He, Y., Qian, Q., Yu, J. 2005.** Silicon-mediated alleviation of Mn toxicity in *Cucumis sativus* in relation to activities of superoxide dismutase and ascorbate peroxidase. *Phytochemistry* 66, 1551-1559.
- Shi, Q., Zhu, Z., Xu, M., Qian, Q., Yu, J. 2006.** Effect of excess manganese on the antioxidant system in *Cucumis sativus* L. under two light intensities. *Environ. Exp. Bot.* 58, 197-205.
- Shigaki, T., Pittman, J., Hirschi, K. 2003.** Manganese specificity determinants in *Arabidopsis* metal/H<sup>+</sup> antiporter CAX2. *J. Biol. Chem.* 278, 6610-6617.
- Sparks, D. 1995.** *Environmental Soil Chemistry.* San Diego, Academic Press, 352 p.
- Sparrow, L., Uren, N. 1987.** Oxidation and reduction of Mn in acidic soils: effect of temperature and soil pH. *Soil Biol. Biochem.* 19, 143-148.
- Takahashi, M., Sugiura, M. 2001.** Strategies for uptake of a soil micronutrient, manganese, by plant roots. *RIKEN Review* 35, 76-77.
- Tsuji, M., Mori, H., Yamamoto, T., Tanaka, S., Kang, Y., Sakurai, K., Iwasaki, K. 2006.** Manganese toxicity of melon plants growing on an isolated soil bed after steam sterilization. *J. Soil Sci. Plant Nutr.* 77, 257-263.
- Thomine, S., Lelievre, F., Debarbieux, E., Schroeder, J., Barbier-Brygoo, H. 2003.** AtNRAMP3, a multispecific vacuolar metal transporter involved in plant responses to iron deficiency. *Plant J.* 34, 685-695.
- Wang, Y., Wisniewski, M., Meilan, R., Uratsu, S., Cui, M., Dandekar, A., Fuchigami, L. 2007.** Ectopic expression of Mn-SOD in *Lycopersicon esculentum* leads to enhanced tolerance to salt and oxidative stress. *J. Applied Hortic.* 9, 3-8.
- Wang, Y., Ying, Y., Chen, J., Wang, X. 2004.** Transgenic Arabidopsis overexpressing Mn-SOD enhanced salt-tolerance. *Plant Sci.* 167, 671-677.
- Watmough, S., Eimer, M., Dillon, P. 2007.** Manganese cycling in central Ontario forests: Response to soil acidification. *Appl. Geochem.* 22, 1241-1247.
- Wissemeier, A., Horst, W. 1991.** Simplified methods for screening cowpea cultivars for manganese leaf-tissue tolerance. *Crop Sci.* 31, 435-439.
- Wissemeier A., Horst, W. 1992.** Effect of light intensity on manganese toxicity symptoms and callose formation in cowpea (*Vigna unguiculata* (L.) Walp.). *Plant Sci.* 143, 299-309.
- Wu, Z., Liang, F., Hong, B., Young, J.C., Sussman, M.R., Harper, J.F., Sze, H. 2002.** An endoplasmic reticulum-bound Ca<sup>2+</sup>/Mn<sup>2+</sup> pump, ECA1, supports plant growth and confers tolerance to Mn<sup>2+</sup> stress. *Plant Physiol.* 130, 128–137.
- Yang, X.E., Baligar, V.C., Foster, J.C., Martens, D.C. 1997.** Accumulation and transport of nickel in relation to organic acids in ryegrass and maize grown with different nickel levels. *Plant Soil* 196: 271–276.
- Zouni, A., Witt, H., Kern, J., Fromme, P., Kraub, N., Saenger, W., Orth, P. 2001.** Crystal structure of photosystem II from *Synechococcus elongatus* at 3.8 Å resolution. *Nature* 409, 739-743.

RESEARCH PAPER

# Excess manganese differentially inhibits photosystem I versus II in *Arabidopsis thaliana*

R. Millaleo<sup>1</sup>, M. Reyes-Díaz<sup>2,3</sup>, M. Alberdi<sup>2,3,\*</sup>, A. G. Ivanov<sup>4</sup>, M. Krol<sup>4</sup> and N. P. A. Hüner<sup>4</sup>

<sup>1</sup> Programa de Doctorado en Ciencias de Recursos Naturales, Universidad de La Frontera, Temuco, Chile

<sup>2</sup> Departamento de Ciencias Químicas y Recursos Naturales; Facultad de Ingeniería, Ciencias y Administración, Universidad de La Frontera, Temuco, Chile

<sup>3</sup> Center of Plant, Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN-UFRO), Universidad de La Frontera, Temuco, Chile

<sup>4</sup> Department of Biology and The Biotron Centre for Experimental Climate Change Research, Western University, London, Ontario, N6A 5B7, Canada

\* To whom correspondence should be addressed. E-mail: [malberdi@ufro.cl](mailto:malberdi@ufro.cl)

Received 2 August 2012; Revised 18 October 2012; Accepted 30 October 2012

## Abstract

The effects of exposure to increasing manganese concentrations (50–1500  $\mu\text{M}$ ) from the start of the experiment on the functional performance of photosystem II (PSII) and photosystem I (PSI) and photosynthetic apparatus composition of *Arabidopsis thaliana* were compared. In agreement with earlier studies, excess Mn caused minimal changes in the PSII photochemical efficiency measured as  $F_v/F_m$ , although the characteristic peak temperature of the  $S_{2/3}Q_B^-$  charge recombinations was shifted to lower temperatures at the highest Mn concentration. SDS-PAGE and immunoblot analyses also did not exhibit any significant change in the relative abundance of PSII-associated polypeptides: PSII reaction centre protein D1, Lhcb1 (major light-harvesting protein of LHCII complex), and PsbO (OEC33, a 33 kDa protein of the oxygen-evolving complex). In addition, the abundance of Rubisco also did not change with Mn treatments. However, plants grown under excess Mn exhibited increased susceptibility to PSII photoinhibition. In contrast, *in vivo* measurements of the redox transients of PSI reaction centre (P700) showed a considerable gradual decrease in the extent of P700 photooxidation (P700<sup>+</sup>) under increased Mn concentrations compared to control. This was accompanied by a slower rate of P700<sup>+</sup> re-reduction indicating a downregulation of the PSI-dependent cyclic electron flow. The abundance of PSI reaction centre polypeptides (PsaA and PsaB) in plants under the highest Mn concentration was also significantly lower compared to the control. The results demonstrate for the first time that PSI is the major target of Mn toxicity within the photosynthetic apparatus of *Arabidopsis* plants. The possible involvement mechanisms of Mn toxicity targeting specifically PSI are discussed.

**Key words:** Chlorophyll fluorescence, Mn toxicity, photosystem I, PSI-associated proteins, PSII-associated proteins, redox state of P700.

## Introduction

Manganese (Mn) is one of the most abundant metals in the Earth's crust and although it is an important essential micro-nutrient for all photosynthetic organisms can be also toxic when it is present in excess (Mukhopadhyay and Sharma,

1991; Marschner, 1995). Mn is considered the second most phytotoxic element, after aluminium (Al), affecting negatively the physiological and biochemical properties of plant species (Foy *et al.*, 1978, Foy, 1984; Millaleo *et al.*, 2010). An

excess of this metal occurs in acid soils with low pH (<5.5) and/or under reducing conditions (Marschner, 1995; Schaaf *et al.*, 2002), where  $Mn^{2+}$  is the predominant solution species and available ion to plant cells (Bradl, 2004). Thus, a Mn excess results in a sharp decrease in shoot height, biomass accumulation, and total leaf area of a woody species (*Populus cathayana*, Lei *et al.*, 2007), a reduction of the dry weights (DW) of both shoots and roots in ryegrass cultivars (*Lolium perenne*, Mora *et al.*, 2009), and *Trifolium repens* (Rosas *et al.*, 2007). Furthermore, excess Mn can result in oxidative stress as indicated by the accumulation of  $H_2O_2$  (Demirevska-Kepova *et al.*, 2004; Lei *et al.*, 2007), high levels of apoplastic  $H_2O_2$ -consuming peroxidases (Fecht-Christoffers *et al.*, 2003), and high level of lipid peroxidation (Mora *et al.*, 2009). Mn stress induced an enhancement of antioxidant enzyme activity in leaves of legumes (González *et al.*, 1998), and it was also demonstrated in perennial ryegrass (Mora *et al.*, 2009) and woody species (Lei *et al.*, 2007). More recently, a proteomic and transcriptomic studies have demonstrated that chloroplastic proteins important for  $CO_2$  fixation and photosynthesis were of lower abundance upon Mn stress of cowpea (Führs *et al.*, 2008).

Mn has an important role in both the structure and functions of the photosynthetic apparatus (Mukhopadhyay and Sharma, 1991). Mn is a constitutive element associated with the oxygen-evolving complex of photosystem II (PSII), an important multiprotein pigment complex embedded in the thylakoid membranes (Hankamer *et al.*, 1997; Enami *et al.*, 2008). Therefore, the Mn cluster, together with other ions and extrinsic proteins that constitute the oxygen-evolving complex, is required to oxidize water and reduce P680, the reaction centre of PSII (Kern and Renger, 2007; Ferreira *et al.*, 2004; Rutherford and Boussac, 2004). In conjunction with photosystem I (PSI) and linear electron transport, these reducing equivalents (electrons) are used primarily in the conversion of  $CO_2$  into carbohydrate (Ferreira *et al.*, 2004). In addition, Mn is indispensable as a cofactor for various enzymes involved in redox reactions such as Mn-superoxide dismutase, an essential enzyme involved in protection against oxidative stress in plants (Burnell, 1988; Bowler *et al.*, 1994).

A number of studies have suggested that chloroplasts and photosynthesis are the major targets of Mn toxicity. Indeed, increased amounts of Mn have been reported for chloroplasts isolated from Mn-stressed common bean (González and Lynch, 1999) and rice leaves (Lidon *et al.*, 2004). Distinctive ultrastructural changes showing swelling of granal and stromal thylakoids have been also observed in the chloroplasts of *Citrus volkameriana* (Papadakis *et al.*, 2007) and maize plants (Doncheva *et al.*, 2009) under Mn excess. It has been demonstrated that high Mn accumulation is associated with inhibition of the net photosynthesis and carboxylation efficiency in various plant species. The decline of photosynthesis is considered as one of the major mechanisms constituting the toxic effects of excess Mn and is proposed as an early indicator for Mn toxicity in tobacco (Nable *et al.*, 1988), rice (Lidon *et al.*, 2004) and wheat (Macfie and Taylor, 1992). Reduced  $CO_2$  assimilation induced by excess Mn was also reported for common bean (González and Lynch, 1997), deciduous broad

leaved trees (Kitao *et al.*, 1997a), and seedlings of *Citrus grandis* (Li *et al.*, 2010). Interestingly, the maximum photochemical efficiency of PSII ( $F_v/F_m$ ) was not substantially affected by Mn accumulation in various plant species over a wide range of leaf Mn concentrations (Nable *et al.*, 1988; Kitao *et al.*, 1997b; Subrahmanyam and Rathore, 2000; Hajiboland and Hasani, 2007; Doncheva *et al.*, 2009).

The reduction in photosynthesis by excess leaf Mn has been generally attributed to modification of ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco; Ohki, 1984; Houtz *et al.*, 1988; McDaniel and Toman, 1994; Kitao *et al.*, 1997b). It has been demonstrated that a high level of Mn affects primarily the activity rather than the amount of Rubisco (Houtz *et al.*, 1988; Chatterjee *et al.*, 1994) and the presence of excess Mn induces enhanced oxygenase activity (Jordan and Ogren, 1983). In addition, the decline of photosynthesis under Mn stress conditions was also ascribed to peroxidative impairment of photosynthetic enzyme activities caused by polyphenol oxidation products (Vaughn and Duke, 1984; Panda *et al.*, 1987).

In spite of these studies, the mechanisms of Mn toxicity causing a decrease in  $CO_2$  assimilation are still not well understood. In addition, very limited information concerning the toxic effect(s) of excess Mn on the polypeptide composition of both PSII and, especially, PSI is available. Therefore, the objective of this study was to evaluate the role of specific Mn-induced changes in the structure and function of PSII and PSI, which could help to understand the mechanisms by which Mn excess may cause a decrease of  $CO_2$  assimilation in *Arabidopsis thaliana*.

## Materials and methods

### Plant material and growth conditions

Seeds of *A. thaliana* (wild type Columbia) were germinated in a substrate mix (82.5% sphagnum peat moss, 12.5% perlite, 5% vermiculite-Pro-Mix, Premier Tech Horticulture) in controlled environment growth cabinets (model GCW15, Environmental Growth Chambers, Chagrin Falls, OH, USA) with a photosynthetical active radiation (PAR) of 250  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , 20/20 °C day/night temperatures, 50% relative humidity, and 8/16 light/dark cycle to prevent flowering. Water was supplied every 5 days. After 15 days, seedlings were transplanted separately in pots with vermiculite and placed in trays. Each tray containing seven pots (one plant per pot) were supplied with Hoagland nutrient solution for 2 weeks before applying the Mn treatments.

### Manganese treatments

Manganese treatments included the final concentrations: 18 (control), 50, 500, 1000, and 1500  $\mu\text{M}$  Mn according to Delhaize *et al.* (2007). Manganese was applied as  $MnCl_2 \cdot 4H_2O$ . Control plants exposed to 18  $\mu\text{M}$  Mn as the optimal dose for Mn for *Arabidopsis* (Cailliatte *et al.*, 2010). The five Mn treatments were grown in five labelled trays, with 500 ml of Hoagland's solution. The trays were maintained in controlled environment growth chambers under the same conditions described above. The pH was adjusted to 5.3 with diluted HCl daily and nutrient solution was changed every 5 days. Plants were subjected to these treatments for 21 days before harvest.

### Plant growth measurements

Prior to beginning the Mn treatments, three plant samples were dried in a forced-air oven (70 °C, 48 h) and weighed to determine dry weight

(W1) at day 0. Similarly, at the end of the experiment, plants were harvested and collected for dry weight measurements (W2). These data were used to determine mean relative growth rate (RGR) according to [Fernando et al. \(2009\)](#):

$$RGR = \frac{(\ln W2 - \ln W1)}{t2 - t1}$$

as g DW d<sup>-1</sup>.

#### Manganese concentration

For Mn chemical analysis, samples of shoot and roots were dry ashed in a muffle furnace at 500 °C for 8 h and digested with 2 M HCl. Manganese was extracted as described by [Sadzawka et al. \(2004\)](#), and the Mn concentration was determined using a simultaneous multi-element atomic absorption spectrophotometer (model 969, Unicam, Cambridge, UK).

#### Thylakoid preparation, SDS-PAGE, and immunoblotting

Thylakoid membranes for SDS-PAGE were isolated as described earlier ([Krol et al., 1999](#)). Leaf material was ground in cold isolation buffer (50 mM Tricine, 0.4 M sorbitol, 10 mM NaCl, 5 mM MgCl<sub>2</sub>, pH 7.8) in a mortar on ice, filtered through two layers of miracloth (typical pore size 22–25 µm; Calbiochem), and centrifuged for 15 min (10,000 g). The supernatant was removed and the pellet was resuspended in cold isolation buffer. Total chlorophyll concentration was measured in 90% (v/v) acetone ([Arnon, 1949](#)). For immunodetection of Rubisco, total leaf proteins were extracted as described in [Rosso et al. \(2009\)](#). Protein content was measured using a BCA protein assay kit (Pierce) by following the absorbance at 562 nm using a spectrophotometer (DU-640, Beckman Coulter). Proteins were separated by SDS-PAGE according to [Laemmli \(1970\)](#), using 15% (w/v) polyacrylamide gel in the presence of 6 M urea in the separating gel. Chloroplast thylakoids were solubilized with SDS (SDS:chlorophyll 20:1) and 15 µg chlorophyll was loaded per lane. All samples for separation of total proteins were loaded on an equal protein basis of 20 µg protein per lane ([Rosso et al. 2009](#)). Immunoblotting was performed by electrophoretically transferring the proteins from SDS-PAGE gel to nitrocellulose membrane (Bio-Rad) according to the method of [Towbin et al. \(1979\)](#). Proteins were probed with antibodies (AgriSera AB, Vanas, Sweden) raised against the reaction centre polypeptides of PSI: PsaA, PsaB (1:2000), the major light-harvesting protein of PSII complex (LHCII) Lhcb1 protein (1:5000), the PSII oxygen-evolving complex extrinsic protein PsbO (33 kDa, 1:2000), the PSII reaction centre protein: D1 and Rubisco (1:5000). As secondary antibodies, goat anti-rabbit IgG conjugated with horseradish peroxidase (Sigma-Aldrich) were used. Polypeptides were detected using enhanced chemiluminescence detection kit (Amersham Biosciences) and visualized by exposing the membrane to X-ray film. Densitometric scanning and analysis of X-ray films from each replicate immunoblot was performed with a Hewlett Packard ScanJet 4200C desktop scanner and ImageJ 1.41o densitometry software (Wayne Rosband, National Institute of Health, USA, <http://rsbweb.nih.gov/ij>).

#### Measurement of the redox state of P<sub>700</sub>

The redox state of P<sub>700</sub> was determined *in vivo*, in dark-adapted (20 min) *Arabidopsis* leaves under growth temperature and ambient O<sub>2</sub> and CO<sub>2</sub> conditions using a PAM-101 modulated fluorometer equipped with a dual-wavelength emitter-detector ED-P700DW unit and PAM-102 units ([Klughammer and Schreiber, 1991](#)) as described in detail by [Ivanov et al. \(1998\)](#). Far-red light (λ<sub>max</sub>=715 nm, 10 W m<sup>-2</sup>, Schott filter RG 715) was provided by an FL-101 light source. The redox state of P700 was evaluated as the absorbance change around 820 nm (ΔA<sub>820–860</sub>) in a custom-designed cuvette. Multiple turnover (MT, 50 ms) and single turnover (ST, half peak 14 µs) saturating flashes were applied with XMT-103 and XST-103 (Walz) power/control units, respectively. The relative functional pool size of intersystem electrons on a P<sub>700</sub> reaction

centre basis was calculated as the complementary area between the oxidation curve of P700 after either ST or MT pulse excitation (ST and MT areas) and the stationary level of P700 under far-red excitation ([Asada et al., 1993](#); [Ivanov et al., 1998](#)). Capacity of PSI cyclic electron (e<sup>-</sup>) flow was determined as the half time for the dark decay of the P700 signal ([Ivanov et al., 1998](#)).

#### Modulated chlorophyll fluorescence measurements

Modulated imaging fluorometer (IMAGING-PAM, Heinz Walz, Efeltrich, Germany) was used for capturing the chlorophyll fluorescence images and estimation of the maximal photochemical efficiency of PSII [F<sub>v</sub>/F<sub>m</sub> = (F<sub>m</sub> - F<sub>o</sub>)/F<sub>m</sub>], effective photochemical efficiency of PSII (Φ<sub>PSII</sub>), photochemical (qP), and non-photochemical (NPQ) fluorescence quenching parameters using the nomenclature of [van Kooten and Snel \(1990\)](#) as described earlier ([Ivanov et al., 2006a](#)). Control and Mn-treated *Arabidopsis* plants were dark adapted (20 min) and all chlorophyll fluorescence measurements were performed *in vivo* at room temperature. Fluorescence images were captured by a CCD camera (IMAG-K, Allied Vision Technologies) featuring a 640 × 480 pixel CCD chip size and CCTV camera lens (Cosmicar/Pentax F1.2, f = 12 mm). Light emitting diode ring array (IMAG-L) consisting of 96 blue LEDs (470 nm) provided standard modulated excitation intensity of 0.5 µmol quanta m<sup>-2</sup> s<sup>-1</sup> (modulation frequency 1–8 Hz) for measuring the basal (F<sub>o</sub>) chlorophyll fluorescence and a saturation pulse of 2400 µmol quanta m<sup>-2</sup> s<sup>-1</sup> PAR for measuring the maximal chlorophyll fluorescence (F<sub>m</sub>). The maximal photochemical efficiency of PSII (F<sub>v</sub>/F<sub>m</sub>) was determined as (F<sub>m</sub> - F<sub>o</sub>)/F<sub>m</sub>. The effective photochemical efficiency of PSII (Φ<sub>PSII</sub>) was calculated from the expression (F'<sub>m</sub> - F<sub>s</sub>)/F'<sub>m</sub> ([Genty et al., 1989](#)), photochemical quenching (qP) was calculated as (F'<sub>m</sub> - F<sub>s</sub>)/(F'<sub>m</sub> - F'<sub>o</sub>) ([Schreiber et al., 1994](#)), electron transport rate (ETR) was calculated as PAR × 0.5Φ<sub>PSII</sub> × 0.84 ([Genty et al., 1989](#)), and regulated non-photochemical quenching (Φ<sub>NPQ</sub>) and constitutive photochemical quenching (Φ<sub>NO</sub>) was determined according to [Kramer et al. \(2004\)](#). All measurements were performed at 0, 5, 10, 15, and 21 days in plants subjected to the different Mn treatments.

At the end of experiments (21 days), leaves of *Arabidopsis* plants growing under the different Mn treatments and 250 µmol m<sup>-2</sup> s<sup>-1</sup> PAR (normal light) were cut and exposed to high light (1000 µmol s<sup>-1</sup> m<sup>-2</sup>, PAR) at 1, 2, and 4 hours. During each time of exposition, F<sub>v</sub>/F<sub>m</sub> was determined.

#### Thermoluminescence measurements

Thermoluminescence (TL) measurements of control and Mn-treated *Arabidopsis* leaves were performed on a personal-computer-based TL data acquisition and analysis system as described earlier ([Ivanov et al., 2001, 2006b](#)). A photomultiplier tube (Hamamatsu R943-02, Hamamatsu Photonics, Shizuoka-ken, Japan) equipped with a photomultiplier power supply (model PS-302, EG&G Electro Optics), and a preamplifier (model C1556-03) was used as a radiation measuring set. The heating rate was 0.6 °C s<sup>-1</sup>. For identifying the S<sub>2/3</sub>Q<sub>B</sub><sup>-</sup> charge recombination peaks, dark-adapted leaf discs were subjected to two consecutive saturating microsecond flashes of white light (1.5 µs peak width at 50% of maximum) applied by a xenon-discharge flash lamp (XST103, Heinz Walz). Dark-adapted leaves (20 minutes at 20 °C) were cooled to 2 °C prior to exposing to the flashes. The nomenclature of [Sane et al. \(2012\)](#) was used for characterization of the TL glow peaks.


#### Experimental design and statistical analysis

The experimental design is a randomized split design, with 1 species × 5 Mn treatments × 7 replicates for the physiological determinations and 1 species × 5 Mn treatments × 3 replicates × 4 times (0, 1, 2, and 4 hours) for the photoinhibition measurements. Data correspond to the means of replicates for each determination as indicated above. Data were tested in their normality and equal variance by the Shapiro–Wilk test and then data were analysed by a one-way ANOVA. Significant differences between means were established by using the multiple comparisons test

of Tukey's ( $P < 0.05$ ). All analyses were performed with Sigma Stat 2.0 software (SPSS), where differences between the values were considered significant at  $P \leq 0.05$ .

## Results

A statistically significant decrease in total biomass of leaves treated with 500 and 1500  $\mu\text{M}$  Mn concentrations was found compared to the control plants (18  $\mu\text{M}$  Mn), although a decrease of the total leaf biomass was found at all Mn concentrations used. In contrast, root biomass exhibited significant decrease in all treatments ( $P < 0.05$ , Fig. 1). It should be noted, however, that while the decrease of leaf biomass was only 25% lower at the highest Mn concentration used (1500  $\mu\text{M}$ ), the biomass of roots was more affected and demonstrated a 2.5-fold decrease even at the lowest Mn concentration tested (50  $\mu\text{M}$ ). This is consistent with the RGR data, where a considerably higher reduction of root RGR was observed compared to leaves across the Mn treatments relative to control plants ( $P < 0.05$ , Table 1). In addition, visual symptoms of Mn toxicity (chlorosis) in leaves were observed predominantly in the highest Mn treatment (data not shown).




**Fig. 1.** Total biomass (leaves and roots) of *Arabidopsis thaliana* plants subjected to Mn treatments at the end of experiment (21 days). Different lower-case letters indicate significant differences between the Mn treatments ( $P \leq 0.05$ ).

**Table 1.** Relative growth rates of *Arabidopsis thaliana* (leaves and roots) growing at increased Mn treatments. Plants were grown under acidic conditions (pH 5.3) for 21 d. Different lower-case letters indicate statistically significant differences between the Mn treatments ( $P < 0.05$ ).

Mn treatment ( $\mu\text{M}$ )	Relative growth rate (g DW d <sup>-1</sup> )	
	Leaves	Roots
Control	0.069 ± 0.001 <sup>a</sup>	0.206 ± 0.006 <sup>a</sup>
50	0.068 ± 0.003 <sup>a</sup>	0.071 ± 0.005 <sup>b</sup>
500	0.046 ± 0.006 <sup>b</sup>	0.053 ± 0.006 <sup>b</sup>
1000	0.059 ± 0.003 <sup>a,b</sup>	0.064 ± 0.007 <sup>b</sup>
1500	0.040 ± 0.004 <sup>b</sup>	0.057 ± 0.004 <sup>b</sup>

Analysis of the total Mn amount in plants exposed to increasing Mn concentrations have demonstrated a gradual increase of Mn in leaves and roots (Fig. 2). However, while Mn concentrations of roots reached values up to ~8100 mg kg<sup>-1</sup> Mn at the highest Mn treatment, which represents a 40-fold increase of Mn, the increased accumulation of Mn in leaves was much lower (15-fold) compared to roots (Fig. 2). Thus, the differential effects of Mn treatments on the total



**Fig. 2.** Mn concentration in leaves and roots of *Arabidopsis thaliana* plants subjected to Mn treatments at the end of experiment (21 days). Different lower-case letters indicate statistically significant differences between the Mn treatments ( $P \leq 0.05$ ).


**Table 2.** Maximum photochemical efficiency of PSII ( $F_v/F_m$ ) in leaves of *Arabidopsis thaliana* measured at 0, 5, 10, 15 and 21 days subjected to increasing Mn treatments. Results are means  $\pm$  SE of five replicates of two plants each. Different lower-case letters indicate significant differences between the Mn treatments at the same time, while different upper-case letters indicate significant differences between the times at a same Mn treatment ( $P < 0.05$ ).

Mn treatment ( $\mu\text{M}$ )	$F_v/F_m$				
	0 days	5 days	10 days	15 days	21 days
Control	0.803 <sup>abA</sup>	0.803 <sup>abA</sup>	0.803 <sup>abA</sup>	0.803 <sup>abA</sup>	0.803 <sup>abA</sup>
50	0.806 <sup>abA</sup>	0.799 <sup>abA</sup>	0.800 <sup>abA</sup>	0.800 <sup>abA</sup>	0.800 <sup>abA</sup>
500	0.803 <sup>abA</sup>	0.801 <sup>abA</sup>	0.804 <sup>abA</sup>	0.787 <sup>ba</sup>	0.791 <sup>abA</sup>
1000	0.797 <sup>abA</sup>	0.805 <sup>abA</sup>	0.800 <sup>abA</sup>	0.778 <sup>bbB</sup>	0.773 <sup>bbB</sup>
1500	0.802 <sup>abA</sup>	0.798 <sup>abA</sup>	0.775 <sup>bbB</sup>	0.769 <sup>bbBC</sup>	0.756 <sup>ccC</sup>

biomass and RGR in leaves and roots could be explained by the higher accumulation of Mn in roots.

The effects of exposure to increasing Mn concentrations on the maximum photochemical efficiency of PSII measured as  $F_v/F_m$  in *Arabidopsis* leaves are presented in Table 2. A time-course measurement did not reveal any statistically significant changes of  $F_v/F_m$  values at the lower Mn doses (50 and 500  $\mu\text{M}$ ) for the entire period of treatment compared to control plants (18  $\mu\text{M}$  Mn). At higher Mn concentrations (1000 and 1500  $\mu\text{M}$ ), a small reduction (6%) of  $F_v/F_m$  was observed only at a later stage (days 15 and 21) of treatment, the statistically significant ( $P < 0.05$ ) differences being registered at the harvest time (21 days, Table 2). Moreover, minimal Mn-induced changes were detected in the light response curves of photochemical quenching (qP), effective photochemical efficiency of PSII ( $\Phi_{\text{PSII}}$ ), and PSII electron transport rates (ETR) even after 21 days of Mn treatments with the highest dose tested (1500  $\mu\text{M}$ , data not shown).

In addition, TL measurements were used as an alternative approach for assessing the effects of Mn excess on the photosynthetic PSII-associated electron transfer reactions especially at its reducing side (Vass and Govindjee, 1996; Sane et al., 2012). Since most of the photosynthetic TL components have been assigned to arise from the reversal of light-driven charge separation in PSII, TL properties of photosynthetic apparatus provide information on the activation energies associated with the back reactions of electron acceptors ( $Q_A$  and  $Q_B$ ) with the electron donors ( $S_2$  and  $S_3$ ) of PSII (Vass and Govindjee, 1996; Sane, 2004; Sane et al., 2012). The temperature maxima ( $T_M$ ) of the TL peaks related to the recombination of these charge pairs reflect the activation energies and hence a measure of the redox potentials of the participating oxidized and reduced donors (de Vault and Govindjee, 1990). Typical TL glow curves representing  $S_{2/3}Q_B^-$  charge recombinations of control and Mn-treated *Arabidopsis* plants obtained following excitation with two consecutive saturating flashes are shown in Fig. 3. The experimental data summarized in Table 3 indicate that treatment with a Mn dose of 500  $\mu\text{M}$  did not exhibit significant differences of the TL



**Fig. 3.** Thermoluminescence (TL) glow curves of  $S_{2/3}Q_B^-$  charge recombinations in control (solid lines) and Mn-treated (dashed lines, 500  $\mu\text{M}$   $\text{Mn}^{2+}$ ; dotted lines, 1500  $\mu\text{M}$   $\text{Mn}^{2+}$ ) *Arabidopsis* leaves after illumination with two single turnover flashes. TL glow curves were recorded immediately after illumination. The presented glow curves are averages from four independent measurements.

peak position, while treatment with the highest concentration (1500  $\mu\text{M}$ ) induced a low temperature shift of the  $T_M$  to 25.8  $^{\circ}\text{C}$  compared to control plants. Besides this, the amplitudes and the integrated areas of the TL peaks representing the  $S_{2/3}Q_B^-$  charge recombination used for assessing the PSII photochemistry was not affected at 500  $\mu\text{M}$  Mn excess, but the overall TL yield was significantly reduced (45%) in plants treated with 1500  $\mu\text{M}$  Mn compared to controls (Table 3).

To further test whether this small but significant Mn-induced effect on PSII photochemistry had physiological implications under additional stress conditions, *Arabidopsis*

**Table 3.** Characteristic thermoluminescence peak emission temperatures ( $T_M$ ) and the overall TL emission area (A) of  $S_{2/3}Q_B^-$  glow peaks of control and Mn-treated (21 days) *Arabidopsis* plants. The samples (leaf disks) were dark adapted for 30 min then cooled to 2  $^{\circ}\text{C}$  and subsequently illuminated with two single turnover flashes of white light. The peak areas are presented as a percentage of the total thermoluminescence light emission in control leaves. Values are mean  $\pm$  SE calculated from four independent experiments.

Mn treatment ( $\mu\text{M}$ )	$T_M$ ( $^{\circ}\text{C}$ )	A (%)
Control	28.7 $\pm$ 1.1	100.0
500	29.1 $\pm$ 1.3	106.2 $\pm$ 7.6
1500	25.8 $\pm$ 0.9	54.5 $\pm$ 7.9


**Table 4.** Effect of high light treatments (photon flux density 1000  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for 1, 2, and 4 hours) on the maximal photochemical efficiency of PSII measured as  $F_v/F_m$  in control *Arabidopsis thaliana* leaves and plants exposed for 21 days to different Mn doses. Results are mean  $\pm$  SE of five repetitions in three leaves of two plants each. Time 0 was measured in plants subjected to 250  $\mu\text{mol m}^{-2} \text{s}^{-1}$  photon flux density. Different upper-case letters indicate significant differences between the times of exposure at the same Mn treatment ( $P < 0.05$ ).

Irradiance exposition time (h)	$F_v/F_m$		
	Control	500 $\mu\text{M Mn}$	1500 $\mu\text{M Mn}$
0	0.815 $\pm$ 0.003 <sup>A</sup>	0.812 $\pm$ 0.004 <sup>A</sup>	0.780 $\pm$ 0.014 <sup>A</sup>
1	0.761 $\pm$ 0.012 <sup>A</sup>	0.750 $\pm$ 0.002 <sup>A</sup>	0.677 $\pm$ 0.014 <sup>B</sup>
2	0.752 $\pm$ 0.007 <sup>A</sup>	0.731 $\pm$ 0.002 <sup>A</sup>	0.619 $\pm$ 0.028 <sup>B</sup>
4	0.685 $\pm$ 0.036 <sup>A</sup>	0.689 $\pm$ 0.004 <sup>A</sup>	0.388 $\pm$ 0.042 <sup>C</sup>

leaves subjected to different Mn treatments for 21 days were exposed to high light stress (Table 4). Indeed, the photoinhibitory effect on PSII, measured as a decrease in  $F_v/F_m$ , was much stronger (49%) at the highest Mn treatment (1500  $\mu\text{M}$ ) compared to control plants, and 500  $\mu\text{M}$  Mn treated plants exhibited only a 15% decrease in  $F_v/F_m$  values after 4 h of exposure to high light (Table 4).

The extent of far-red light-induced absorbance decrease at 820 nm ( $\Delta A_{820-860}$ ) of *Arabidopsis* leaves (Klughammer and Schreiber, 1991; Ivanov et al., 1998, 2006a) was used to estimate the potential functional differences of PSI and photosynthetic electron transport pathways between plants exposed to different Mn treatments. Typical traces representing *in vivo* measurements of oxidation–reduction transients of  $P_{700}$  in control and Mn-treated plants are shown in Fig. 4. The relative amount of  $P_{700}^+$ , measured as  $\Delta A_{820-860}$ , gradually decreased with increasing Mn concentrations and was 30% lower in Mn-treated plants at the highest concentration used (1500  $\mu\text{M}$ , Fig. 4 and Table 5). Concomitantly, kinetic measurements of dark re-reduction of  $P_{700}^+$  after turning off the far-red light, which is thought to reflect the extent and/or capacity for cyclic electron flow around PSI (Maxwell and Biggins, 1976; Ravenel et al., 1994), indicated significantly slower (46%) re-reduction of  $P_{700}^+$  in Mn (1500  $\mu\text{M}$ )-treated plants compared to control plants (Table 5). In addition, the apparent electron donor pool size to PSI ( $e^-/P_{700}$ ) estimated by measuring single- and multiple-turnover flash-induced  $\Delta A_{820-860}$  under steady-state oxidation of PSI by far-red light (Asada et al., 1993; Ivanov et al., 1998) demonstrated a significant decrease in Mn-treated plants (Table 5). This indicates that the pool size of electrons that can be donated to photooxidized  $P_{700}$  ( $P_{700}^+$ ) from the stroma in control plants was 37% higher compared to plants treated with the highest Mn dose (13 electrons per  $P_{700}$ , Table 4).

The major photosynthetic components within the thylakoid membranes of control and Mn-treated *Arabidopsis* plants were compared by SDS-PAGE and immunodetection to quantify their relative abundance. Immunoblot analyses did not exhibit any significant Mn-stress-induced changes



**Fig. 4.** Typical traces of *in vivo* measurements of  $P_{700}$  oxidation by far-red light (FR) in control (A, 18  $\mu\text{M Mn}$ ) and Mn-treated (B, 1500  $\mu\text{M Mn}$ ) *Arabidopsis* plants. After reaching a steady state level of  $P_{700}^+$  by FR light, single turnover (ST) and multiple turnover (MT) pulses of white light were applied. Arrows indicate application of ST, MT, and FR light sources. The measurements were performed at the growth temperature of 20  $^{\circ}\text{C}$ .

in the relative abundance of PSII-associated polypeptides, as revealed by the densitometry analysis of the immunoblot bands for D1 (the PSII reaction centre protein), Lhcb1 (major light-harvesting protein of LHCII), and PsbO (extrinsic protein of the oxygen-evolving complex (Fig. 5). The abundance of Rubisco was only marginally affected by the Mn treatments. In contrast, the abundance of reaction centre polypeptides of PSI (PsaA and PsaB) was significantly reduced in Mn-treated compared with control thylakoids (Fig. 5A). The densitometric analysis demonstrated that the relative abundance of PsaA and PsaB in Mn-treated *Arabidopsis* was only about 20 and 60%, respectively, of that observed in the control plants (Fig. 5B). Thus, the quantitative analysis of photosynthetic polypeptides clearly indicates that excess Mn has

**Table 5.** Effects of Mn treatments on far-red light-induced steady state oxidation of P700 ( $\Delta A_{820-860}$ , P700<sup>+</sup>), the relative intersystem electron donor pool size to PSI ( $e^-/P700$ ) and half times for P700<sup>+</sup> reduction ( $t_{1/2}$ ) of *Arabidopsis* leaves at 21 days of treatment. Different lower-case letters indicate significant differences between the Mn treatments ( $P < 0.05$ ). MT, multiple turnover; ST, single turnover.

Mn treatment ( $\mu\text{M}$ )	$\Delta A_{820-860}$ (P700 <sup>+</sup> )	$e^-/P700$ (MT <sub>area</sub> /ST <sub>area</sub> )	$t_{1/2}$ (s)
Control	423.2 ± 7.9 <sup>a</sup>	20.7 ± 0.9 <sup>a</sup>	0.670 ± 0.05 <sup>b</sup>
50	416.4 ± 8.3 <sup>a</sup>	15.7 ± 0.6 <sup>b</sup>	0.812 ± 0.06 <sup>a,b</sup>
500	368.4 ± 8.3 <sup>b</sup>	16.4 ± 0.6 <sup>b</sup>	0.938 ± 0.08 <sup>a,b</sup>
1000	293.4 ± 8.4 <sup>c</sup>	15.6 ± 0.5 <sup>b</sup>	0.908 ± 0.07 <sup>a,b</sup>
1500	292.9 ± 4.2 <sup>c</sup>	13.0 ± 0.3 <sup>c</sup>	0.984 ± 0.08 <sup>a</sup>


a greater effect on PSI-associated proteins rather than PSII. Consequently, the differences in the extent of P700 oxidation (P700<sup>+</sup>) and the kinetics of P700<sup>+</sup> reduction between control and Mn-treated plants are consistent with the lower levels of PSI-associated immunodetectable PsaA and PsaB protein complexes.

## Discussion

In agreement with a number of previous studies examining the effects of Mn stress on various plant species (Alam *et al.*, 2006; Lei *et al.*, 2007; Doncheva *et al.*, 2009; Mora *et al.*, 2009; Stoyanova *et al.*, 2009; Khabaz-Saberi *et al.*, 2010) *Arabidopsis* plants subjected to increasing Mn concentrations, also exhibited a reduction in dry weight of both shoots and roots at doses between 500 and 1500  $\mu\text{M}$  Mn (Fig. 1). The decline in biomass corresponded with a gradual increase of Mn concentrations in both shoot and roots of *Arabidopsis* subjected to excess Mn supply (Fig. 2). It should be noted that the decline of biomass was more pronounced in roots, where even at the lowest Mn treatment (50  $\mu\text{M}$ ) the dry weight was 2.5-fold lower compared to leaves. Similar results have been reported by Delhaize *et al.* (2007), where transporter proteins were implicated in the endogenous Mn tolerance of wild-type *Arabidopsis*. More recently, Mora *et al.* (2009) demonstrated that Mn-tolerant ryegrass cultivars accumulated higher Mn concentrations in roots than shoots, while Mn-sensitive cultivars exhibited a greater Mn translocation from roots to shoots. These results are also consistent with studies in legumes such as white clover (*T. repens* L., Rosas *et al.*, 2007). However, in two contrasting populations of *P. cathayana*, acclimated to wet and dry climate exposure to excess Mn caused an increase in Mn content of plant tissues especially in leaves and a visual symptoms of Mn toxicity (chlorosis) at high Mn concentrations (Lei *et al.*, 2007). The chlorosis observed in the present experiments (data not shown) also correspond to a decreased amounts of both Chl a and Chl b could be due to a higher Mn accumulation in leaves after exposure to Mn excess, thus suggesting a damage to the photosynthetic apparatus as reported by Demirevska-Kepova *et al.* (2004).

Interestingly, while reduced CO<sub>2</sub> assimilation induced by excess Mn has been reported in many species and is considered one of the major physiological effects of Mn toxicity (Nable *et al.*, 1988; Macfie and Taylor, 1992; González and Lynch, 1997; Kitao *et al.*, 1997a; Lidon *et al.*, 2004; Li *et al.*, 2010), the functional integrity of the photosynthetic apparatus assessed by the maximum photochemical efficiency of PSII ( $F_v/F_m$ ) did not decline as a result of exposure to excess Mn in various plant species (Nable *et al.*, 1988; Kitao *et al.*, 1997b; Subrahmanyam and Rathore, 2000; Hajiboland and Hasani, 2007; Doncheva *et al.*, 2009). However, some studies have reported a substantial decrease in  $F_v/F_m$  as a result of excess Mn treatment in *Citrus* species (Papadakis *et al.*, 2007; Li *et al.*, 2010), rice (Lidon *et al.*, 2004), Mn-sensitive maize (Doncheva *et al.*, 2009), and cucumber (Feng *et al.*, 2009). Furthermore, Kitao *et al.* (1997b) have shown that the potential maximum photochemical efficiency of PSII ( $F_v/F_m$ ) is not affected by excess Mn in white birch, although the reduction state of PSII primary electron acceptor ( $Q_A$ ) was increased at high Mn concentrations. These results clearly indicate that the excess Mn-induced decline in CO<sub>2</sub> assimilation may or may not be accompanied by changes in PSII photochemistry and this response is species dependent. The experimental data presented in this study also failed to demonstrate any substantial effects of excess Mn within a wide range of Mn concentrations on the maximum photochemical efficiency of PSII in *Arabidopsis* plants (Table 2). Assessing the relative abundance of PSII-associated proteins also showed no changes in the immunodetectable amounts of PSII reaction centre protein D1, the light-harvesting chlorophyll-protein complex of PSII (Lhcb1), and the manganese-stabilizing 33-kDa protein of the water splitting complex of PSII (PsbO) polypeptides (Fig. 5) in plants exposed to high Mn concentrations compared to control *Arabidopsis*. However, the observed low temperature shift of the  $S_{2/3}Q_B^-$  charge recombination and much lower overall TL emission implies lower redox potential of  $Q_B$  (Fig. 3 and Table 3) confirms the suggestion of altered reduction state of PSII acceptor side in Mn-stressed plants (Kitao *et al.*, 1997b). Since  $Q_A$  is in quasi-equilibrium with  $Q_B$  and the PQ pool, the present results imply that lowering the redox potential of  $Q_B$  will decrease the probability for forward electron transfer between the two quinone acceptors by shifting the redox equilibrium between  $Q_A^-Q_B$  and  $Q_AQ_B^-$  towards  $Q_A^-Q_B$  (Minagawa *et al.*, 1999; Ivanov *et al.*, 2002, 2003) in plants exposed to high Mn concentrations.

In addition to the lack of significant inhibitory effects of excess Mn on PSII photochemistry discussed above, an earlier study reported that the photochemistry of photosystem I (PSI) and the photosynthetic electron transport were not significantly affected during early development of Mn toxicity in tobacco plants (Nable *et al.*, 1988). However, a decreased Hill activity in isolated chloroplasts was found in mungbean leaves exposed to toxic Mn concentrations (Sinha *et al.*, 2002). More recently, Li *et al.* (2010) have suggested that Mn excess can effectively impair the whole photosynthetic electron transport chain, thus restricting the production of reducing equivalents and limiting the rate of CO<sub>2</sub> assimilation in *Citrus grandis* seedlings. Despite these few studies,



**Fig. 5.** (A) Representative immunoblots of SDS-PAGE separated proteins from thylakoid membranes probed with antibodies raised against PSI (PsaA and PsaB) and PSII (D1, PsbO, Lhcb1)-associated polypeptides and Rubisco (PbcL) in control (C, 18 μM Mn) and *Arabidopsis* plants treated with 500 and 1500 μM Mn. (B) The relative polypeptide abundance was quantified by densitometric analysis (area × intensity bands) of the immunoblots and the presented data were normalized to the relative abundance of PsaA, PsaB, D1, PsbO, Lhcb1, and PbcL in control *Arabidopsis* plants. Mean ± SE were calculated from three independent experiments. Different lower-case letters indicate significant differences between the Mn treatments ( $P < 0.05$ ).

the potential effect(s) of excess Mn on the functional/structural integrity of PSI remains elusive. As far as is known, the results presented in this study are the first report of an *in vivo* assessment of high Mn concentrations on PSI photochemistry. In contrast to PSII photochemistry, *in vivo* measurements of the oxidation state of P700 (P700<sup>+</sup>) (Klughammer and Schreiber, 1991; Ivanov et al., 1998), the primary donor of PSI demonstrated that the relative amount of oxidizable P700 (P700<sup>+</sup>) decreased by 30% in Mn-treated *Arabidopsis* plants at concentrations above 1000 μM (Fig. 4 and Table 5). The functional impairment of PSI photochemistry by excess Mn was

accompanied by a significant reduction in the abundance of PSI reaction centre polypeptides (PsaA and PsaB, Fig. 5). This clearly indicates that the major target of Mn toxicity within the photosynthetic electron transport chain of *Arabidopsis* is PSI- rather than PSII-related components. The reduced amounts of PSI reaction centre polypeptides PsaA and PsaB would imply acceptor side limitations of the photosynthetic electron transport and this could explain the increased reduction state Q<sub>A</sub> in Mn-stressed plants reported earlier (Kitao et al., 1997b). Moreover, the decreased expression of another Fe-containing chloroplastic protein precursor, ferredoxin-1

serving as a terminal electron acceptor of the photosynthetic electron transport, observed in Mn-treated young rice leaves also supports Mn-induced limitations at the acceptor side of PSI (Fühns *et al.*, 2010).

One of the major mechanisms considered for Mn toxicity involves the inhibition of other essential cations including Fe, thus suggesting that a Mn-induced Fe deficiency may play a key role in the physiological responses to excess Mn (Foy *et al.*, 1978; Foy, 1984; Kohno *et al.*, 1984). More recently, chloroplast alterations in maize plants exposed to excess Mn (Doncheva *et al.*, 2009) and Mn toxicity in young rice leaves (Fühns *et al.*, 2010) have been also ascribed to Mn-induced Fe deficiency rather than to direct Mn-induced oxidative stress. Given that about 80% of the plant Fe is located in the chloroplast (Terry and Abadia, 1986) and that the functional photosynthetic apparatus requires 22–23 iron atoms, of which PSI is the most Fe-abundant component (Ferreira and Straus, 1994), it seems reasonable to assume that the observed lower abundance of PSI reaction centre polypeptides and the associated decline of PSI photochemistry in Mn-treated *Arabidopsis* plants were consequences of a Mn-induced moderate Fe deficiency.

Although light energy is important for photosynthetic processes in plants, an excess of light can be also harmful because it can result in photoinhibition, which can be exacerbated when it is combined with other stresses (Powles, 1984; Aro *et al.*, 1993; Sonoike, 1996). Photoinhibition is a complex phenomenon that may cause damage to the photosynthetic apparatus reducing the photosynthetic efficiency when light conditions exceed the photon requirements for photosynthesis (Murata *et al.*, 2007). It is considered that PSII is the main site of photoinhibition (Aro *et al.*, 1993; Sonoike, 1996; Takahashi and Murata, 2008), being more unstable than PSI, because the D1 protein, one of the two major heterodimeric polypeptides of the PSII reaction centre complex, has a very high light-dependent turnover rate (Aro *et al.*, 1993; Burnap, 2004; Scheller and Haldrup, 2005; Takahashi and Murata, 2008). Earlier reports have shown that the susceptibility to Mn toxicity is strongly dependent on the light intensity and exposure of Mn-treated plants to high light can exacerbate the toxic effect of Mn (Horiguchi, 1988; Nable *et al.*, 1988; González *et al.*, 1998; Clair and Lynch, 2004; Hajiboland and Hasani, 2007). The present results also demonstrate that *Arabidopsis* plants predisposed to high Mn concentrations are more susceptible to photoinhibitory damage of PSII photochemistry in a concentration-dependent manner (Table 4).

Apart from the radiation-less dissipation of excess excitation energy in the chlorophyll pigment bed of LHCII, associated with the formation of the xanthophylls pigment zeaxanthin, which is considered one of the major protective mechanisms against photoinhibitory damage (Horton *et al.*, 1996; Niyogi, 1999), PSI-dependent cyclic electron transport has been also suggested to play a significant role in preventing the photoinhibitory damage of the photosynthetic apparatus during exposure of plants to high light conditions (Munekage *et al.*, 2002; Takahashi *et al.*, 2009). Considering the increased re-reduction rate of P700<sup>+</sup> in *Arabidopsis* (Table 5), the higher susceptibility of plants exposed to excess

Mn to photoinhibition (Table 4) might be due to lower capacity of PSI-driven cyclic electron flow under conditions of Mn toxicity.

In summary, the results presented in this research demonstrate for the first time that exposure of *Arabidopsis* plants to excess Mn causes specific negative effects on the abundance of polypeptides comprising the reaction centre of PSI, thus resulting in decreased PSI photochemistry and lower capacity for cyclic electron transport, which may be due to a Mn-induced Fe deficiency and may have critical physiological implications under conditions of Mn toxicity in higher plants.

## Acknowledgements

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NPAH), a CONICYT internship ‘Pasantías Doctorales en el Extranjero 2010’, the Department of Biology and the Biotron, University of Western Ontario, London, Ontario, Canada, and the Fondecyt project, Government of Chile (no. 1080372).

## References

- Alam S, Kodama R, Akiha F, Kamei S, Kawai S. 2006. Alleviation of manganese phytotoxicity in barley with calcium. *Journal of Plant Nutrition* **29**, 59–74.
- Arnon DI. 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* **24**, 1–15.
- Aro E-M, Virgin I, Andersson B. 1993. Photoinhibition of photosystem II. Inactivation, protein damage and turnover. *Biochimica et Biophysica Acta* **1143**, 113–134.
- Asada K, Heber U, Schreiber U. 1993. Electron flow to the intersystem chain from stromal components and cyclic electron flow in maize chloroplasts, as determined in intact leaves by monitoring redox change of P700 and chlorophyll fluorescence. *Plant and Cell Physiology* **34**, 39–50.
- Bowler C, VanCamp W, VanMontagu M, Inze D. 1994. Superoxide dismutase in plants. *Critical Reviews in Plant Sciences* **13**, 199–218.
- Bradi H. 2004. Adsorption of heavy metal ions on soils and soils constituents. *Journal of Colloids and Interface Science* **277**, 1–18.
- Burnap RL. 2004. D1 protein processing and Mn cluster assembly in light of the emerging Photosystem II structure. *Physical Chemistry Chemical Physics* **6**, 4803–4809.
- Burnell JN. 1988. The biochemistry of manganese in plants. In: RD Graham, J Hannam, NC Uren, eds, *Manganese in soils and plants*. Dordrecht, The Netherlands: Kluwer Academic Publishers, pp 125–133.
- Cailliatte R, Schikora A, Briat JF, Mari S, Curie C. 2010. High-affinity manganese uptake by the metal transporter NRAMP1 is essential for *Arabidopsis* growth in low manganese conditions. *The Plant Cell* **22**, 904–917.
- Chatterjee C, Nautiyal N, Agarwala SC. 1994. Influence of changes in manganese and magnesium supply on some aspects of wheat physiology. *Soil Science and Plant Nutrition* **40**, 191–197.

- Clair SB St, Lynch JP.** 2004. Photosynthetic and antioxidant enzyme responses of sugar maple and red maple seedlings to excess manganese in contrasting light environments. *Functional Plant Biology* **31**, 1005–1014.
- Delhaize E, Gruber BD, Pittman JK, White RG, Leung H, Miao Y, Jiang L, Ryan PR, Richardson AE.** 2007. A role for the AtMTP11 gene of *Arabidopsis* in manganese transport and tolerance. *The Plant Journal* **51**, 198–210.
- Demirevska-Kepova K, Simova-Stoilova L, Stoyanova Z, Holzer R, Feller U.** 2004. Biochemical changes in barley plants after excessive supply of copper and manganese. *Environmental and Experimental Botany* **52**, 253–266.
- de Vault D, Govindjee.** 1990. Photosynthetic glow peaks and their relationship with the free energy changes. *Photosynthesis Research* **24**, 175–181.
- Doncheva S, Poschenrieder C, Stoyanova Z, Georgieva K, Velichkova M, Barceló J.** 2009. Silicon amelioration of manganese toxicity in Mn-sensitive and Mn-tolerant maize varieties. *Environmental and Experimental Botany* **65**, 189–197.
- Enami I, Okumura A, Nagao R, Suzuki T, Iwai M, Shen JR.** 2008. Structures and functions of the extrinsic proteins of photosystem II from different species. *Photosynthesis Research* **98**, 349–363.
- Fecht-Christoffers MM, Maier P, Horst W.** 2003. Apoplastic peroxidase and ascorbate are involved in manganese toxicity and tolerance of *Vigna unguiculata*. *Physiologia Plantarum* **117**, 237–244.
- Feng J-P, Shi Q-H, Wang X-F.** 2009. Effects of exogenous silicon on photosynthetic capacity and antioxidant enzyme activities in chloroplast of cucumber seedlings under excess manganese. *Agricultural Science in China* **8**, 40–50.
- Fernando DR, Baker AJM, Woodrow IE.** 2009. Physiological responses in *Macadamia integrifolia* on exposure to manganese treatment. *Australian Journal of Botany* **57**, 406–413.
- Ferreira F, Straus NA.** 1994. Iron deprivation in cyanobacteria. *Journal of Applied Phycology* **6**, 199–210.
- Ferreira KN, Iverson TM, Maghlaoui K, Barber J, Iwata S.** 2004. Architecture of the photosynthetic oxygen-evolving center. *Science* **303**, 1831–1838.
- Foy CD.** 1984. Physiological effects of hydrogen, aluminum, and manganese toxicities in acid soils. In: F Adams, editor, *Soil acidity and liming*, 2nd ed. Madison: American Society of Agronomy. pp 57–97.
- Foy CD, Chaney RL, White MC.** 1978. The physiology of metal toxicity in plants. *Annual Review of Plant Physiology* **29**, 511–566.
- Führs H, Behrens C, Gallien S, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ.** 2010. Physiological and proteomic characterization of manganese sensitivity and tolerance in rice (*Oryza sativa*) in comparison with barley (*Hordeum vulgare*). *Annals of Botany* **105**, 1129–1140.
- Führs H, Hartwig M, Molina LEB, Heintz D, Van Dorsselaer A, Braun HP, Horst WJ.** 2008. Early manganese-toxicity response in *Vigna unguiculata* L. – a proteomic and transcriptomic study. *Proteomics* **8**, 149–159.
- Genty B, Briantais J-M, Baker NR.** 1989. The relationship between quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica et Biophysica Acta* **990**, 87–92.
- González A, Lynch JP.** 1997. Effects of manganese toxicity on leaf CO<sub>2</sub> assimilation of contrasting common bean genotypes. *Physiologia Plantarum* **101**, 872–880.
- González A, Lynch JP.** 1999. Subcellular and tissue Mn compartmentation in bean leaves under Mn toxicity stress. *Australian Journal of Plant Physiology* **26**, 811–822.
- González A, Steffen KL, Lynch JP.** 1998. Light and excess manganese. Implication for oxidative stress in common bean. *Plant Physiology* **18**, 493–504.
- Hajiboland R, Hasani BD.** 2007. Effect of Cu and Mn toxicity on chlorophyll fluorescence and gas exchange in rice and sunflower under different light intensities. *Journal of Stress Physiology and Biochemistry* **3**, 4–17.
- Hankamer B, Barber J, Boekema EJ.** 1997. Structure and membrane organization of photosystem II in green plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **48**, 641–671.
- Horiguchi T.** 1988. Mechanism of manganese toxicity and tolerance of plants. VII. Effect of light intensity on manganese-induced chlorosis. *Journal of Plant Nutrition* **11**, 235–245.
- Horton P, Ruban AV, Walters RG.** 1996. Regulation of light harvesting in green plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **47**, 655–684.
- Houtz RL, Nable RO, Cheniae GM.** 1988. Evidence for effects on the *in vivo* activity of ribulose-bisphosphate carboxylase/oxygenase during development of Mn toxicity in tobacco. *Plant Physiology* **86**, 1143–1149.
- Ivanov AG, Hendrickson L, Krol M, Selstam E, Öquist G, Hurry V, Hüner NPA.** 2006a. Digalactosyl-diacylglycerol deficiency impairs the capacity for photosynthetic intersystem electron transport and state transitions in *Arabidopsis thaliana* due to photosystem I acceptor-side limitations. *Plant and Cell Physiology* **47**, 1146–1157.
- Ivanov AG, Morgan R, Gray GR, Velitchkova MY, Hüner NPA.** 1998. Temperature/light dependent development of selective resistance to photoinhibition of photosystem I. *FEBS Letters* **430**, 288–292.
- Ivanov AG, Sane P, Hurry V, Krol M, Sveshnikov D, Hüner NPA, Öquist G.** 2003. Low-temperature modulation of the redox properties of the acceptor side of photosystem II: photoprotection through reaction centre quenching of excess energy. *Physiologia Plantarum* **119**, 376–383.
- Ivanov AG, Sane PV, Krol M, Gray GR, Balseris A, Savitch LV, Öquist G, Hüner NPA.** 2006b. Acclimation to temperature and irradiance modulates PSII charge recombination. *FEBS Letters* **580**, 2797–2802.
- Ivanov AG, Sane PV, Zeinalov Y, Malmberg G, Gardeström P, Hüner NPA, Öquist G.** 2001. Photosynthetic electron transport adjustments in overwintering Scots pine (*Pinus sylvestris* L.). *Planta* **213**, 575–585.
- Ivanov AG, Sane PV, Zeinalov Y, Simidjiev I, Hüner NPA, Öquist G.** 2002. Seasonal responses of photosynthetic electron transport in Scots pine (*Pinus sylvestris* L.) studied by thermoluminescence. *Planta* **215**, 457–465.
- Jordan DB, Ogren WL.** 1983. Species variation in kinetic properties of Rubisco. *Archives of Biochemistry and Biophysics* **227**, 425–433.

- Kern J, Renger G.** 2007. Photosystem II: Structure and mechanism of the water:plastoquinone oxidoreductase. *Photosynthesis Research* **94**, 183–202.
- Khabaz-Saberi H, Rengel Z, Wilson R, Setter TL.** 2010. Variation of tolerance to manganese toxicity in Australian hexaploid wheat. *Journal of Plant Nutrition and Soil Science* **173**, 103–112.
- Kitao M, Lei TT, Koike T.** 1997a. Comparison of photosynthetic responses to manganese toxicity of deciduous broad leaved trees in northern Japan. *Environmental Pollution* **97**, 113–118.
- Kitao M, Lei TT, Koike T.** 1997b. Effects of manganese toxicity on photosynthesis of white birch (*Betula platyphylla* var. *japonica*) seedlings. *Physiologia Plantarum* **101**, 249–256.
- Klughammer C, Schreiber U.** 1991. Analysis of light-induced absorbancy changes in the near-infrared spectral region. 1. Characterization of various components in isolated chloroplasts. *Zeitschrift für Naturforschung C* **46**, 233–244.
- Kohno Y, Foy CD, Fleming AL, Krizek DT.** 1984. Effect of Mn concentration on the growth and distribution of Mn and Fe in two bush bean cultivars grown in solution culture. *Journal of Plant Nutrition* **7**, 547–566.
- Kramer DM, Johnson G, Kuirats O, Edwards GE.** 2004. New fluorescence parameters for the determination of QA redox state and excitation energy fluxes. *Photosynthesis Research* **79**, 209–218.
- Krol M, Ivanov AG, Jansson S, Kloppstech K, Hüner NPA.** 1999. Greening under high light and cold temperature affects the level of xanthophyll-cycle pigments, early light-inducible proteins, and light harvesting polypeptides in wild-type barley and the Chlorina *f2* mutant. *Plant Physiology* **120**, 193–203.
- Laemmli U.** 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Lei Y, Korpelainen H, Li C.** 2007. Physiological and biochemical responses to high Mn concentrations in two contrasting *Populus cathayana* populations. *Chemosphere* **68**, 686–694.
- Li Q, Chen L-S, Jiang H-X, Tang N, Yang L-T, Lin Z-H, Li Y, Yang G-H.** 2010. Effects of manganese-excess on CO<sub>2</sub> assimilation ribulose-1,5-bisphosphate carboxylase/oxygenase, carbohydrates and photosynthetic electron transport of leaves, and antioxidant systems of leaves and roots in *Citrus grandis* seedlings. *BMC Plant Biology* **10**, 42.
- Lidon FC, Barreiro MG, Ramalho JC.** 2004. Manganese accumulation in rice: implications for photosynthetic functioning. *Journal of Plant Physiology* **161**, 1235–1244.
- Macfie S, Taylor G.** 1992. The effects of excess manganese on photosynthetic rate and concentration of chlorophyll in *Triticum aestivum* grown in solution culture. *Physiologia Plantarum* **85**, 467–475.
- McDaniel KL, Toman FR.** 1994. Short-term effects of manganese toxicity on ribulose-1,5-bisphosphate carboxylase in tobacco chloroplasts. *Journal of Plant Nutrition* **17**, 523–536.
- Marschner H.** 1995. *Mineral nutrition of higher plants*. San Diego: Academic Press.
- Maxwell PC, Biggins J.** 1976. Role of cyclic electron transport in photosynthesis as measured by turnover of P700 *in vivo*. *Biochemistry* **15**, 3975–3981.
- Millaleo R, Reyes-Díaz M, Ivanov AG, Mora ML, Alberdi M.** 2010. Manganese as essential and toxic element for plants: Transport, accumulation and resistance mechanisms. *Journal of Soil Science and Plant Nutrition* **10**, 476–494.
- Minagawa J, Narusaka Y, Inoue Y, Satoh K.** 1999. Electron transfer between Q<sub>A</sub> and Q<sub>B</sub> in photosystem II is thermodynamically perturbed in phototolerant mutants of *Synechocystis* sp. PCC 6803. *Biochemistry* **38**, 770–775.
- Mora M, Rosas A, Ribera A, Rengel R.** 2009. Differential tolerance to Mn toxicity in perennial ryegrass genotypes: involvement of antioxidative enzymes and root exudation of carboxylates. *Plant and Soil* **320**, 79–89.
- Mukhopadhyay MJ, Sharma A.** 1991. Manganese in cell metabolism of higher plants. *Botanical Review* **57**, 117–149.
- Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T.** 2002. *PGR5* is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. *Cell* **110**, 361–371.
- Murata N, Takahashi S, Nishiyama Y, Allakhverdiev SI.** 2007. Photoinhibition of photosystem II under environmental stress. *Biochimica et Biophysica Acta* **1767**, 414–421.
- Nable RO, Houtz RL, Cheniae GM.** 1988. Early inhibition of photosynthesis during development of Mn toxicity in tobacco. *Plant Physiology* **86**, 1136–1142.
- Niyogi KK.** 1999. Photoprotection revisited: Genetic and molecular approaches. *Annual Review of Plant Physiology and Plant Molecular Biology* **50**, 333–359.
- Ohki K.** 1984. Manganese deficiency and toxicity effects on growth, development and nutrient composition in wheat. *Agronomy Journal* **76**, 213–218.
- Panda S, Mishra AK, Biswal UC.** 1987. Manganese induced peroxidation of thylakoid lipids and changes in chlorophyll- $\alpha$  fluorescence during aging of cell free chloroplasts in light. *Phytochemistry* **26**, 3217–3219.
- Papadakis IE, Giannakoula A, Therios IN, Bosabalidis AM, Moustakas M, Nastou A.** 2007. Mn-induced changes in leaf structure and chloroplast ultrastructure of *Citrus volkameriana* (L) plants. *Journal of Plant Physiology* **164**, 100–103.
- Powles SB.** 1984. Photoinhibition of photosynthesis induced by visible light. *Annual Review of Plant Physiology* **35**, 15–44.
- Ravenel J, Peltier G, Havaux M.** 1994. The cyclic electron pathways around photosystem I in *Chlamydomonas reinhardtii* as determined *in vivo* by photoacoustic measurements of energy storage. *Planta* **193**, 251–259.
- Rosas A, Rengel Z, Mora M.** 2007. Manganese supply and pH influence growth, carboxylate exudation and peroxidase activity of ryegrass and white clover. *Journal of Plant Nutrition* **30**, 253–270.
- Rosso D, Bode R, Li W, Krol M, Saccon D, Wang S, Schillaci LA, Rodermeil SR, Maxwell DP, Hüner NPA.** 2009. Photosynthetic redox imbalance governs leaf sectoring in the *Arabidopsis thaliana* variegation mutants *immutans*, *spotty*, *var1*, and *var2*. *The Plant Cell* **21**, 3473–3492.
- Rutherford AW, Boussac A.** 2004. Water photolysis in biology. *Science* **303**, 1782–1784.

- Sadzwaka A, Carrasco M, Grez R, Mora M.** 2004. *Métodos de análisis de tejidos vegetales. Comisión de normalización y acreditación.* Sociedad Chilena de la Ciencia del Suelo (SCCS). Available at: [http://www.inia.cl/platina/descarga/docs/libros/asadzwaka\\_tejveg\\_2004.pdf](http://www.inia.cl/platina/descarga/docs/libros/asadzwaka_tejveg_2004.pdf).
- Sane PV.** 2004. Thermoluminescence: a technique for probing photosystem II. In: R Carpentier, ed, *Photosynthesis research protocols*. Totowa, NJ: Humana Press. pp 229–248.
- Sane PV, Ivanov AG, Öquist G, Hüner NPA.** 2012. Thermoluminescence. In: JJ Eaton-Rye, BC Tripathy, TD Sharkey, eds, *Photosynthesis: plastid biology, energy conversion and carbon assimilation. Advances in photosynthesis and respiration*, vol 34. Dordrecht, The Netherlands: Springer Science+Business Media. pp 445–474.
- Schaaf G, Catoni E, Fitz M, Schwake R, Schneider A, von Wirén N, Frommer WB.** 2002. A putative role for vacuolar calcium/manganese proton antiporter *AtCAX2* in heavy metal detoxification. *Plant Biology* **2**, 612–618.
- Scheller H, Haldrup A.** 2005. Photoinhibition of photosystem I. *Planta* **221**, 5–8.
- Schreiber U, Bilger W, Neubauer C.** 1994. Chlorophyll fluorescence as a noninvasive indicator for rapid assessment of in vivo photosynthesis. In: ED Schulze, MM Caldwell, eds, *Ecophysiology of photosynthesis*. Berlin: Springer-Verlag. pp 49–70.
- Sinha S, Mukherji S, Dutta J.** 2002. Effect of manganese toxicity on pigment content, Hill activity and photosynthetic rate of *Vigna radiata* L. Wilczek seedlings. *Journal of Environmental Biology* **23**, 253–257.
- Sonoike K.** 1996. Photoinhibition of photosystem I: its physiological significance in the chilling sensitivity of plants, *Plant and Cell Physiology* **37**, 239–247.
- Stoyanova Z, Poschenrieder C, Tzvetkova N, Doncheva S.** 2009. Characterization of the tolerance to excess manganese in four maize varieties. *Soil Science and Plant Nutrition* **55**, 747–753.
- Subrahmanyam D, Rathore VS.** 2000. Influence of manganese toxicity on photosynthesis in ricebean (*Vigna umbellata*) seedlings. *Photosynthetica* **38**, 449–453.
- Takahashi S, Murata N.** 2008. How do environmental stresses accelerate photoinhibition? *Trends in Plant Science* **13**, 178–182.
- Takahashi S, Milward SE, Fan DY, Chow WS, Badger MR.** 2009. How does cyclic electron flow alleviate photoinhibition in *Arabidopsis*? *Plant Physiology* **149**, 1560–1567.
- Terry N, Abadia J.** 1986. Function of iron in chloroplasts. *Journal of Plant Nutrition* **9**, 609–646.
- Towbin H, Staehelin T, Gordon J.** 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences, USA* **76**, 4350–4354.
- van Kooten O, Snel JFH.** 1990. The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosynthesis Research* **25**, 147–150.
- Vass I, Govindjee.** 1996. Thermoluminescence from the photosynthetic apparatus. *Photosynthesis Research* **48**, 117–126.
- Vaughn KC, Duke SO.** 1984. Function of polyphenol oxidase in higher plants. *Physiologia Plantarum* **60**, 106–112.